

GEOMETRY AND ARITHMETIC OF INTEGRABLE HIERARCHIES OF KDV TYPE. I. INTEGRALITY

BORIS DUBROVIN[†], DI YANG, AND DON ZAGIER

ABSTRACT. For each of the simple Lie algebras $\mathfrak{g} = A_l, D_l$ or E_6 , we show that the all-genera one-point FJRW invariants of \mathfrak{g} -type, after multiplication by suitable products of Pochhammer symbols, are the coefficients of an algebraic generating function and hence are integral. Moreover, we find that the all-genera invariants themselves coincide with the coefficients of the unique calibration of the Frobenius manifold of \mathfrak{g} -type evaluated at a special point. For the A_4 (5-spin) case we also find two other normalizations of the sequence that are again integral and of at most exponential growth, and hence conjecturally are the Taylor coefficients of some period functions.

CONTENTS

1. Introduction and description of the results for the 5-spin case	1
2. Cohomological field theories and FJRW invariants of \mathfrak{g} -type	6
3. Differential equation	9
4. Computing residues	13
5. The A series	19
6. The D series	26
7. The E_6 case	32
8. Explicit relation for FJRW invariants of \mathfrak{g} -type	35
Appendix A. Wave functions	45
Appendix B. Frobenius manifolds	47
References	50

1. INTRODUCTION AND DESCRIPTION OF THE RESULTS FOR THE 5-SPIN CASE

In this paper, we will study certain intersection numbers $\tau_{\mathfrak{g}}(g)$ (the precise definition will be given in Section 2) on the moduli space of stable algebraic curves $\overline{\mathcal{M}}_{g,n}$ [32] associated to simple Lie algebras, the case of A_{r-1} -type simple Lie algebra being essentially one-point r -spin intersection numbers, introduced by Witten [112]. (What we call $\tau_{A_{r-1}}(g)$ would be $\langle \tau_{n,m} \rangle$ in Witten's notation, where $2(r+1)g = r(n+1) + m + 2$ with $n \geq 0$, $0 \leq m \leq r-2$; our notation in later sections will be slightly different from Witten's.) In particular, we will give recursive, closed, and asymptotic formulas for these numbers. Using these formulas, we will show for $\mathfrak{g} = A_l$ ($l \geq 1$), or D_l ($l \geq 4$), or E_6 that by multiplying $\tau_{\mathfrak{g}}(g)$ by appropriate gamma factors (products of Pochhammer symbols) we obtain new numbers whose generating functions are *algebraic*. In particular, these renormalized numbers are integral and grow only exponentially in g . Moreover, for the case of A_4 , we find that there are *different* normalizations of the $\tau_{\mathfrak{g}}(g)$, obtained by multiplying by other gamma factors, that are again integral and of exponential growth, so that each of the corresponding generating series is conjecturally a period function for

[†]Deceased on March 19, 2019.

some family of algebraic varieties (or equivalently, a solution of some Picard–Fuchs differential equation). This latter point is also of interest from the point of view of the general arithmetic theory of differential equations (see for instance [115], where the conjecture relating integrality and geometric origin is discussed on pp. 728–729 and the A_4 example on pp. 768–769) and will be discussed from this point of view in the later paper [114].

For most of this introduction, we will assume that $\mathfrak{g} = A_4$ and describe our results in detail only for that case, indicating briefly at the end of the introduction where the statements of the general results can be found in the paper. For convenience, we write $\tau_g = \tau_{A_4}(g)$ for $g > 0$, and also set $\tau_0 = 1$ and $\tau_g = 0$ for $g < 0$. The first values are

g	0	1	2	3	4	5	6	7	8	9
τ_g	1	$\frac{1}{6}$	$\frac{11}{3600}$	0	$\frac{341}{25920000}$	$\frac{161}{777600000}$	$\frac{3397}{93312000000}$	$\frac{3421}{4199040000000}$	0	$\frac{1670581}{84652646400000000}$

One-point 5-spin intersection numbers

The following theorem, which will be proved in Section 5, gives three different integrality statements about the numbers τ_g . One of these statements (the integrality of the numbers a_g) will be generalized to all A_l , D_l and E_6 in Sections 5–7 below. The two others will be given in this paper for the A_4 case only, with a discussion of the integrality properties of $\tau_{\mathfrak{g}}(g)$ for other simple Lie algebras postponed to the later paper [114] mentioned above.

Theorem 1. *Each of the renormalized values*

$$a_{5n} = 2^{2n} \left(\frac{2}{5}\right)_n \left(\frac{-1}{10}\right)_n \tau_{5n}, \quad b_{5n} = \left(\frac{4}{5}\right)_n \left(\frac{1}{5}\right)_n \tau_{5n}, \quad c_{5n} = \left(\frac{4}{5}\right)_n \left(\frac{3}{5}\right)_n \tau_{5n}, \quad (1)$$

$$a_{5n-1} = \frac{2^{2n}}{3} \left(\frac{1}{5}\right)_n \left(\frac{-3}{10}\right)_n \tau_{5n-1}, \quad b_{5n-1} = \left(\frac{2}{5}\right)_n \left(\frac{3}{5}\right)_n \tau_{5n-1}, \quad c_{5n-1} = \left(\frac{2}{5}\right)_n \left(\frac{4}{5}\right)_n \tau_{5n-1}, \quad (2)$$

$$a_{5n-3} = \frac{5 \cdot 2^{2n}}{14} \left(\frac{-1}{5}\right)_n \left(\frac{-7}{10}\right)_n \tau_{5n-3}, \quad b_{5n-3} = \left(\frac{3}{5}\right)_n \left(\frac{2}{5}\right)_n \tau_{5n-3}, \quad c_{5n-3} = \left(\frac{3}{5}\right)_n \left(\frac{1}{5}\right)_n \tau_{5n-3}, \quad (3)$$

$$a_{5n-4} = \frac{5 \cdot 2^{2n}}{36} \left(\frac{-2}{5}\right)_n \left(\frac{-9}{10}\right)_n \tau_{5n-4}, \quad b_{5n-4} = \left(\frac{1}{5}\right)_n \left(\frac{4}{5}\right)_n \tau_{5n-4}, \quad c_{5n-4} = \left(\frac{1}{5}\right)_n \left(\frac{2}{5}\right)_n \tau_{5n-4} \quad (4)$$

belongs to $\mathbb{Z}\left[\frac{1}{30}\right]$. Here $(x)_k := x(x+1)\cdots(x+k-1)$ denotes the ascending Pochhammer symbol.

The point here is that the numbers τ_g decay like $1/\Gamma(\frac{2g}{5})$, as we will see in a moment, and that therefore each of the numbers a_g, b_g and c_g , as well as being integral (away from the primes 2, 3 and 5), is of only exponential growth in g and hence is expected to be the g th Taylor coefficient of some period function. It would be very interesting to identify the two generating functions $\sum b_g x^g$ and $\sum c_g x^g$ as explicit period functions, but we have not yet been able to do this. For a_g , on the other hand, the next theorem includes the stronger statement that the generating function $\sum a_g x^g$ is not only a period function, but is in fact algebraic.

Remark 1. The formulas (1)–(4) can be written uniformly as

$$a_g = \frac{(-1)^m}{5} (A)_m \tau_g, \quad b_g = (A)_n (B)_n \tau_g, \quad c_g = (A)_n (C)_n \tau_g, \quad (5)$$

where $m = \lfloor \frac{2g-1}{5} \rfloor$, $n = \lfloor \frac{g}{5} \rfloor$, and A, B, C are the fractional parts of $\frac{2g-1}{5}$, $\frac{2g+1}{5}$, $\frac{2g+3}{5}$, respectively. Recall that for a real number x , its fractional part is defined as $\{x\} := x - [x]$. Note that $A = 0$ if $g \equiv 3 \pmod{5}$, and that both a_g and τ_g vanish in this case.

In the following theorem we collect many further properties of τ_g . All of the statements of this theorem will be generalized in the main body of the paper to the A_l, D_l , and E_6 cases.

Theorem 2. *The numbers $\tau_g = \tau_{A_4}(g)$ have the following properties:*

(i) [recursion] *The numbers τ_g satisfy the recursion relation*

$$\begin{aligned} & 2^8 3^4 5^{17} 31 g (g-1) (g-2) (g-4) \tau_g \\ & - 5^{11} (2^8 3^4 g^4 - 2^{13} 3^4 g^3 + 2^4 3^2 54331 g^2 - 2^4 3^2 43^1 6329 g + 5^1 7^1 2013229) \tau_{g-5} \\ & + 2^2 5^6 (2^2 3^2 5 g^2 - 2^2 3^3 5^1 7 g + 19739) \tau_{g-10} - \tau_{g-15} = 0, \quad g \in \mathbb{Z} \end{aligned} \quad (6)$$

with the initial conditions given by the above table.

(ii) [dual topological ODE] *The generating series*

$$\varphi(X) := \sum_{g \geq 0} \tau_g X^{\frac{g}{5}} \quad (7)$$

satisfies the fourth-order linear differential equation $Q\varphi = 0$, where

$$\begin{aligned} Q = & 2^8 3^4 5^{15} X^3 (X - 5^6 31) \frac{d^4}{dX^4} + 2^8 3^4 5^{14} X^2 (2^1 3^2 X - 5^6 23^1 31) \frac{d^3}{dX^3} \\ & - 2^4 3^2 5^9 X (X^2 - 5^4 6091 X + 2^6 3^3 5^{10} 7^1 31) \frac{d^2}{dX^2} \\ & - 2^5 3^2 5^8 (2 X^2 - 5^4 3209 X + 2^5 3^3 5^{10} 31) \frac{d}{dX} \\ & + (X^2 + 2^2 5^6 61 X + 5^{13} 7^1 23^1 31). \end{aligned} \quad (8)$$

(iii) [algebraicity] *The generating function of the numbers a_g defined in (5) is algebraic. More precisely, we have*

$$y := \sum_{g \geq 0} a_g z^g \quad \Rightarrow \quad y^5 - \frac{z}{6} y^3 + \frac{z^2}{400} y = 1. \quad (9)$$

(iv) [closed formula] *Denote $m = \lfloor (2g-1)/5 \rfloor$ as above, and define $c_{p,j} \in \mathbb{Q}$ ($0 \leq p \leq j$) by*

$$c_{p,j} := \text{coefficient of } x^j \text{ in } \frac{1}{p!} \left(\frac{(1+x)^6 - 1 - 6x}{6x} \right)^p. \quad (10)$$

Then for all $g \geq 0$ with $g \not\equiv 3 \pmod{5}$, we have

$$\tau_g = \Gamma\left(\left\{\frac{2g-1}{5}\right\}\right) \frac{(-1)^{g+m-1}}{5g} \sum_{p=0}^{2g} \frac{c_{p,2g}}{\Gamma\left(\frac{2g-1}{5} - p + 1\right)}. \quad (11)$$

(v) [product formula] *Let*

$$w(u) = 1 + \sum_{n \geq 0} C_n u^{n+1} = 1 + u - \frac{2}{3} u^2 + \frac{11}{18} u^3 - \dots \quad (12)$$

be the unique power-series solution in $1 + u + u^2 \mathbb{Q}[[u]]$ to the sextic equation

$$\frac{w^6}{30} - \frac{w}{5} + \frac{1}{6} = \frac{u^2}{2}, \quad (13)$$

and let $f(T) := \sum_{k \geq 0} (2k-1)!! C_{2k} (-T)^k$. Then

$$f(T) f(-T) = \sum_{g \geq 0} \left(1 + \frac{2g-1}{5}\right)_{2g} (-1)^{g-1} 5^{3g} a_g T^{2g} \quad (14)$$

with a_g as in (5) or in part (iii).

(vi) [terminating hypergeometric sum] For all $g \in \mathbb{Z}$ with $g \not\equiv 3 \pmod{5}$, we have

$$\tau_g = \frac{6^{-g}}{\Gamma(1 - \{\frac{2g-1}{5}\})} \sum_{0 \leq s \leq g/2} (-1)^s \left(\frac{3}{10}\right)^{2s} \frac{\Gamma(\frac{1+3g-5s}{5})}{s!(g-2s)!}. \quad (15)$$

(vii) [asymptotics] For $g \not\equiv 3 \pmod{5}$ and as $g \rightarrow \infty$, τ_g is given asymptotically by

$$\tau_g \sim \frac{5}{\sqrt{6^{\frac{3}{5}} \sqrt{\pi g^3}}} \frac{\sin(\{\frac{2g-1}{5}\})}{(\{\frac{2g-1}{5}\})_m} \left(6^{\frac{2}{5}} 20 \sin^2\left(\frac{\pi}{5}\right)\right)^{-g}, \quad (16)$$

where $m = [(2g-1)/5]$ is again as above.

The proof of this theorem is given in Sections 2 and 5.

Remark 2. Of course, parts (i) and (ii) are equivalent, by the usual bijection between power series satisfying a linear differential equation with polynomial coefficients and sequences of numbers satisfying a recursion with polynomial coefficients. More explicitly, since the recursion (6) involves only $\tau_g, \tau_{g-5}, \tau_{g-10}$ and τ_{g-15} , it is equivalent to five separate recursions for τ_g with $g \equiv s \pmod{5}$, $0 \leq s \leq 4$, the values for $s = 3$ being uninteresting because $\tau_g = 0$. Similarly, by the Frobenius method, the differential equation in part (ii) has four fundamental solutions in $x^\lambda \mathbb{Q}[[x]]$ with $\lambda = 0, \frac{1}{5}, \frac{2}{5}, \frac{4}{5}$, and the generating function (7) is simply a certain linear combination of these with rational coefficients.

Remark 3. The integrality (away from 2, 3, and 5) of a_g in Theorem 1 follows immediately from part (iii) of Theorem 2, since the Taylor coefficients of any algebraic function are integral after some fixed rescaling. The integrality of b_g and c_g is harder and will follow from formula (15), as a consequence of the stronger statement that the product of either $(A)_n(B)_n$ or $(A)_n(C)_n$ with each summand on the right-hand side of (15) is p -adically integral for each $p > 5$.

Remark 4. We say something here about the origin of the problem and about some of the various approaches that can be used to solve it. The way that we are approaching intersection numbers is to use integrable systems. In 1990, Witten [110] proposed his famous conjecture that the partition function of ψ -class intersection numbers is a tau-function for the Korteweg-de Vries (KdV) integrable hierarchy. This conjecture was later proved by Kontsevich [82]. The r -spin Witten conjecture [112], stating that the partition function of the r -spin intersection numbers gives a tau-function for the Gelfand–Dickey integrable hierarchy [34], was proved by Faber–Shadrin–Zvonkine [64]. (The $r = 2$ case corresponds to Witten’s conjecture in 1990.) More generally, the ADE Witten conjecture [112], given its precise form by Fan–Jarvis–Ruan [67] (cf. also Givental–Milanov [72]), states that the partition function of the FJRW invariants associated to a certain simple singularity [3] gives a tau-function for the Drinfeld–Sokolov (DS) integrable hierarchy [13, 25, 31, 38, 49]. The ADE Witten conjecture was proved by Fan–Jarvis–Ruan [67] (cf. also [68, 72, 89]) with the D_4 case confirmed in [65]. For all these cases, the main mathematical object of the study is the so-called *topological solution* u^{top} [48, 57] to the corresponding integrable system together with its tau-function. Perhaps, the simplest way to describe this particular solution is to use its initial value (observe that each member of the DS hierarchy is an evolutionary PDE), which in terms of the normal coordinates [57] r_1, \dots, r_l (where l is the rank) reads:

$$r_\alpha^{\text{top}}|_{\text{higher times}=0} = \eta_{\alpha 1} t^{1,0}, \quad (17)$$

where $\alpha = 1, \dots, l$ and $\eta_{\alpha 1}$ are constants. One could then apply methods in integrable systems to compute u^{top} and more importantly its *tau-function*. These methods include the wave-function approach [5, 8, 9, 14, 20, 25, 34, 39, 53, 77, 106, 107], the Ψ DOs [29, 34, 79, 86], the Sato

Grassmannian approach [14, 30, 34, 106, 107], the Dubrovin–Zhang approach [24, 43, 48, 50, 55, 57], the Givental quantization [24, 69, 70, 71, 72], the topological recursion of Chekhov–Eynard–Orantin type (or of Bouchard–Eynard type) (cf. [15, 59, 116, 118] and the references therein), the matrix-resolvent method [11, 12, 13, 52, 53] (cf. also [21, 87, 88, 96]), etc. One of the keys for several of these methods is the *Lax pair* (cf. [5, 34, 38, 39]). In the second paper [54] of the series, the mathematical object will be a different solution to the DS hierarchy, whose tau-function has a different topological meaning; more general cases are considered in joint papers currently in preparation with Daniele Valeri. Taking the dispersionless limit of the DS hierarchy, one can also obtain an interesting geometric structure, suitable for studying *arbitrary* solutions, i.e., the Frobenius structure [41, 42, 43, 48, 57], which plays one of the central roles in understanding the above-mentioned methods. Besides the methods from integrable systems, there are other important methods for computing the intersection numbers from other theories, including the theory of matrix models [1, 2, 16, 17, 18, 33, 73, 82, 94], vertex algebras [6, 35, 68, 78, 91, 117, 120], emergent geometry [118, 119, 121], etc.

Our next result (see Theorem 8 and its corollary) gives an explicit relationship between the all-genera FJRW intersection numbers of \mathfrak{g} -type (with $g = A_l, D_l$ or E_6) and genus zero, which again will be stated in the introduction only for the 5-spin case. Let B be the Frobenius manifold associated to the A_4 Coxeter group [43, 45, 104, 122], and let $(\theta_{\alpha,m})_{\alpha=1,\dots,4,m \geq 0}$ be the unique *calibration* [50, 57] on B . The reader who is not familiar with the theory of Frobenius manifolds could simply identify $\theta_{\alpha,m}$ with the following formal power series of infinitely many variables:

$$\theta_{\alpha,m} = \frac{\partial^2 \mathcal{F}_0(\mathbf{t})}{\partial t^{\alpha,m} \partial t^{1,0}}, \quad (18)$$

where $\mathbf{t} = (t^{\alpha,q})_{\alpha=1,\dots,4,q \geq 0}$ and the definition of \mathcal{F}_0 can be found in (37). Denote $v_\alpha = \theta_{\alpha,0}$, $\alpha = 1, \dots, 4$, and denote $v = (v_1, \dots, v_4)$. Then from the theory of Frobenius manifolds (see Appendix B) we know that all the $\theta_{\alpha,m}$ are *polynomials in v* , with the first values being

$$\begin{aligned} \theta_{\alpha,0} &= v_\alpha, \\ \theta_{1,1} &= v_1 v_4 + v_2 v_3, \\ \theta_{2,1} &= \frac{v_3^2}{2} + \frac{1}{10} v_1^2 v_3 + v_4 v_2 + \frac{1}{10} v_2^2 v_1, \\ \theta_{3,1} &= \frac{v_2^3}{15} + \frac{1}{75} v_1^3 v_2 + \frac{1}{5} v_3 v_1 v_2 + v_4 v_3, \\ \theta_{4,1} &= \frac{v_1^5}{2500} + \frac{1}{50} v_2^2 v_1^2 + \frac{1}{10} v_3^2 v_1 + \frac{v_4^2}{2} + \frac{1}{10} v_3 v_2^2, \\ \theta_{1,2} &= \frac{v_1^6}{3750} + \frac{1}{50} v_2^2 v_1^3 + \frac{1}{10} v_3^2 v_1^2 + \frac{1}{2} v_4^2 v_1 + \frac{1}{5} v_3 v_2^2 v_3 + \frac{v_2^4}{30} + \frac{v_3^3}{6} + v_4 v_3 v_2. \end{aligned}$$

Theorem 3. *Define*

$$v_1^* = \frac{1}{6}, \quad v_2^* = 0, \quad v_3^* = \frac{11}{3600}, \quad v_4^* = 0.$$

Then for all $g \geq 1$ with $g \not\equiv 3 \pmod{5}$, we have

$$\tau_g = \theta_{\alpha,m}(v^*), \quad (19)$$

where $\alpha \in \{1, 2, 3, 4\}$ and $m \geq 0$ are such that $2g - 1 = \alpha + 5m$.

Organization of the paper. In Section 2, we recall the definition of $\tau_{\mathfrak{g}}(g)$ for all \mathfrak{g} . In Section 3 we generalize parts (i), (ii) of Theorem 2 from A_4 to an arbitrary \mathfrak{g} . In Section 4, we provide several technical preparations for the subsequent sections. In Section 5 we prove the

generalization of the rest of Theorem 2 for the A series, as well as proving Theorem 1. The analogues of Theorem 2 for the D_l and E_6 cases are given in Sections 6 and 7, respectively, while Theorem 3 is generalized to the A_l , D_l and E_6 cases in Section 8. The necessary material on the wave-function-pair approach for computing residues of pseudodifferential operators and a brief review of the theory of Frobenius manifolds are provided in Appendices A and B.

Acknowledgements. We would like to thank Giordano Cotti, Yuri Manin, Maxim Smirnov, Jian Zhou and Wadim Zudilin for their interest and helpful discussions. D.Y. is grateful to Youjin Zhang and Marco Bertola for their advice. We also thank Elba Garcia-Failde for a very careful reading of this paper and for very helpful comments. Part of the work of D.Y. was done during his postdoctoral studies and visits at SISSA and MPIM; he thanks both institutes for their excellent working conditions, warm hospitality and financial support. The work was partially supported by the National Key Research and Development Project ‘‘Analysis and Geometry on Bundles’’ SQ2020YFA070080, and by NSFC No. 12061131014.

2. COHOMOLOGICAL FIELD THEORIES AND FJRW INVARIANTS OF \mathfrak{g} -TYPE

Let \mathfrak{g} be a simply-laced simple Lie algebra. In this section, we review the definitions of the enumerative invariants $\tau_{\mathfrak{g}}(g)$, called the one-point FJRW (Fan–Jarvis–Ruan, Witten) invariants, that are studied in this paper.

We start with the definitions for the A case. For $g, n \geq 0$, let $1 \leq \alpha_1, \dots, \alpha_n \leq r$ be integers satisfying the divisibility condition $r \mid \sum_{i=1}^n \alpha_i - n - (2g - 2)$. For an algebraic curve C of genus g with n distinct marked points x_1, \dots, x_n , there exists a line bundle \mathcal{T} on C such that

$$\mathcal{T}^{\otimes r} = K_C \otimes \mathcal{O}((1 - \alpha_1)x_1) \otimes \cdots \otimes \mathcal{O}((1 - \alpha_n)x_n), \quad (20)$$

where K_C is the canonical class of C . For C smooth there are r^{2g} such line bundles. A choice of such an ‘‘ r -th root’’ of the bundle (20) is called an r -spin structure, and it defines a point in a covering of $\overline{\mathcal{M}}_{g,n}$. After a suitable compactification, this covering is denoted by

$$p : \overline{\mathcal{M}}_{g,n}^{1/r}(\alpha_1, \dots, \alpha_n) \rightarrow \overline{\mathcal{M}}_{g,n}.$$

In genus zero, for a point $(C, x_1, \dots, x_n, \mathcal{T})$ in the covering space $\overline{\mathcal{M}}_{0,n}^{1/r}(\alpha_1, \dots, \alpha_n)$, denote $V = H^1(C, \mathcal{T})$. This gives a vector bundle $\mathcal{V} \rightarrow \overline{\mathcal{M}}_{0,n}^{1/r}(\alpha_1, \dots, \alpha_n)$ as the space V has constant dimension thanks to the fact that $H^0(C, \mathcal{T})$ vanishes. Put

$$c_0(\alpha_1, \dots, \alpha_n) := p_* e(\mathcal{V}^\vee) \in H^{2(s-1)}(\overline{\mathcal{M}}_{0,n}), \quad (21)$$

where $e(\mathcal{V}^\vee)$ is the Euler class of the dual vector bundle \mathcal{V}^\vee , and $s := \frac{\sum_{i=1}^n \alpha_i - n + 2}{r}$. The cohomology class $c_0(\alpha_1, \dots, \alpha_n)$ is called the *Witten class*. For higher genus, $H^0(C, \mathcal{T})$ is only generically zero and the vector bundle can only be defined on a generic stratum. The Witten class $c_g(\alpha_1, \dots, \alpha_n)$ could still be defined as a particular cohomology class in $H^{2(s+g-1)}(\overline{\mathcal{M}}_{g,n})$ with $s = \frac{\sum_{i=1}^n \alpha_i - n - (2g-2)}{r}$, but the construction is more involved (cf. [26, 27, 28, 66, 67, 75, 95, 100, 101, 112]). The genus g r -spin intersection numbers are defined as the following integrals:

$$\int_{\overline{\mathcal{M}}_{g,n}} c_g(\alpha_1, \dots, \alpha_n) \psi_1^{q_1} \cdots \psi_n^{q_n} =: \langle \tau_{\alpha_1, q_1} \cdots \tau_{\alpha_n, q_n} \rangle_g, \quad q_1, \dots, q_n \geq 0, \quad (22)$$

where ψ_i ($1 \leq i \leq n$) denotes the first Chern class of the i th tautological line bundle over $\overline{\mathcal{M}}_{g,n}$. These integrals vanish unless the degree and the dimension match:

$$s + g - 1 + q_1 + \cdots + q_n = 3g - 3 + n.$$

The so-called Vanishing Axiom, conjectured in [75] and proved in [100, 101], says that the Witten class $c_g(\alpha_1, \dots, \alpha_n)$ vanishes if any of $\alpha_1, \dots, \alpha_n$ reaches r . Therefore we assume that $\alpha_1, \dots, \alpha_n$ are in $\{1, \dots, r-1\}$. The numbers $\tau_{A_{r-1}}(g)$ that we are looking at are the r -spin intersection numbers with $n = 1$ (one-point). More precisely, they are defined by

$$\tau_{A_{r-1}}(g) := \langle \tau_{\alpha, q} \rangle_g, \quad (23)$$

where $\alpha \in \{1, \dots, r-1\}$ and $q \geq 0$ are uniquely determined by

$$2(r+1)g - 1 = \alpha + r(q+1). \quad (24)$$

The Witten class gives a particular *cohomological field theory (CohFT)* [83, 93] of rank $(r-1)$. Let us recall the definition of a general CohFT. A rank l CohFT is a quadruple $(V^l, \eta, \mathbb{1}, \Omega_{g,n})$, where V is a \mathbb{C} -vector space, η is a symmetric non-degenerate bilinear form, $\mathbb{1}$ is a particular element in V , called the unity, and $\{\Omega_{g,n}\}_{2g-2+n>0}$ is a collection of linear maps from $V^{\otimes n}$ to $H^{\text{even}}(\overline{\mathcal{M}}_{g,n}; \mathbb{C})$, satisfying the following axioms:

- C1. (total symmetry) Each $\Omega_{g,n}$ is S_n -invariant, where the action of S_n permutes both the marked points of $\overline{\mathcal{M}}_{g,n}$ and the tensor products $V^{\otimes n}$.
- C2. (splitting) Denote by q and s be the gluing maps

$$q: \overline{\mathcal{M}}_{g-1, n+2} \rightarrow \overline{\mathcal{M}}_{g,n}, \quad (25)$$

$$s: \overline{\mathcal{M}}_{g, n_1+1} \times \overline{\mathcal{M}}_{g, n_2+1} \rightarrow \overline{\mathcal{M}}_{g,n}. \quad (26)$$

Then it is required that $\forall x_1, \dots, x_n \in V$,

$$q^* \Omega_{g,n}(x_1, \dots, x_n) = \Omega_{g-1, n+2}(x_1, \dots, x_n, e_\alpha, e^\alpha), \quad (27)$$

$$s^* \Omega_{g,n}(x_1, \dots, x_n) = \Omega_{g_1, n_1+1}(x_1, \dots, x_{n_1}, e_\alpha) \Omega_{g_2, n_2+1}(e^\alpha, x_{n_1+1}, \dots, x_n). \quad (28)$$

- C3. (unity) Let $p: \overline{\mathcal{M}}_{g, n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ be the forgetful map. Then $\forall x_1, \dots, x_n \in V$,

$$\Omega_{g, n+1}(x_1, \dots, x_n, \mathbb{1}) = p^* \Omega_{g,n}(x_1, \dots, x_n), \quad (29)$$

$$\Omega_{0,3}(x_1, x_2, \mathbb{1}) = \eta(x_1, x_2). \quad (30)$$

Choose $e_1 = \mathbb{1}, e_2, \dots, e_l$ a basis of V , and denote $\eta_{\alpha\beta} := \eta(e_\alpha, e_\beta)$, $\eta = (\eta_{\alpha\beta})$, $(\eta^{\alpha\beta}) := \eta^{-1}$. It is natural to view V as the complex coordinate space $\{(v^1, \dots, v^l) \mid v^\alpha \in \mathbb{C}\}$ and therefore as the complex manifold \mathbb{C}^l . Define a power series $F = F(v) \in \mathbb{C}[[v^1, \dots, v^l]]$ by

$$F := \sum_{n \geq 3} \int_{\overline{\mathcal{M}}_{0,n}} \Omega_{0,n}(e_{\alpha_1}, \dots, e_{\alpha_n}) \frac{v^{\alpha_1} \dots v^{\alpha_n}}{n!}. \quad (31)$$

We call F the *genus-zero primary potential* (cf. [43, 57, 83, 93, 97, 98, 109]). Denote by $B \subset \mathbb{C}^l$ the domain of convergence for F around $v = 0$. Throughout the paper we assume that B contains an open ball centered at $v = 0$.

Remark 5. For a projective variety X , the Gromov–Witten classes associated to X give rise to a CohFT. For this case, it is sometimes helpful or even necessary to replace the ring $H^{\text{even}}(\overline{\mathcal{M}}_{g,n}; \mathbb{C})$ with $\mathcal{N} \otimes H^{\text{even}}(\overline{\mathcal{M}}_{g,n}; \mathbb{C})$, where \mathcal{N} is the Novikov ring [83, 93]. We did not write the axioms involving the Novikov ring because for the specific CohFTs that are considered in this paper the power series F is actually a polynomial and so $B = \mathbb{C}^l$. Nevertheless, it will be interesting to generalize the results of this paper to the situation when the Novikov ring is introduced.

A CohFT $(V, \eta, \mathbb{1}, \Omega_{g,n})$ is called *homogeneous* of *charge* d if it satisfies the following axiom:

C4. There is a vector field E on B which for some choice of the basis e_α of V has the form

$$E = \left(1 - \frac{d}{2}\right) v^\alpha \partial_\alpha + r^\alpha \partial_\alpha - \sum_{\alpha=1}^l \mu_\alpha v^\alpha \partial_\alpha, \quad (32)$$

and such that if we define the action of E on Ω by

$$\begin{aligned} (E\Omega)_{g,n}(e_{\alpha_1}, \dots, e_{\alpha_n}) &:= \left(\text{gr} + \frac{2-d}{2}n - \sum_{i=1}^n \mu_{\alpha_i} \right) \Omega_{g,n}(e_{\alpha_1}, \dots, e_{\alpha_n}) \\ &\quad + p_* \Omega_{g,n+1}(e_{\alpha_1}, \dots, e_{\alpha_n}, r^\alpha e_\alpha), \end{aligned} \quad (33)$$

where p denotes the forgetful map, and gr is the grading operator defined by

$$\text{gr} \phi := q \phi, \quad \text{if } \phi \in H^{2q}(\overline{\mathcal{M}}_{g,n}; \mathbb{C}), \quad (34)$$

then

$$(E\Omega)_{g,n} = ((g-1)d + n) \Omega_{g,n}, \quad \forall 2g - 2 + n > 0. \quad (35)$$

The genus g correlators of the CohFT are defined by

$$\int_{\overline{\mathcal{M}}_{g,n}} \Omega_{g,n}(e_{\alpha_1}, \dots, e_{\alpha_n}) \psi_1^{q_1} \cdots \psi_n^{q_n} =: \langle \tau_{\alpha_1, q_1} \cdots \tau_{\alpha_n, q_n} \rangle_g^\Omega, \quad q_1, \dots, q_n \geq 0. \quad (36)$$

Below, the label Ω will often be omitted. Here and throughout the paper, free Greek indices take the integer values from 1 to l , the Einstein summation convention will be used for repeated Greek indices with one up and one down, and the tensors $\eta^{\alpha\beta}$ and $\eta_{\alpha\beta}$ will be used to raise and lower the Greek indices. The genus g free energy of the CohFT is defined by

$$\mathcal{F}_g(\mathbf{t}) := \sum_{n \geq 0} \sum_{q_1, \dots, q_n \geq 0} \frac{t^{\alpha_1, q_1} \cdots t^{\alpha_n, q_n}}{n!} \langle \tau_{\alpha_1, q_1} \cdots \tau_{\alpha_n, q_n} \rangle_g, \quad (37)$$

where $\mathbf{t} := (t^{\alpha, q})_{\alpha=1, \dots, l, q \geq 0}$ denotes the infinite vector of indeterminates. The exponential

$$e^{\sum_{g \geq 0} \epsilon^{2g-2} \mathcal{F}_g(\mathbf{t})} =: Z \quad (38)$$

is called the *partition function* of the CohFT, where ϵ is an indeterminate. It satisfies the following string equation

$$\sum_{q \geq 1} t^{\alpha, q} \frac{\partial Z}{\partial t^{\alpha, q-1}} + \frac{1}{2\epsilon^2} \eta_{\alpha\beta} t^{\alpha, 0} t^{\beta, 0} Z = \frac{\partial Z}{\partial t^{1, 0}}. \quad (39)$$

An immediate consequence of the string equation is that

$$\langle \tau_{\alpha, q-1} \rangle_g = \langle \tau_{\alpha, q} \tau_{1, 0} \rangle_g. \quad (40)$$

The CohFT $\Omega_{g,n}$ given by the Witten class $c_g(\alpha_1, \dots, \alpha_n)$ is also recognized as the FJRW CohFT of A_{r-1} -type. This is the reason for the notation used in (23). For each simply-laced simple Lie algebra \mathfrak{g} of rank l , Fan, Jarvis and Ruan [66, 67] constructed a rank l homogeneous CohFT from a certain simple singularity such that the corresponding Frobenius manifold is isomorphic to the Frobenius manifold associated to the Weyl group of \mathfrak{g} [45, 122]. This CohFT will be referred to as the FJRW CohFT of \mathfrak{g} -type, whose partition function is shown to be a particular tau-function for the DS hierarchy of \mathfrak{g} -type [38, 67, 110, 112] (cf. [65, 68, 71, 72, 78, 82, 89, 113]). In [89] Liu, Ruan and Zhang introduced the notion of a *partial CohFT*, and constructed certain partial CohFTs associated to simple singularities, whose partition functions are proved *ibid.* to be tau-functions for the DS hierarchies of BCFG-type. Correlators of these CohFTs or partial CohFTs are referred to as the FJRW–LRZ invariants. The main focus of this paper will be on the ADE cases, leaving the more detailed studies of the BCFG cases to future publications.

It should be noted that the terminology “DS hierarchy of \mathfrak{g} -type” refers to the DS hierarchy, under the choice of a principal nilpotent element, associated to the untwisted affine Kac–Moody algebra $\hat{\mathfrak{g}}^{(1)}$ (see page 1402 of [12] for more details; cf. also [13]).

Now let \mathfrak{g} be a *simply-laced* simple Lie algebra of rank l with the normalized Cartan–Killing form $(\cdot|\cdot)$. Denote by r the Coxeter number of \mathfrak{g} , and by m_1, \dots, m_l the exponents of \mathfrak{g} . Here we order m_1, \dots, m_l such that $1 = m_1 \leq m_2 \leq \dots \leq m_l = r - 1$, except for the D_l case, where we set $m_\alpha = 2\alpha - 1$ for $\alpha = 1, \dots, l - 1$ and $m_l = l - 1$. (Observe the difference with [12, 13], where for the D_l case the exponents are numbered by $1 = m_1 \leq m_2 \leq \dots \leq m_l = r - 1$.) Let $(V^l, \eta, \mathbb{1}, \Omega_{g,n})$ denote the FJRW CohFT of \mathfrak{g} -type. Choose a basis e_α of V satisfying

$$\Omega_{g,n}(e_{\alpha_1}, \dots, e_{\alpha_n}) \in H^{2(s+g-1)}(\overline{\mathcal{M}}_{g,n}), \quad s := \frac{\sum_{i=1}^l m_{\alpha_i} - n - (2g - 2)}{r}. \quad (41)$$

The existence of such a choice can be verified case by case from the data provided in [67], and this implies that the CohFT $\Omega_{g,n}$ is *homogeneous* of charge $d = \frac{r-2}{r}$. The associated correlators (36) are called *genus g FJRW invariants of \mathfrak{g} -type*. The degree-dimension matching implies that $\langle \tau_{\alpha_1, q_1} \cdots \tau_{\alpha_n, q_n} \rangle_g$ vanishes unless

$$\frac{\sum_{i=1}^n m_{\alpha_i} - n - (2g - 2)}{r} + g - 1 + \sum_{i=1}^n q_i = 3g - 3 + n. \quad (42)$$

For given (α_i, q_i) , $i = 1, \dots, n$, for simplicity we sometimes omit the subindex g in $\langle \cdots \rangle_g$, because for all possibly non-zero invariants this subindex can be reconstructed by (42); in such an omission, if the reconstructed g is not an integer, $\langle \tau_{\alpha_1, q_1} \cdots \tau_{\alpha_n, q_n} \rangle$ is defined as 0. It is also convenient to take $\epsilon = 1$ in (38) for the definition of the partition Z , i.e., we have

$$Z = Z(\mathbf{t}) := e^{\sum_{g \geq 0} \mathcal{F}_g(\mathbf{t})}, \quad (43)$$

and the string equation reads

$$\sum_{q \geq 1} t^{\alpha, q} \frac{\partial Z(\mathbf{t})}{\partial t^{\alpha, q-1}} + \frac{1}{2} \eta_{\alpha\beta} t^{\alpha, 0} t^{\beta, 0} Z(\mathbf{t}) = \frac{\partial Z(\mathbf{t})}{\partial t^{1, 0}}. \quad (44)$$

The numbers $\tau_{\mathfrak{g}}(g)$ that we are studying in this paper are defined by

$$\tau_{\mathfrak{g}}(g) := \langle \tau_{\alpha, q} \rangle, \quad g \geq 0, \quad (45)$$

where

$$2(r+1)g - 1 = m_\alpha + r(q+1). \quad (46)$$

It should be noticed that for the case $\mathfrak{g} = D_l$ with l being an even number, there are two equal exponents $m_{l/2}$ and m_l , so for this case, the different $(\alpha = l/2, q)$ and $(\alpha = l, q)$ correspond to the same g . However, with an appropriate choice of the basis, the numbers $\langle \tau_{l, q} \rangle$ vanish, and we use $\tau_{\mathfrak{g}}(g)$ to denote $\langle \tau_{l/2, q} \rangle$.

3. DIFFERENTIAL EQUATION

The topological and dual topological ODEs of \mathfrak{g} -type are introduced in [12] for computing the FJRW–LRZ invariants, which are obtained as a result of the theorems of Fan–Jarvis–Ruan and Liu–Ruan–Zhang together with an application of the matrix-resolvent method [11, 12, 13] for the Drinfeld–Sokolov hierarchy of \mathfrak{g} -type [38]. In this section, via reducing the dual topological ODE to a scalar differential equation, we generalize parts (i) and (ii) of Theorem 2 of the Introduction from A_4 to an arbitrary simple Lie algebra \mathfrak{g} .

Fix \mathfrak{h} a Cartan subalgebra of \mathfrak{g} and let Δ be the root system. Choose a set of simple roots Π , and let $E_1, \dots, E_l, F_1, \dots, F_l$ be the Weyl generators. Denote by θ the highest root with respect

to Π , and by $E_{-\theta}$ a root vector associated to $-\theta$. The semisimple element $\Lambda = \Lambda(\lambda)$ (aka the Kostant element) is defined by $\Lambda(\lambda) = \sum_{\alpha=1}^l E_{\alpha} + \lambda E_{-\theta}$, $\lambda \in \mathbb{C}$. Kostant [84] shows that for any $\lambda \neq 0$, \mathfrak{g} has the orthogonal decomposition with respect to $(\cdot|\cdot)$, i.e. $\mathfrak{g} = \text{Ker ad}_{\Lambda(\lambda)} \oplus \text{Im ad}_{\Lambda(\lambda)}$, $\text{Ker ad}_{\Lambda(\lambda)} \perp \text{Im ad}_{\Lambda(\lambda)}$. Denote by $L(\mathfrak{g}) = \mathfrak{g} \otimes \mathbb{C}[\lambda, \lambda^{-1}]$ the loop algebra, and by ρ^{\vee} the Weyl co-vector (which is the unique element in \mathfrak{h} satisfying $[\rho^{\vee}, E_{\alpha}] = E_{\alpha}$). Introduce a grading operator on $L(\mathfrak{g})$

$$\text{gr} := \text{ad}_{\rho^{\vee}} + h \lambda \frac{d}{d\lambda}. \quad (47)$$

An element q in $L(\mathfrak{g})$ is called homogeneous of principal degree k if $\text{gr } q = k q$. It was proven by Kac [76] that the kernel of $\text{ad}_{\Lambda(\lambda)}$ in $L(\mathfrak{g})$ has the form $\bigoplus_{j \in E} \mathbb{C} \Lambda_j(\lambda)$. Here, E is the exponent set of $L(\mathfrak{g})$, and $\Lambda_j(\lambda) \in L(\mathfrak{g})$ are homogeneous of principal degree j , normalized by $\Lambda_1 = \Lambda$, $\Lambda_{m_{\alpha} + \ell h}(\lambda) = \lambda^{m_{\alpha}}(\lambda) \lambda^{\ell}$, $\ell \geq 0$, and $(\Lambda_{m_{\alpha}} | \Lambda_{m_{\beta}}) = h \lambda \eta_{\alpha\beta}$ for some non-degenerate symmetric constant matrix η .

Write

$$\rho^{\vee} = \sum_{i=1}^n x_i H_i, \quad x_i \in \mathbb{C},$$

and define $I_- = 2 \sum_{i=1}^n x_i F_i$. Then I_+, I_-, ρ^{\vee} form an $sl_2(\mathbb{C})$ Lie algebra:

$$[\rho^{\vee}, I_+] = I_+, \quad [\rho^{\vee}, I_-] = -I_-, \quad [I_+, I_-] = 2\rho^{\vee}. \quad (48)$$

According to [7, 84], there exist elements $\gamma^1, \dots, \gamma^n \in \mathfrak{g}$ such that

$$\text{Ker ad}_{I_-} = \text{Span}_{\mathbb{C}}\{\gamma^1, \dots, \gamma^n\}, \quad [\rho^{\vee}, \gamma^i] = -m_i \gamma^i. \quad (49)$$

Fix $\{\gamma^1, \dots, \gamma^n\}$, then the lowest weight decomposition of \mathfrak{g} has the form

$$\mathfrak{g} = \bigoplus_{i=1}^n \mathcal{L}^i, \quad \mathcal{L}^i = \text{Span}_{\mathbb{C}}\{\gamma^i, \text{ad}_{I_+} \gamma^i, \dots, \text{ad}_{I_+}^{2m_i} \gamma^i\}. \quad (50)$$

Here each \mathcal{L}^i is an $sl_2(\mathbb{C})$ -module. Any \mathfrak{g} -valued function $M(\lambda)$ can be uniquely represented as

$$M(\lambda) = \sum_{i=1}^n S_i(\lambda) \text{ad}_{I_+}^{2m_i} \gamma^i + \sum_{i=1}^n \sum_{m=0}^{2m_i-1} K_{im}(\lambda) \text{ad}_{I_+}^m \gamma^i, \quad (51)$$

where $S_i(\lambda)$, $K_{im}(\lambda)$ are certain complex-valued functions. Note that $\text{ad}_{I_+}^{2m_i} \gamma^i$ is the highest weight vector of \mathcal{L}^i , $i = 1, \dots, n$.

The dual topological ODE of \mathfrak{g} -type [12] is an ODE for a \mathfrak{g} -valued function G defined by

$$\left[\frac{dG}{dx}, E_{-\theta} \right] + [G, I_+] + xG = 0. \quad (52)$$

Write

$$G(x) = \sum_{\alpha=1}^l \phi_{\alpha}(x) \text{ad}_{I_+}^{2m_{\alpha}} \gamma^{\alpha} + \sum_{\alpha=1}^l \sum_{m=0}^{2m_{\alpha}-1} \tilde{K}_{\alpha m}(x) \text{ad}_{I_+}^m \gamma^{\alpha}. \quad (53)$$

It is shown in [12] that the ODE (52) is equivalent to an ODE for $\phi = (\phi_1, \dots, \phi_l)^T$

$$\frac{d\phi}{dx} = \sum_{i=-1}^{2h-2} x^i V_i \phi, \quad (54)$$

where V_i ($i = -1, \dots, 2h-2$) are constant $l \times l$ matrices and $V_{-1} = \text{diag}(-m_{l+1-\alpha}/h)_{\alpha=1, \dots, l}$. (Equivalence between systems of ODEs means that their solutions have a one-to-one correspondence.) Here we note that $x = 0$ is a Fuchsian singular point of (54) and that the matrices V_{-1} ,

\dots, V_{2h-2} are determined by the lowest-weight-structure-constants of \mathfrak{g} . It is obvious from (54) that the dimension of the space of solutions to (52) is equal to the rank of \mathfrak{g} .

Denote by $(G_\alpha)_{\alpha=1,\dots,l}$ a solution basis of (52), and by $\phi_{\alpha;1}, \dots, \phi_{\alpha;l}$ the ϕ -coefficients of G_α . The matrix Φ defined by $\Phi_{\beta\alpha} := \phi_{\alpha;\beta}$ is called the fundamental solution matrix, which can be normalized by using the following initial condition near $x = 0$:

$$\Phi = D \left(I + \sum_{m \geq 1} \Phi_m x^m \right), \quad D = \text{diag} \left(x^{-\frac{m_{l+1}-\alpha}{h}} \right)_{\alpha=1,\dots,l}. \quad (55)$$

For each fixed α we allow G_α to have a multiplicative non-zero constant and keep using the notations G_α and $\phi_{\alpha;\beta}$. The following proposition is proved in [12], and we review its proof here.

Proposition 1. *The series $\phi_{\alpha;l}$ can be expressed in terms of the FJRW invariants of \mathfrak{g} -type by*

$$\phi_{\alpha;l}(x) = \frac{x^{-\frac{1}{r}}}{\Gamma(\frac{m_\alpha}{h})} \delta_{\alpha,l} + \frac{x^{\frac{m_\alpha}{r}}}{\Gamma(\frac{m_\alpha}{r})} \sum_{w \geq 0} c_{\alpha,w} x^{m_\alpha + (r+1)w+1}, \quad (56)$$

where

$$c_{\alpha,w} = (-1)^{m_\alpha + (r+1)w} \langle \tau_{\alpha, m_\alpha + (r+1)w} \rangle (-r)^{\frac{m_\alpha + 1 + rw}{2}}, \quad w \geq 0. \quad (57)$$

Here we note that when \mathfrak{g} is a non-simply-laced simple Lie algebra, the $\langle \tau_{\alpha, m_\alpha + (r+1)w} \rangle$ in the right-hand side of (57) should be considered as the one-point correlators of the corresponding Liu–Ruan–Zhang partial CohFT (cf. [12, 13, 89]).

Proof of Proposition 1. Recall that the topological ODE of \mathfrak{g} -type [12] is the \mathfrak{g} -valued ODE

$$\frac{dM(\lambda)}{d\lambda} = [M(\lambda), \Lambda(\lambda)]. \quad (58)$$

About this ODE, the following statements are proved in [12, 13].

- (a) The dimension of the formal Puiseux series solutions to (58) is equal to the rank of \mathfrak{g} .
- (b) There exists a unique basis M_1, \dots, M_l of the formal solutions to (58) such that

$$M_\alpha(\lambda) = \lambda^{-\frac{m_\alpha}{h}} \left[\Lambda_{m_\alpha}(\lambda) + \sum_{k \geq 1} M_{\alpha,k}(\lambda) \right], \quad (59)$$

$$M_{\alpha,k}(\lambda) \in L(\mathfrak{g}), \quad \text{gr } M_{\alpha,k}(\lambda) = [m_\alpha - (h+1)k] M_{\alpha,k}(\lambda). \quad (60)$$

- (c) Denote $\kappa = (\sqrt{-r})^{-r}$. The following identity for one-point FJRW invariants of \mathfrak{g} -type is true:

$$\kappa^{\frac{2}{r+1}} \sqrt{-r} \sum_{g, q \geq 0} (-1)^q \frac{\binom{m_\alpha}{r} q+2}{(\kappa^{\frac{1}{r+1}} \lambda)^{\frac{m_\alpha}{r} + q+2}} \langle \tau_{\alpha,q} \rangle_g = (E_{-\theta} | M_\alpha(\lambda)) - \lambda^{-\frac{r-1}{r}} \delta_{\alpha,l}. \quad (61)$$

Write

$$(E_{-\theta} | M_\alpha(\lambda)) = \lambda^{-\frac{m_\alpha}{r}} \sum_{q \geq 0} S_{\alpha,q} \lambda^{-q} =: S_\alpha(\lambda), \quad S_{\alpha,q} \in \mathbb{Q}. \quad (62)$$

We have for $w \in \mathbb{Z}$ and $(1 + m_\alpha + rw)/2 \in \mathbb{Z}_{\geq 0}$,

$$\langle \tau_{\alpha, m_\alpha + (r+1)w} \rangle = (-1)^{m_\alpha + (r+1)w} \frac{S_{\alpha, m_\alpha + (r+1)w+2}}{(-r)^{\frac{m_\alpha + 1 + rw}{2}} \binom{m_\alpha}{r}_{m_\alpha + (r+1)w+2}}. \quad (63)$$

According to the definition given in [12], solutions to topological and dual topological ODEs are related via the Laplace transform $M(\lambda) = \int G(x) e^{-\lambda x} dx$. In particular, the series $\phi_{\alpha;l}$ are related to S_α by

$$S_\alpha(\lambda) = \int_C \phi_{\alpha;l}(x) e^{-\lambda x} dx, \quad (64)$$

where C is a carefully chosen contour (which can depend on α) on the x -plane. We have

$$\phi_{\alpha;l}(x) = \sum_{q \geq 0} S_{\alpha,q} \frac{x^{q + \frac{m_\alpha}{r} - 1}}{\Gamma(q + \frac{m_\alpha}{r})} = \frac{x^{\frac{m_\alpha}{r} - 1}}{\Gamma(\frac{m_\alpha}{r})} \sum_{q \geq 0} S_{\alpha,q} \frac{x^q}{\left(\frac{m_\alpha}{r}\right)_q}. \quad (65)$$

The proposition is proved. \square

An immediate consequence of Proposition 1 is that the intersections numbers $\tau_{\mathfrak{g}}(g)$ grow at most exponentially. Let us introduce the series $a(x)$ by

$$a(x) := (-r)^{-\frac{1}{2(r+1)}} \sum_{\alpha=1}^l \Gamma\left(\frac{m_\alpha}{r}\right) (-1)^{\frac{m_\alpha - r}{r}} \phi_{\alpha;l}(x), \quad (66)$$

where $\phi_{\alpha;l}(x)$ are given by (56). More explicitly,

$$\begin{aligned} a(x) &= (-r)^{-\frac{1}{2(r+1)}} (-x)^{-\frac{1}{r}} + \sum_{\alpha=1}^l \sum_{w \geq 0} (-1)^{m_\alpha + (r+1)w + \frac{m_\alpha - r}{r}} \\ &\quad \langle \tau_{\alpha, m_\alpha + (r+1)w} \rangle (-r)^{\frac{m_\alpha + 1 + rw}{2}} x^{\frac{(r+1)m_\alpha + r(r+1)w + r}{r}} \\ &= \left(-(-r)^{\frac{r}{2(r+1)}} x \right)^{-\frac{1}{r}} + \left(-(-r)^{\frac{r}{2(r+1)}} x \right)^{-\frac{1}{r}} \sum_{g \geq 0} \tau_{\mathfrak{g}}(g) \left(-(-r)^{\frac{r}{2(r+1)}} x \right)^{2g + \frac{2g}{r}}. \end{aligned}$$

Denote $t = -(-r)^{\frac{r}{2(r+1)}} x$ and

$$b(t) = t^{-\frac{1}{r}} + t^{-\frac{1}{r}} \sum_{g \geq 1} \tau_{\mathfrak{g}}(g) t^{2g + \frac{2g}{r}} = \sum_{g \geq 0} \tau_{\mathfrak{g}}(g) t^{2g \frac{r+1}{r} - \frac{1}{r}}.$$

We have $a(x) = b(t)$. We are ready to generalize part (i) of Theorem 2 by showing that $b(t)$ satisfies an ODE with polynomial coefficients.

Theorem 4. *Let \mathfrak{g} be an arbitrary simple Lie algebra of rank l . The component ϕ_l satisfies a linear ODE with polynomial coefficients of order at most l . In other words, for every fixed α , the series $\phi_{\alpha;l}$ satisfies this ODE. In particular, $b(t)$ satisfies an ODE of order at most l .*

Proof. For \mathfrak{g} with $l = 1$, the statement is trivial. Below we consider $l \geq 2$. As we have mentioned, the dual topological ODE is equivalent to system (54). This system consists of l linear equations for ϕ_1, \dots, ϕ_l . Denote these equations by e_1, \dots, e_l . Let N be a positive integer. Now consider the following linear combination of equations

$$e := \sum_{k=0}^N c_k^l \frac{d^k e_l}{dx^k} + \sum_{\alpha=1}^{l-1} \sum_{k=1}^N c_k^\alpha \frac{d^{k-1} e_\alpha}{dx^{k-1}}. \quad (67)$$

Requiring that the coefficients of $\phi_1, \dots, \phi_{l-1}, \frac{d\phi_1}{dx}, \dots, \frac{d\phi_{l-1}}{dx}, \dots, \frac{d^{N-1}\phi_1}{dx^{N-1}}, \dots, \frac{d^N\phi_{l-1}}{dx^N}$ vanish gives rise to a system of linear equations for c_0^l and $c_k^\alpha, k = 1, \dots, N$. The number of equations is equal to $(l-1)(N+1)$, while the number of unknowns is $1 + Nl$. Take $N = l-1$, we have

$$1 + Nl - (l-1)(N+1) = 2 + N - l = 1. \quad (68)$$

According to the Gauss elimination we know that this linear system over the field $\mathbb{C}(x)$ has a non-zero solution c_0^l and $c_k^\alpha, k = 1, \dots, N$. The theorem is proved by further noticing that $\frac{d^k e_l}{dx^k}$ ($k = 0, \dots, N$) and $\frac{d^{k-1} e_\alpha}{dx^{k-1}}$ ($\alpha = 1, \dots, l-1, k = 1, \dots, N$) together are linearly independent. \square

It is clear that part (i) of Theorem 2 is a special case of Theorem 4 with the particular form of the linear ODE (8) computed from the standard $sl_5(\mathbb{C})$ realization with the $\Lambda(\lambda)$ given by

$$\Lambda(\lambda) = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ \lambda & 0 & 0 & 0 & 0 \end{pmatrix}, \tag{69}$$

and from the change of independent variable $t \mapsto X = t^{\frac{2(r+1)}{r}}$ and $\varphi(X) = b(t) t^{\frac{1}{r}}$.

4. COMPUTING RESIDUES

In this section, we carry out several technical preparations for the later sections. It is very convenient here to permit the variable r , which originally came from r -spin classes and hence was a positive integer, to take on arbitrary complex values, since essentially all of the identities we need are polynomial in r (at least at the level of the individual coefficients of the generating functions appearing).

Let r be a complex number and let L be the pseudodifferential operator

$$L = \partial^r + Cx, \tag{70}$$

where $\partial = d/dx$ and C is an arbitrarily given constant. We are to compute the residues $z_k(x)$ of the pseudodifferential operators $L^{k/r}$, i.e.

$$z_k(x) := \text{res } L^{\frac{k}{r}}. \tag{71}$$

We note that most of the time we consider k as a nonnegative integer, although sometimes we give results valid for any $k \in \mathbb{C}$. For the definition of a pseudodifferential operator and its residue we refer to [29, 34, 79].

4.1. Closed formula. Let us first introduce some notations. Define polynomials $c_{p,j}(r) \in \mathbb{Q}[r]$ ($0 \leq p \leq j$) by the generating function

$$\frac{1}{p!} \left(\frac{(1+x)^{r+1} - 1 - (r+1)x}{(r+1)x} \right)^p = \sum_{j=p}^{\infty} c_{p,j}(r) x^j \tag{72}$$

(then $c_{p,j}(r)$ has degree j and is divisible by r^p), the first few values being

	$p = 0$	$p = 1$	$p = 2$	$p = 3$	$p = 4$	$p = 5$
$j = 0$	1					
$j = 1$	0	$r/2$				
$j = 2$	0	$\frac{1}{3} \binom{r}{2}$	$\frac{(r/2)^2}{2!}$			
$j = 3$	0	$\frac{1}{4} \binom{r}{3}$	$\frac{r^2(r-1)}{12}$	$\frac{(r/2)^3}{3!}$		
$j = 4$	0	$\frac{1}{5} \binom{r}{4}$	$\frac{r(5r-8)}{72} \binom{r}{2}$	$\frac{r^3(r-1)}{48}$	$\frac{(r/2)^4}{4!}$	
$j = 5$	0	$\frac{1}{6} \binom{r}{5}$	$\frac{r(4r-7)}{60} \binom{r}{3}$	$\frac{r^3(r-1)(7r-10)}{576}$	$\frac{r^4(r-1)}{288}$	$\frac{(r/2)^5}{5!}$

The polynomials $c_{p,j}(r)$ ($0 \leq p \leq j \leq 5$)

The result of this subsection is given by the following proposition.

Proposition 2. *Let $L = \partial^r + Cx$. For any $r, \lambda \in \mathbb{C}$, we have*

$$L^\lambda = \sum_{j,s \geq 0} C^{j+s} d_{j,s}(\lambda, r) x^s \partial_x^{r(\lambda-s)-(r+1)j}, \quad (73)$$

where $d_{j,s}(\lambda, r)$ is a polynomial in λ and r with rational coefficients, given explicitly by

$$d_{j,s}(\lambda, r) = \frac{1}{s!} \sum_{p=0}^j c_{p,j}(r) (\lambda)_{s+p+j}^-. \quad (74)$$

Here $(\lambda)_n^- := \prod_{m=1}^n (\lambda - m + 1)$ denotes the descending Pochhammer symbol.

Proof. By the rules for manipulating pseudodifferential operators [29, 34, 79], it is clear that L^λ has the form (73) for some polynomials $d_{j,s}(\lambda, r)$. It then suffices to prove (74) for $\lambda \in \mathbb{Z}_{\geq 0}$, which we do by induction, the case $\lambda = 0$ being trivial (both sides of (74) then reduce to $\delta_{j,0} \delta_{s,0}$). Expanding the identity $L^{\lambda+1} = L^\lambda L$, we find that $d_{j,s}$ satisfies the recursive relation

$$d_{j,s}(\lambda + 1, r) - d_{j,s-1}(\lambda, r) - d_{j,s}(\lambda, r) = (r(\lambda - s) - (r + 1)(j - 1)) d_{j-1,s}(\lambda, r).$$

If we multiply both sides of this by $s!$ and replace $d_{j,s}$ everywhere by the expression given in (74), then the left-hand side equals

$$\sum_{p \geq 0} c_{p,j} [(\lambda + 1)_{s+p+j}^- - s(\lambda)_{s+p+j-1}^- - (\lambda)_{s+p+j}^-] = \sum_{p \geq 0} (p + j) c_{p,j} (\lambda)_{s+p+j-1}^-$$

and the right-hand side equals

$$\sum_{p \geq 0} c_{p,j-1} [r(\lambda - s) - (r + 1)(j - 1)] (\lambda)_{s+p+j-1}^- = \sum_{p \geq 0} c_{p,j-1} [r(\lambda)_{s+p+j}^- + (rp - j + 1)(\lambda)_{s+p+j-1}^-].$$

(Define $c_{p,j} = 0$ if $p > j$.) Comparing the coefficients of $(\lambda)_n^-$ on both sides, we see that the claim follows from the recursion $(p + j) c_{p,j} = (rp - j + 1) c_{p,j-1} + r c_{p-1,j-1}$, which follows easily by differentiating (72).

We observe that there is a slightly different proof, which does not depend on the polynomiality, obtained by using the identity $L L^\lambda = L^\lambda L$ instead of $L^\lambda L = L^{\lambda+1}$. Expanding both sides and comparing coefficients, we find the identity

$$(r(\lambda - s + 1) - (r + 1)j) d_{j,s-1}(\lambda, r) - r s d_{j,s}(\lambda, r) = \sum_{m=1}^j \binom{r}{m+1} (s+m)_{m+1}^- d_{j-m,s+m}(\lambda, r),$$

which determines the $d_{j,s}$ completely by a double induction (first on j , and then for a given j on s). The proposition then follows by verifying that the right-hand side of (74) satisfies the same identity, which is an elementary exercise using the generating function (72). \square

Corollary. *For arbitrary positive integers r and k , the polynomial $z_k(x) = \text{res } L_r^{\frac{k}{r}}$ is given by $z_k(0) = C^j d_{j,0}(k/r, r)$ if $k = -1 + (r + 1)j$ with $j \geq 0$ and is 0 otherwise, and similarly for any $s \geq 0$, $z_k^{(s)}(0)$ is equal to $C^{s+j} d_{j,s}(k/r, r)$ if $k = -1 + (r + 1)j$ with $j \geq 0$ and vanishes otherwise.*

4.2. Product formula. In this subsection we will derive another formula for $z_k(0)$ using the wave-function-pair approach (see Appendix A or [53]). Before doing this, we will first introduce some functions and prove several lemmas that are useful for the construction.

Definition 1. Define a power series (algebraic if r is rational)

$$w = 1 + u - \frac{r-1}{6} u^2 + \frac{(r-1)(2r+1)}{72} u^3 - \dots \quad (75)$$

by

$$\frac{w^{r+1}}{r(r+1)} - \frac{w}{r} + \frac{1}{r+1} = \frac{u^2}{2} \quad (76)$$

and define coefficients $C_n(r, j) \in \mathbb{Q}[r, j]$, $n \geq 0$ by

$$\frac{w^{j+1} - 1}{j+1} = \sum_{n \geq 0} C_n(r, j) u^{n+1} \quad (\text{replace LHS by } \log w \text{ if } j = -1), \quad (77)$$

the first few values being

$$\begin{aligned} C_0(r, j) &= 1, & C_1(r, j) &= \frac{j}{2} - \frac{r-1}{6}, & C_2(r, j) &= \frac{j(j-r)}{6} + \frac{(r-1)(2r+1)}{72}, \\ C_3(r, j) &= \frac{j(j-r)(j-r-1)}{24} - \frac{(r-1)(r+2)(2r+1)}{540}. \end{aligned}$$

Define $C_{-2}(r, j) = C_{-1}(r, j) = 0$.

Remark 6. Immediately from the definition we find a further property for the coefficients $C_n(r, j)$, that is, they have an S_3 -symmetry generated by two involutions

$$C_n(r, j) = (-1)^n C_n\left(-r-1, \frac{n-3}{2} - j\right) = (r+1)^n C_n\left(\frac{-r}{r+1}, \frac{j-r}{r+1}\right). \quad (78)$$

We hope to investigate this very intriguing property and its applications later.

We now define a power series $f_j(T) = f_{r,j}(T) \in \mathbb{Q}[r, j][[T]]$ (we usually omit r) by

$$f_j(T) := \sum_{k \geq 0} (2k+1)!! C_{2k}(r, j) (-T)^k. \quad (79)$$

Note that this is defined for all r and j in \mathbb{C} , not just for non-negative integers, because the $C_n(r, j)$ are polynomials.

Lemma 1. *The power series $f_j(T)$ ($j \in \mathbb{C}$) satisfy the following two identities:*

$$f_{j+1}(T) = \left(1 + \left(\frac{r-1}{2} - j\right)T + (r+1)T^2 \frac{d}{dT}\right) f_j(T), \quad (80)$$

$$f_{j+r}(T) = f_j(T) - r j T f_{j-1}(T). \quad (81)$$

Proof. We have by definition

$$w^j \frac{dw}{du} = \sum_{n \geq 0} (n+1) C_n(r, j) u^n. \quad (82)$$

Applying d/du to (76) gives

$$\left(\frac{w^r}{r} - \frac{1}{r}\right) \frac{dw}{du} = u \quad \Rightarrow \quad w^{r+j} \frac{dw}{du} - w^j \frac{dw}{du} = r u w^j.$$

Substituting (82) and (77) into this identity, comparing the Taylor coefficients of u , and noticing that $C_0(r, j) = 1$, $C_1(r, j) = \frac{j}{2} - \frac{r-1}{6}$, we obtain that for all $n \geq 0$,

$$(n+1) C_n(r, r+j) = (n+1) C_n(r, j) + r j C_{n-2}(r, j-1) + r \delta_{n,1},$$

which proves (81). Similarly, we have

$$\begin{aligned} (w^{j+1} - w^j) \frac{dw}{du} &= w^{j+1} \left(w^r \frac{dw}{du} - r u \right) - w^j \frac{dw}{du} \\ &= w^j \left(\frac{r(r+1)}{2} u^2 - r + (r+1)w \right) \frac{dw}{du} - r u w^{j+1} - w^j \frac{dw}{du}. \end{aligned}$$

Namely,

$$(w^{j+1} - w^j) \frac{dw}{du} = -\frac{r+1}{2} u^2 w^j \frac{dw}{du} + u w^{j+1},$$

which implies (80). The lemma is proved. \square

We note that the above relations (80) and (81) determine $f_{r,j}(T) \in 1 + T\mathbb{C}[r, j][[T]]$ uniquely for all $r, j \in \mathbb{C}$. To prove this, we may assume that $j \geq 0$ and $r \geq 2$ are integers (since these countably many values define the polynomial coefficients uniquely). Then (80) implies by induction that $f_1, f_2, f_3, \dots, f_r$ are uniquely determined by f_0 and then (81) with $j = 0$ gives a differential equation for f_0 , which has a unique solution in $1 + T\mathbb{C}[r, j][[T]]$. (In fact, this proof shows that the $f_{j,r}(T)$ are uniquely determined by just (80) and the equation $f_0 = f_r$.)

We now define an odd Laurent series (again algebraic if r is rational)

$$X = \frac{1}{u} - \frac{r-1}{6} u + \frac{(r-1)(2r+1)(r-3)}{360} u^3 + \dots \in \frac{1}{u} \mathbb{Q}[r][[u^2]]$$

by the equation

$$\frac{(X+1)^{r+1} - (X-1)^{r+1}}{2(r+1)} = \frac{1}{u^r}, \quad (83)$$

and define coefficients $\tilde{C}_n(r, i, j) \in \mathbb{Q}[r, i, j]$ by

$$-(X+1)^i (X-1)^j \frac{dX}{du} = \sum_{n \geq 0} \tilde{C}_n(r, i, j) u^{n-i-j-2}. \quad (84)$$

The first two $\tilde{C}_n(r, i, j)$ are given by

$$\tilde{C}_0(r, i, j) = 1, \quad \tilde{C}_1(r, i, j) = i - j. \quad (85)$$

Taking $i = j = 0$ in (84), we have

$$-\frac{dX}{du} = \sum_{n \geq 0} \tilde{C}_n(r, 0, 0) u^{n-2}. \quad (86)$$

Integrating this equality with respect to u we find that

$$-X = -\frac{1}{u} + \sum_{n \geq 2} \frac{1}{n-1} \tilde{C}_n(r, 0, 0) u^{n-1}. \quad (87)$$

Lemma 2. *The numbers $\tilde{C}_n(r, i, j)$ satisfy the following relations:*

$$\tilde{C}_n(r, i+1, j) - \tilde{C}_n(r, i, j+1) = 2\tilde{C}_{n-1}(r, i, j), \quad (88)$$

$$\tilde{C}_n(r, i+r+1, j) - \tilde{C}_n(r, i, j+r+1) = 2(r+1)\tilde{C}_{n-1}(r, i, j). \quad (89)$$

Proof. We have from the defining equation (84) that

$$\begin{aligned} -(X+1)^{i+1} (X-1)^j \frac{dX}{du} &= (X+1) \sum_{n \geq 0} \tilde{C}_n(r, i, j) u^{n-i-j-2}, \\ -(X+1)^i (X-1)^{j+1} \frac{dX}{du} &= (X-1) \sum_{n \geq 0} \tilde{C}_n(r, i, j) u^{n-i-j-2}. \end{aligned}$$

Subtracting these two identities, using again (84), and comparing the coefficients of powers of u gives (88). Similarly, multiplying $(X+1)^i (X-1)^j$ to both sides of the defining equation (83), using (84), and comparing the coefficients of powers of u , we obtain (89). \square

Proposition 3. *We have the following identities: $\forall i, j \in \mathbb{C}$,*

$$f_i(T) f_j(-T) = \sum_{n \geq 0} \left(1 + \frac{n-i-j-1}{r}\right)_n \tilde{C}_n(r, i, j) \left(\frac{rT}{2}\right)^n, \quad (90)$$

where $(s)_n = s(s+1) \cdots (s+n-1)$ denotes the ascending Pochhammer symbol.

Proof. Write

$$f_i(T) f_j(-T) = \sum_{n \geq 0} \left(1 + \frac{n-i-j-1}{r}\right)_n P_{i,j,n}(r) \left(\frac{rT}{2}\right)^n. \quad (91)$$

It then suffices to show that the $P_{i,j,n}(r)$ satisfy the same recursion relation as $\tilde{C}_n(r, i, j)$. This can be verified straightforwardly using (79)–(81). The proposition is proved. \square

Let us now assume that r is a positive integer and use the wave-function-pair approach (see Appendix A) to compute the residue of $L^{k/r}$, where we remind the reader that

$$L = \partial^r + Cx.$$

First we construct a particular pair of wave functions (ψ, ψ^*) of L . Start with solving the equation $L\psi = z^r\psi$, that is,

$$(\partial^r - C(z^r/C - x))\psi = 0. \quad (92)$$

Denote $X = z^r/C - x$. Then we have

$$((-\partial_X)^r - CX)\psi = 0. \quad (93)$$

Using the formal saddle point method, we know that this linear ODE has a unique formal Puiseux series solution of the form:

$$P_1(X) = e^{-\frac{r}{r+1}C^{\frac{1}{r}}X^{\frac{r+1}{r}}} X^{-\frac{r-1}{2r}} \sum_{m \geq 0} \frac{c_m}{X^{\frac{(r+1)m}{r}}}, \quad c_0 := 1. \quad (94)$$

Therefore, any wave function $\psi(x, z)$ for L can be expressed as $\alpha_1(z) P_1(X)$ for some $\alpha_1(z)$. Although as we know that the choice of $\alpha_1(z)$ is not unique, let us show that the following function gives a particular choice:

$$\alpha_1^{\text{bisp}}(z) := \frac{1}{e^{-\frac{r}{r+1}C^{\frac{1}{r}}X^{\frac{r+1}{r}}} X^{-\frac{r-1}{2r}} \Big|_{x=0}} = C^{-\frac{r-1}{2r}} e^{\frac{r}{r+1}C^{-1}z^{r+1}} z^{\frac{r-1}{2}}. \quad (95)$$

Indeed, define

$$\psi = \psi(x, z) := \alpha_1^{\text{bisp}}(z) P_1(X). \quad (96)$$

Then it is easy to see that ψ has the form $\psi = \Phi_1(e^{xz})$, where $\Phi_1 = \sum_{k \geq 0} \phi_{1,k}(x) \partial^{-k}$ with $\phi_{1,0} \equiv 1$. Hence the function ψ constructed by (96) is indeed a wave function of L . We call this choice of $\alpha_1(z)$ the *bispectral* one. Similarly, denote by

$$P_2(X) := e^{\frac{r}{r+1}C^{\frac{1}{r}}X^{\frac{r+1}{r}}} X^{-\frac{r-1}{2r}} \sum_{m \geq 0} \frac{c_m^*}{X^{\frac{(r+1)m}{r}}}, \quad c_0^* := 1 \quad (97)$$

the unique formal solution to the linear ODE

$$(\partial_X^r - CX)\psi^* = 0. \quad (98)$$

Define

$$\alpha_2^{\text{bisp}}(z) := C^{-\frac{r-1}{2r}} e^{-\frac{r}{r+1}C^{-1}z^{r+1}} z^{\frac{r-1}{2}}, \quad (99)$$

and construct

$$\psi^* = \psi^*(x, z) := \alpha_2^{\text{bisp}}(z) P_2(X). \quad (100)$$

Then it is easy to see that ψ^* can be written as $\psi^* = \Phi_2(e^{-xz})$, where $\Phi_2 = \sum_{k \geq 0} \phi_{2,k}(x) \partial^{-k}$ with $\phi_{2,0} \equiv 1$.

Proposition 4. *The above ψ, ψ^* form a particular pair of wave and dual wave functions of L .*

Proof. Since we already know that ψ is a wave function and ψ^* is a dual wave function, we are left to show that ψ, ψ^* form a pair. From the definition it then suffices to show that $\Phi_1 \circ \Phi_2^* \equiv 1$. Note that for any two pseudo-differential operators P, Q , $\text{res}_z P(e^{xz}) Q(e^{-xz}) dz = \text{res} P \circ Q^*$. Taking $P = \partial^i \circ \Phi_1$ and $Q = \Phi_2$ in this identity we find $\text{res} \partial^i \circ \Phi_1 \circ \Phi_2^* = \text{res}_z \partial^i \circ \Phi_1(e^{xz}) \Phi_2(e^{-xz}) dz$. Therefore, showing $\Phi_1 \circ \Phi_2^* \equiv 1$ is further equivalent to showing that for all $i \geq 0$,

$$\text{res}_z \partial^i (\psi(z, x)) \psi^*(z, x) dz = 0. \quad (101)$$

Before continuing the proof let us do preparations. Following [14, 20] (see also [77]), introduce the following linear operators S_z and S_z^* :

$$S_z := \frac{C}{rz^{r-1}} \partial_z - \frac{r-1}{2r} C z^{-r} - z = \frac{C}{r} z^{-\frac{r-1}{2}} \circ \partial_z \circ z^{-\frac{r-1}{2}} - z, \quad (102)$$

$$S_z^* := -\frac{C}{rz^{r-1}} \partial_z + \frac{r-1}{2r} C z^{-r} - z = \frac{C}{r} z^{-\frac{r-1}{2}} \circ (-\partial_z) \circ z^{-\frac{r-1}{2}} - z. \quad (103)$$

Then we have the following lemma.

Lemma 3. *For any $i \geq 0$, we have*

$$\partial^i (\psi(x, z)) = (-S_z)^i (\psi(x, z)), \quad \partial^i (\psi^*(x, z)) = (S_z^*)^i (\psi^*(x, z)). \quad (104)$$

Proof. By straightforward calculations using the definitions (96), (100). \square

As in Appendix A, define

$$c(z) := \psi(0, z), \quad c^*(z) := \psi^*(0, z).$$

We then further have the following lemma.

Lemma 4. *The ψ and ψ^* defined by (96) and (100) have the expressions:*

$$\psi(x, z) = \sum_{i \geq 0} \frac{(-1)^i}{i!} S_z^i (c(z)) x^i = \sum_{i \geq 0} \frac{(xz)^i}{i!} f_i \left(\frac{C/r}{z^{r+1}} \right), \quad (105)$$

$$\psi^*(x, z) = \sum_{i \geq 0} \frac{1}{i!} (S_z^*)^i (c^*(z)) x^i = \sum_{i \geq 0} (-1)^i \frac{(xz)^i}{i!} f_i \left(\frac{-C/r}{z^{r+1}} \right), \quad (106)$$

where we recall that f_i are given by (79).

Proof. Performing the Taylor expansion of ψ with respect to x at $x = 0$ and using Lemma 3 we immediately get the first equality of (105). By using (94) we find that $c(z)$ has the form

$$c(z) = \sum_{m \geq 0} C \frac{(\frac{r+1}{r})^m}{r} \frac{c_m}{z^{(r+1)m}}, \quad c_0 = 1. \quad (107)$$

Define

$$\tilde{f}_i := z^{-i} (-S_z)^i (c(z)).$$

Then by using the definition of S_z we observe that $\tilde{f}_i \in \mathbb{C}[[1/z^{r+1}]]$. Write

$$\tilde{f}_i = \tilde{f}_i(T), \quad T = \frac{C}{r z^{r+1}}.$$

Using again the definition of S_z as well as (92), we find that

$$\begin{aligned}\tilde{f}_{j+1}(T) &= \left(1 + \left(\frac{r-1}{2} - j\right)T + (r+1)T^2 \frac{d}{dT}\right) \tilde{f}_j(T), \\ \tilde{f}_{j+r}(T) &= \tilde{f}_j(T) - rjT \tilde{f}_{j-1}(T).\end{aligned}$$

Comparing these with (80)–(81) and using the uniqueness of the recursion we conclude that $f_i = \tilde{f}_i$. This proves (105). The proof of (106) is similar. The lemma is proved. \square

End of the proof of Proposition 4. Using the above Lemma 4 and (90), we have

$$\begin{aligned}& \operatorname{res}_z \partial_x^i (\psi(z, x)) \psi^*(z, x) dz \\ &= \operatorname{res}_z \sum_{m \geq 0} \frac{z^{m+i} x^m}{m!} f_{m+i} \left(\frac{C/r}{z^{r+1}}\right) \sum_{\ell \geq 0} (-1)^\ell \frac{(xz)^\ell}{\ell!} f_\ell \left(\frac{-C/r}{z^{r+1}}\right) dz \\ &= \operatorname{res}_z \sum_{m, \ell, q \geq 0} (-1)^\ell \frac{z^{m+\ell+i-q(r+1)} x^{m+\ell}}{m! \ell!} \left(1 + \frac{q-i-m-\ell-1}{r}\right)_q \tilde{C}_q(r, m+i, \ell) \left(\frac{r}{2}\right)^q dz = 0.\end{aligned}$$

The proposition is proved. \square

Corollary. *Let $L = \partial^r + Cx$ and $z_k(x)$ be defined by (71). We have*

$$z_k(0) = \begin{cases} (-1)^{(r+1)n} \left(1 + \frac{n-1}{r}\right)_n \tilde{C}_n(r, 0, 0) \frac{C^n}{2^n}, & \text{if } k = -1 + (r+1)n \text{ with } n \geq 0, \\ 0, & \text{otherwise.} \end{cases} \quad (108)$$

Proof. According to (279) and (278), and using Lemma 4, we have

$$\sum_{k \geq -1} (-1)^{k+1} z_k(0) z^{-(k+1)} = f_0 \left(\frac{C/r}{z^{r+1}}\right) f_0 \left(-\frac{C/r}{z^{r+1}}\right). \quad (109)$$

Here $z_{-1}(0) := 1$. Substituting (90) in this equality we find

$$\sum_{n \geq 0} \left(1 + \frac{n-1}{r}\right)_n \frac{\tilde{C}_n(r, 0, 0) C^n}{2^n z^{n(r+1)}} = 1 + \sum_{k \geq 0} (-1)^{k-1} z_k(0) z^{-(k+1)}. \quad (110)$$

The corollary is proved by comparing the coefficients of powers of z^{-1} . \square

5. THE A SERIES

In this section, we will prove Theorem 5, which generalizes all parts of Theorem 2 (except for (i) and (ii), which were generalized to all \mathfrak{g} , not just A_l , in Section 3) from A_4 to A_{r-1} for all r . We will also prove the integrality result Theorem 1, which we have obtained for $r = 5$ only. The odd-looking numbering is meant to correspond to the various parts of Theorem 2.

Theorem 5. *Let $m := \lceil (2g-1)/r \rceil$. The following statements are true:*

(iii) [algebraicity] *Define $\tilde{a}_g(r)$ from the generating function*

$$y(x) = \sum_{g \geq 0} \tilde{a}_g(r) (2x)^{2g} = 1 - \frac{r-1}{6} x^2 + \frac{(r-1)(r-3)(2r+1)}{360} x^4 + \dots, \quad (111)$$

where $y(x)$ is the unique solution in $1 + x^2 \mathbb{Q}[r][[x^2]]$ of the polynomial equation

$$\frac{(y+x)^{r+1} - (y-x)^{r+1}}{2x} - r - 1 = 0. \quad (112)$$

Then for all $g \geq 1$ with $2g - 1 \not\equiv 0 \pmod{r}$,

$$\tau_{A_{r-1}}(g) = \frac{(-1)^{m+g} r^{1-g}}{\left(\left\{\frac{2g-1}{r}\right\}\right)_m} \tilde{a}_g(r). \quad (113)$$

(iv) [closed formula] Define $c_{p,j}(r)$ as in (72). Then for all $g \geq 0$, we have

$$\tau_{A_{r-1}}(g) = \frac{(-1)^{g+m-1} r^{-g}}{\left(\left\{\frac{2g-1}{r}\right\}\right)_{2g+m+1}} \sum_{p=0}^{2g} \left(\frac{2g(r+1)-1}{r}\right)_{p+2g}^{-} c_{p,2g}(r). \quad (114)$$

(v) [product formula] Define $C_n(r, j) \in \mathbb{Q}[r, j]$ by (75)–(77) and $f_j(T)$ by (79), i.e., $f_j(T) = \sum_{k \geq 0} (2k+1)!! C_{2k}(r, j) (-T)^k$. Then the following identity holds true:

$$f_0(T) f_0(-T) = \sum_{g \geq 0} \left(1 - \frac{1-2g}{r}\right)_{2g} (1-2g) r^{2g} \tilde{a}_g(r) T^{2g}. \quad (115)$$

(vi) [terminating hypergeometric sum] Set $R = [r/2]$. For all $g \geq 1$ with $2g - 1 \not\equiv 0 \pmod{r}$, we have

$$\tau_{A_{r-1}}(g) = \frac{(-1)^{g+m}}{\left(\left\{\frac{2g-1}{r}\right\}\right)_m} \frac{r^{1-g}}{1-2g} \sum_{d \geq 0} \binom{\frac{2g-1}{r}}{d} K_{g,d}, \quad (116)$$

where

$$K_{g,d} = \frac{1}{4^g (r+1)^d} \sum_{\substack{m_1+m_2+\dots+m_R=d \\ m_1+2m_2+\dots+Rm_R=g}} \binom{d}{m_1, \dots, m_R} \prod_{i=1}^R \binom{r+1}{2i+1}^{m_i}. \quad (117)$$

(vii) [asymptotics] For $r \geq 3$, as $g \rightarrow \infty$ with $2g - 1 \not\equiv 0 \pmod{r}$, we have

$$\tau_{A_{r-1}}(g) \sim \frac{r \sqrt{\pi}}{\sqrt{r+1}^{\frac{r-2}{r}}} \frac{1}{\Gamma(1 - \left\{\frac{2g-1}{r}\right\})} \frac{1}{\Gamma(\frac{2g-1}{r})} g^{-\frac{3}{2}} \left(4r(r+1)^{\frac{2}{r}} \sin^2\left(\frac{\pi}{r}\right)\right)^{-g}. \quad (118)$$

For $r = 2$ (the A_1 case) the asymptotic formula is the same with an extra factor of $1/2$.

Proof. Let us start from recalling the r -spin Witten conjecture [112] (the Faber–Shadrin–Zvonkine theorem [64]). Introduce the Lax operator

$$L = \partial^r + \sum_{\alpha=1}^{r-1} u_\alpha \partial^{r-1-\alpha}. \quad (119)$$

The Gelfand–Dickey hierarchy [34] is the following infinite family of PDEs for $(r-1)$ unknown functions u_1, \dots, u_{r-1} :

$$\frac{\partial L}{\partial s^{\alpha,q}} = \left[(L^{k/r})_+, L \right], \quad q \geq 0, \quad (120)$$

where $k = \alpha + rq$. Similarly as before, denote $\mathbf{s} := (s^{\alpha,q})_{\alpha=1, \dots, r-1, q \geq 0}$. Since

$$\frac{\partial u_\alpha}{\partial s^{1,0}} = \frac{\partial u_\alpha}{\partial x},$$

we identify $s^{1,0}$ with x . Denote by $Z(\mathbf{t})$ the r -spin partition function defined by (38) with Ω being the Witten class. Then the r -spin Witten conjecture [112] is stated as follows: the partition function $Z = Z(\mathbf{t}(\mathbf{s}))$ is a particular tau-function for the Gelfand–Dickey hierarchy (120), where

$$t^{\alpha,q} = (-1)^{q+1} (\sqrt{-r})^{\frac{3k}{r+1}+1} \left(\frac{\alpha}{r}\right)_{q+1} s^{\alpha,q}, \quad q \geq 0; \quad (121)$$

moreover, the partition function Z satisfies the string equation (44) with $\eta_{\alpha\beta} = \delta_{\alpha+\beta,r}$. Note that the string equation (44) written in terms of the time variables $s^{\alpha,q}$ reads

$$\sum_{\alpha=1}^l \sum_{q \geq 0} (\alpha + rq + r) s^{\alpha,q+1} \frac{\partial Z}{\partial s^{\alpha,q}} + \frac{1}{2} \sum_{\alpha=1}^l \alpha (r - \alpha) s^{\alpha,0} s^{l+1-\alpha,0} Z = \frac{\partial Z}{\partial s^{1,0}}. \quad (122)$$

It is known that the differential polynomials h_k defined by

$$h_k := \frac{(-1)^{q+1}}{(\sqrt{-r})^3 \frac{1+k}{r+1} \left(\frac{\alpha}{r}\right)_{q+1}} \operatorname{res} L^{\frac{k}{r}} \quad (k \in \mathbb{Z}_{>0} \setminus r\mathbb{Z}_{>0}) \quad (123)$$

are the tau-symmetric Hamiltonian densities for the Gelfand–Dickey hierarchy (see [34, 112]; also cf. [14, 23, 24, 57]). Here $k = \alpha + rq$. Denote by $u_\alpha(\mathbf{s})$ the solution corresponding to the particular tau-function $Z(\mathbf{t}(\mathbf{s}))$. The Lax operator L given by (119) is now also subjected to this solution. We have

$$h_k(\mathbf{s}(\mathbf{t})) = \frac{\partial^2 \log Z(\mathbf{t})}{\partial t^{\alpha,q} \partial t^{1,0}}. \quad (124)$$

Then using the string equation together with a Miura-type transformation we find that the initial Lax operator $L|_{s^{\alpha,q}=x \delta^{\alpha,1} \delta^{q,0}}$, still denoted by L , has the following explicit expression [14]:

$$L = \partial^r + r x. \quad (125)$$

Now by using the above (40), (123), (124) we have

$$\tau_{A_{r-1}}(g) = (-1)^{q+g} \frac{z_{2g(r+1)-1}^{(0)}}{r^{3g} \left(\frac{\alpha}{r}\right)_{q+2}}, \quad \forall g \geq 0, \quad (126)$$

where $\alpha \in \{1, \dots, l\}$, $q \geq 0$ are the unique integers satisfying (24), and we recall that $z_k(x) := \operatorname{res} L^{k/r}$. Therefore, by using (108), Proposition 3, and the corollary to Proposition 2 from the previous section, we arrive at parts (iii)–(v). We remark in passing that equation (112) multiplied out, takes the form

$$\sum_{i=0}^{r/2} \binom{r+1}{2i+1} x^{2i} y^{r-2i} - r - 1 = 0, \quad (127)$$

which for $r = 5$ agrees with the equation in (9) (part (iii) of Theorem 2) if we set $z = -20x^2$.

To show (116), we will prove a more general statement (suitable for the later as well). Denote

$$\lambda(p; s) = p^r + \sum_{i=1}^{r-1} s_i p^{r-1-i},$$

where $s = (s_1, \dots, s_l)$. Consider the following algebraic equation for p :

$$\lambda(p; s) = \xi^r. \quad (128)$$

This equation has a unique formal solution $p(\xi; s)$ in $\xi + \mathbb{Q}[s][[\xi^{-1}]]$. Write

$$p(\xi; s) = \xi + \sum_{k \geq 1} u_k(s) \xi^{-k} \quad (129)$$

with $u_k(s) \in \mathbb{Q}[s]$, $k \geq 1$. Differentiating (129) with respect to ξ , we find

$$\frac{dp(\xi; s)}{d\xi} = 1 - \sum_{k \geq 1} k u_k(s) \xi^{-k-1}. \quad (130)$$

Therefore,

$$k u_k(s) = \operatorname{res}_{\xi=\infty} \frac{dp(\xi; s)}{d\xi} \xi^k d\xi = \operatorname{res}_{\xi=\infty} \rho^k dp(\xi; s) = \operatorname{res}_{p=\infty} f(p; s) \frac{k}{r} dp. \quad (131)$$

Noting that

$$\left(p^r + \sum_{i=1}^{r-1} s_i p^{r-1-i} \right)^{\frac{k}{r}} = p^k \sum_{j \geq 0} \binom{\frac{k}{r}}{j} \sum_{i_1, \dots, i_j=1}^{r-1} s_{i_1} \dots s_{i_j} p^{-j-i_1-\dots-i_j}, \quad (132)$$

we obtain the following expressions for $u_k(s)$:

$$k u_k(s) = - \sum_{j \geq 0} \binom{\frac{k}{r}}{j} \sum_{\substack{1 \leq i_1, \dots, i_j \leq r-1 \\ j+i_1+\dots+i_j=k+1}} s_{i_1} \dots s_{i_j} \quad (133)$$

$$= - \sum_{\substack{n_1, \dots, n_{r-1} \geq 0 \\ 2n_1+3n_2+\dots+rn_{r-1}=k+1}} \binom{\frac{k}{r}}{\sum_i n_i} \binom{\sum_i n_i}{n_1, \dots, n_{r-1}} s_1^{n_1} \dots s_{r-1}^{n_{r-1}}. \quad (134)$$

Now specializing the above to $s = s^*$ given by

$$s_{2i}^* = 0, \quad s_{2i-1}^* = \frac{1}{4^i(r+1)} \binom{r+1}{2i+1},$$

we obtain

$$(1-2g) u_{2g-1}(s^*) = \sum_{d \geq 0} \frac{\binom{2g-1}{d}}{4^g (r+1)^d} \sum_{\substack{m_1+\dots+m_R=d \\ m_1+2m_2+\dots+Rm_R=g}} \binom{d}{m_1, \dots, m_R} \prod_{i=1}^R \binom{r+1}{2i+1}^{m_i}. \quad (135)$$

Here $m_i = n_{2i-1}$. Combining with (112) we obtain (116).

It remains to prove part (vii). In view of (113), (118) is equivalent to the asymptotic formula

$$\tilde{a}_g(r) \sim \frac{(r+1)^{\frac{1}{r}-\frac{1}{2}}}{\sqrt{\pi g^3}} \frac{\sin\left(\frac{2g-1}{r}\pi\right)}{\left(-4(r+1)^{\frac{2}{r}} \sin^2\left(\frac{\pi}{r}\right)\right)^g}, \quad r > 2 \quad (136)$$

for the coefficients $\tilde{a}_g(r)$ defined in (111). By a standard principle, this is in turn equivalent to studying the asymptotic properties (to lowest order) of the generating function $y(x)$ near its singularities of smallest absolute value. Let $P = P(x, y)$ be the polynomial on the left-hand side of (112) and denote by P_x and P_y its partial derivatives with respect to x and y , respectively. The singularities of $y(x)$ are located at the points where its graph becomes vertical, i.e., where both P and P_y vanish (and where P_x doesn't vanish, but one sees easily that these three polynomials have no common zeros). Since $P_y/(r+1) = ((y+x)^r - (y-x)^r)/(2x)$, we see that the curve $\{P_y = 0\}$ has $(r-1)$ components, parametrized by the non-trivial r th roots of unity ζ and given by $\frac{y+x}{y-x} = \zeta$. Setting $(x, y) = c\left(\frac{\zeta-1}{2}, \frac{\zeta+1}{2}\right)$ with $c \in \mathbb{C}$, and substituting this into the equation $P(x, y) = 0$, we find that $c^r = r+1$. Thus, setting $c_0 = (r+1)^{1/r}$, we have that the $r(r-1)$ common points of $P = 0$ and $P_y = 0$ are given parametrically by $(x, y) = c_0\left(\frac{\alpha-\beta}{2}, \frac{\alpha+\beta}{2}\right)$ with (α, β) ranging over all pairs of distinct r th roots of unity. Of these, the ones with x nearest to the origin are those with $\alpha/\beta = \zeta_r^{\pm 1}$, where $\zeta_r = e^{2\pi i/r}$, i.e., they are the $2r$ points $(\pm x_j, y_j)$, where $j \in \mathbb{Z}/r\mathbb{Z}$ and $(x_j, y_j) := \frac{c_0}{2}(\zeta_r^j - \zeta_r^{j-1}, \zeta_r^j + \zeta_r^{j-1})$, all of them with $|x_j| = c_0 \sin(\pi/r)$.

To complete the proof, we must (a) compute the potential contribution from each nearest singularity and (b) see that only the four singularities $(\pm x_0, y_0)$ and $(\pm x_1, y_1)$ belong to the subset of $\{P = 0\}$ that is parametrized by $x \mapsto (x, y(x))$ in the closed disk $|x| \leq c_0 \sin(\pi/r)$.

For (a), we note that from the Taylor expansion of P near the singularity (x_j, y_j) we find that for a point $(x, y) = (x_j - \varepsilon, y_j + \delta)$ near (x_j, y_j) lying on the curve $P = 0$ we must have $P_x(x_j, y_j)\varepsilon \approx P_{yy}(x_j, y_j)\delta^2/2$ or $y \approx y_j + C_j \sqrt{1 - x/x_j}$ with $C_j = \pm \sqrt{(2x P_x/P_{yy})(x_j, y_j)}$. A short computation gives that $x_j P_x(x_j, y_j) = r(r+1)$ and $P_{yy}(x_j, y_j) = -\zeta_r^{1-2j} r(r+1)^{2-2/r}$, so $C_j = \pm i \sqrt{2} \zeta_r^{j-\frac{1}{2}} (r+1)^{\frac{1}{r}-\frac{1}{2}}$. The contribution from this singularity to the coefficient $2^{2g} \tilde{a}_g(r)$ of x^{2g} in $y(x)$ is then asymptotically equal to $C_j \binom{1/2}{2g} x_j^{-2g}$, and of course the contribution from $(-x_j, y_j)$ is the same since P is even in X . Since $\binom{1/2}{2g}$ is asymptotically equal to $-1/\sqrt{32\pi g^3}$, we find that the sum of the four contributions coming from $(\pm x_0, y_0)$ and $(\pm x_1, y_1)$ indeed gives the asymptotic formula stated in (136). For (b), we first note that since P is a polynomial in x^2 and y having degree r in y , the map $\pi : (x, y) \mapsto X = x^2$ represents the Riemann surface $\{P = 0\}$ as an r -sheeted branched cover of $\mathbb{P}^1(\mathbb{C})$. By the above calculations, this map is unramified over the open disc $D = \{|X| < |x_j^2| = c_0^2 \sin^2(\pi/r)\}$, but has r ramification points $(X_j = x_j^2, y_j)$, at each of which exactly two of the r sheets come together, over the closure \bar{D} . The inverse image of the disk D consists of r disjoint disks D_ζ , indexed by the r th roots of unity ζ , where D_ζ is the component of $\pi^{-1}(D)$ containing $(0, \zeta)$ and is parametrized by $y = \zeta y(x/\zeta)$ with $x \in D$. If we write D_j ($j \in \mathbb{Z}/r\mathbb{Z}$) for D_{ζ^j} , then it is not hard to see that \bar{D}_j meets \bar{D}_{j-1} at the point (X_j, y_j) and, of course (replacing j by $j+1$) also meets \bar{D}_{j+1} at (X_{j+1}, y_{j+1}) , and that the closed disks \bar{D}_j have no other points in common. In particular, the component D_0 parametrized by $y = y(x)$ contains the two ramification points (X_0, y_0) and (X_1, y_1) in its closure, and no others, and this means that the two points nearest to the origin where the series $y(\sqrt{X})$ is not analytic are X_0 and X_1 , as claimed. For $r = 2$, this proof of the asymptotics also applies, but the situation on how the various branches over the closed disk \bar{D} meet at their boundaries is slightly different, and we get the extra factor $1/2$. This can also be seen directly, since $y = \sqrt{1 - x^2/3}$ in this case, but in any case there is no need to do any of this since the well-known formula $\tau_{A_1}(g) = \frac{1}{24^g g!}$ immediately gives the asymptotics.

This completes the proof of all parts of Theorem 5. \square

Remark 7. A different approach for computing the r -spin intersection numbers, using the theory of matrix models, was obtained by Brézin–Hikami [16, 17]. For the one-point numbers $\tau_{A_{r-1}}(g)$, Brézin–Hikami discovered the following integral formula:

$$\tau_{A_{r-1}}(g) = \frac{(-1)^g r^{1-g}}{1 - \frac{\alpha}{r}} [t^{2g}] \int_0^\infty \exp\left(-\frac{(s+t/2)^{r+1} - (s-t/2)^{r+1}}{(r+1)t}\right) ds, \quad (137)$$

where $g \geq 0$ with $2g - 1 \equiv \alpha \pmod{r}$, $\alpha = 1, \dots, r-1$. We note that formula (116) of part (vi) of Theorem 5 can also be proved by using (137). Indeed, expanding the integrand suitably and integrating term-by-term, one can obtain that

$$\tau_{A_{r-1}}(g) = \frac{(-1)^g r^{-g}}{4^g \Gamma(1 - \{\frac{2g-1}{r}\})} \sum_{d \geq 0} (-1)^d \frac{\Gamma(d - \frac{2g-1}{r})}{(r+1)^d} \sum_{\substack{m_1+m_2+\dots+m_R=d \\ m_1+2m_2+\dots+Rm_R=g}} \prod_{i=1}^R \frac{\binom{r+1}{2i+1}^{m_i}}{m_i!}, \quad (138)$$

which by Euler's formula is equivalent to (116). Liu–Vakil–Xu [86] obtained formula (138) by using the integral formula (137); this was also our original derivation of (116). Therefore, part (vi) of Theorem 5 is not new; however, our current proof for (116) using the wave-function-pair approach from the theory of integrable systems is a self-contained one. We also note that the integral formula (137) in principle could also be used together with the method of steepest descent to give a different proof for part (vii) of Theorem 5. (The recursion given by the dual

topological ODE, cf. Theorem 4, could give a third proof of part (vii) of Theorem 5 with the constant term in the right-hand side of (136) undetermined.)

By using (111)–(113) one immediately gets the first few values of $\tau_{A_{r-1}}(g)$:

$$\tau_{A_{r-1}}(1) = \frac{r-1}{24}, \quad r \geq 2, \quad (139)$$

$$\tau_{A_{r-1}}(2) = \begin{cases} 1/1152, & r = 2, \\ \frac{(r-3)(r-1)(2r+1)}{5760r}, & r \geq 3, \end{cases} \quad (140)$$

$$\tau_{A_{r-1}}(3) = \begin{cases} 1/82944, & r = 2, \\ 1/31104, & r = 3, \\ 3/20480, & r = 4, \\ \frac{(r-5)(r-1)(2r+1)(8r^2-13r-13)}{2903040r^2}, & r \geq 5. \end{cases} \quad (141)$$

Corollary. *For $r \geq 2g$, the value of $\tau_{A_{r-1}}(g)$ is a Laurent polynomial in r . Moreover, the value of this Laurent polynomial at $r = -1$ is equal to $\frac{B_{2g}}{2g}$, where B_n is the n th Bernoulli number.*

Proof. If $r \geq 2g$, then $m = 0$, so the right-hand side of (113) reduces to $-\tilde{a}_g(r)/(-r)^{g-1}$. The first statement then follows from the fact that $\tilde{a}_g(r) \in \mathbb{Q}[r]$. Now by taking the $r \rightarrow -1$ limit in (112) we find the unique solution $y = \frac{x}{\tanh x}$. The second statement then follows. \square

According to Witten [111, 112] the second statement in the corollary should give a new proof of the Harer–Zagier formula [73, 99] on the orbifold Euler characteristic of $\mathcal{M}_{g,1}$ (cf. also [17, 86]).

We now give the proof of the three integrality statements for $\tau_{A_4}(g)$ stated in the introduction.

Proof of Theorem 1. The algebraicity of the generating function of the numbers a_g as given in part (iii) of Theorem 2 implies their integrality away from the primes 2, 3, and 5. To prove the integrality of b_g and c_g defined by equation (5), we use (15). It follows from (15) that

$$c_g = 6^{-g} \sum_{0 \leq s \leq 5g/2} 2^{-2s} (-9)^s c_g^{[s]}, \quad (142)$$

where the numbers $c_g^{[s]}$ are given according to the value of $g \pmod{5}$ by

$$\begin{aligned} c_{5n}^{[s]} &:= \frac{5^{-2s} \binom{3}{5}_n \binom{4}{5}_n \binom{1}{5}_{3n-s}}{s! (5n-2s)!}, & c_{5n-1}^{[s]} &:= \frac{5^{-2s} \binom{2}{5}_n \binom{4}{5}_n \binom{3}{5}_{3n-1-s}}{s! (5n-1-2s)!}, \\ c_{5n-3}^{[s]} &:= \frac{5^{-2s} \binom{3}{5}_n \binom{1}{5}_n \binom{2}{5}_{3n-2-s}}{s! (5n-3-2s)!}, & c_{5n-4}^{[s]} &:= \frac{5^{-2s} \binom{1}{5}_n \binom{2}{5}_n \binom{4}{5}_{3n-3-s}}{s! (5n-4-2s)!}. \end{aligned}$$

To show that c_g belongs to $\mathbb{Z}[1/30]$ we will actually show the stronger statement that each $c_g^{[s]}$ belongs to $\mathbb{Z}[1/5]$.

Consider first the case $g = 5n$. For each prime $p \neq 5$, the p -adic valuation of $c_{5n}^{[s]}$ is given by

$$\nu_p(c_{5n}^{[s]}) = \nu_p\left(\binom{3}{5}_n\right) + \nu_p\left(\binom{4}{5}_n\right) + \nu_p\left(\binom{1}{5}_{3n-s}\right) - \nu_p(s!) - \nu_p((5n-2s)!).$$

Namely,

$$\nu_p(c_{5n}^{[s]}) = \sum_{k \geq 1} \left[u\left(\binom{3}{5}_n, p^k\right) + u\left(\binom{4}{5}_n, p^k\right) + u\left(\binom{1}{5}_{3n-s}, p^k\right) - u(s!, p^k) - u((5n-2s)!, p^k) \right],$$

where $u((a/5)_n, p^k)$ denotes the number of elements in $\{a, a+5, \dots, a+5n-5\}$ that are divisible by p^k ($a = 1, \dots, 5, k \geq 1$). Counting these numbers we find

$$\begin{aligned} u\left(\left(\frac{3}{5}\right)_n, p^k\right) + u\left(\left(\frac{4}{5}\right)_n, p^k\right) + u\left(\left(\frac{1}{5}\right)_{3n-s}, p^k\right) - u(s!, p^k) - u((5n-2s)!, p^k) \\ = f_j\left(\frac{n}{p^k}, \frac{s}{p^k}\right) \quad \text{if } p^k \equiv j \pmod{5}, \quad 1 \leq j \leq 4, \end{aligned} \quad (143)$$

where the functions $f_j : \mathbb{R}^2 \rightarrow \mathbb{Z}$ are defined by

$$f_1(x, y) := \left[x + \frac{2}{5}\right] + \left[x + \frac{1}{5}\right] + \left[3x - y + \frac{4}{5}\right] - [y] - [5x - 2y], \quad (144)$$

$$f_2(x, y) := \left[x + \frac{1}{5}\right] + \left[x + \frac{3}{5}\right] + \left[3x - y + \frac{2}{5}\right] - [y] - [5x - 2y], \quad (145)$$

$$f_3(x, y) := \left[x + \frac{4}{5}\right] + \left[x + \frac{2}{5}\right] + \left[3x - y + \frac{3}{5}\right] - [y] - [5x - 2y], \quad (146)$$

$$f_4(x, y) := \left[x + \frac{3}{5}\right] + \left[x + \frac{4}{5}\right] + \left[3x - y + \frac{1}{5}\right] - [y] - [5x - 2y]. \quad (147)$$

Therefore it suffices to show that each of f_1, f_2, f_3, f_4 is nowhere negative on \mathbb{R}^2 . To show this, we observe that each $f_\alpha(x, y)$ is periodic in both variables and is also piecewise constant, with jumps only along finitely many lines in the unit square $[0, 1]^2$, so one only has to compute the values of $f_\alpha(x, y)$ in each component of the complement of the union of these lines. (The value of f_i at a point lying on one of these lines is always equal to the value of f_i in one of the adjacent open regions.) This is most easily seen graphically, as illustrated in Figure 1, which shows that




FIGURE 1. The values of f_1 and f_2 in $[0, 1]^2$

each of f_1 and f_2 assumes only the values 0, 1, and 2, and this also gives the non-negativity of f_3 and f_4 as well since $f_3(x, y) = 2 - f_2(-x, -y)$ and $f_4(x, y) = 2 - f_1(-x, -y)$.

The proof for each other residue class of $g \pmod{5}$ is exactly similar and again reduces to the non-negativity of four piecewise continuous functions on $\mathbb{R}^2/\mathbb{Z}^2$, but when one goes through the details it turns out that these four are a permutation of the same four functions f_j as above in each case, so that we do not need to make new graphs. The proofs for b_g are also similar, and again a priori involve the non-negativity of 16 periodic functions on \mathbb{R}^2 , which again turn out to be repetitions of only four functions:

$$g_1(x, y) = \left[x + \frac{1}{5}\right] + \left[x + \frac{4}{5}\right] + \left[3x - y + \frac{4}{5}\right] - [y] - [5x - 2y], \quad (148)$$

$$g_2(x, y) = \left[x + \frac{3}{5}\right] + \left[x + \frac{2}{5}\right] + \left[3x - y + \frac{2}{5}\right] - [y] - [5x - 2y], \quad (149)$$

$$g_3(x, y) = \left[x + \frac{2}{5}\right] + \left[x + \frac{3}{5}\right] + \left[3x - y + \frac{3}{5}\right] - [y] - [5x - 2y], \quad (150)$$

$$g_4(x, y) = \left[x + \frac{4}{5}\right] + \left[x + \frac{1}{5}\right] + \left[3x - y + \frac{1}{5}\right] - [y] - [5x - 2y]. \quad (151)$$

Their non-negativity can be proved as for the f_j . The theorem is proved. \square

6. THE D SERIES

In this section, we derive explicit formulas for the numbers $\tau_{\mathfrak{g}}(g)$ with \mathfrak{g} being D_l ($l \geq 4$). Denote by $r = 2l - 2$ the Coxeter number of \mathfrak{g} , and recall that $\tau_{D_l}(g) = \langle \tau_{\alpha, q} \rangle$ are the one-point FJRW invariants of D_l -type, where α, q are determined by (46). Denote by $Z = Z(\mathbf{t})$ the partition function for the FJRW invariants of D_l -type as in (38).

Theorem 6. *Let $l \geq 4$ be an integer and $r = 2l - 2$. Set $m := [(2g - 1)/r]$. Then the following statements are true:*

(iii) [algebraicity] *Define $n_g(r)$ from the generating function*

$$y(t) = \sum_{g \geq 0} n_g(r) t^{2g} = 1 - \frac{r+2}{24} t^2 + \frac{(r+2)(r-6)(2r+1)}{5760} t^4 + \dots, \quad (152)$$

where $y(t)$ is the unique solution in $1 + t^2 \mathbb{Q}[r][[t^2]]$ to the algebraic equation

$$\sum_{j=0}^{r/2} \frac{1}{2j+1} \binom{j+r/2}{2j} y^{r-2j} t^{2j} - 1 = 0 \quad (153)$$

or alternatively t and $y = y(t)$ are related algebraically by

$$t/y = Z - Z^{-1}, \quad t^r \frac{Z^{r+1} - Z^{-r-1}}{r+1} = (Z - Z^{-1})^{r+1}. \quad (154)$$

Then for all $g \geq 1$ we have

$$\tau_{D_l}(g) = \frac{(-1)^{m+g} r^{1-g}}{\left(\left\{\frac{2g-1}{r}\right\}\right)_m} n_g(r). \quad (155)$$

(iv) [closed formula] *For all $g \geq 1$, we have*

$$\tau_{D_l}(g) = \frac{(-1)^{g+m-1} r^{-g}}{\left(\left\{\frac{2g-1}{r}\right\}\right)_{2g+m+1}} \sum_{j=0}^{2g} \sum_{p=0}^j c_{p,j}(r) \left(\frac{2g(r+1)-1}{r}\right)_{2g+p}^{-} \binom{-\frac{1}{2}}{2g-j}. \quad (156)$$

(v) [product formula] *Define $C_n(r, j) \in \mathbb{Q}[r, j]$ by (75)–(77) and $f_j(T)$ by (79), i.e., $f_j(T) = \sum_{k \geq 0} (2k+1)!! C_{2k}(r, j) (-T)^k$. Then the following identity holds true:*

$$\left[f_{\frac{1}{2}}(T) f_{-\frac{1}{2}}(-T) \right]_{\text{even}} = \sum_{g \geq 0} \left(1 - \frac{1-2g}{r}\right)_{2g} (1-2g) r^{2g} n_g(r) T^{2g}. \quad (157)$$

Here for a power series $v(T)$, $[v(T)]_{\text{even}}$ means taking the even degree part of $v(T)$.

(vi) [terminating hypergeometric sum] *For all $g \geq 1$,*

$$\tau_{D_l}(g) = \frac{(-1)^{g+m}}{\left(\left\{\frac{2g-1}{r}\right\}\right)_m} \frac{r^{1-g}}{1-2g} \sum_{d \geq 0} \binom{\frac{2g-1}{r}}{d} K_{g,d}, \quad (158)$$

where

$$K_{g,d} = \sum_{\substack{m_1+m_2+\dots+m_{l-1}=d \\ m_1+2m_2+\dots+(l-1)m_{l-1}=g}} \binom{d}{m_1, \dots, m_{l-1}} \prod_{i=1}^R \left[\frac{\binom{r+i+1}{i+1}}{i+2} \right]^{m_i}. \quad (159)$$

(vii) [asymptotics] *For $l > 4$, as $g \rightarrow \infty$, $\tau_{D_l}(g)$ is given asymptotically by*

$$\tau_{D_l}(g) \sim \frac{r \sqrt{\pi} \cos\left(\frac{\pi}{r}\right)}{\sqrt{r+1} \frac{r-2}{r}} \frac{1}{\Gamma\left(1 - \left\{\frac{2g-1}{r}\right\}\right)} \frac{1}{\Gamma\left(\frac{2g-1}{r}\right)} g^{-\frac{3}{2}} \left(4r(r+1) \frac{2}{r} \sin^2\left(\frac{\pi}{r}\right)\right)^{-g}. \quad (160)$$

For $l = 4$ the formula is the same with an extra factor of 3.

Proof. Recall that the DS hierarchy of D_l -type admits the following scalar Lax operator [38]:

$$L = \partial^r + \partial^{-1} \circ \sum_{\alpha=1}^{l-1} (u^\alpha \circ \partial^{2\alpha-1} + \partial^{2\alpha-1} \circ u^\alpha) + \partial^{-1} \circ u^l \circ \partial^{-1} \circ u^l. \quad (161)$$

For $\alpha = 1, \dots, l-1$, the corresponding DS flows can be written in terms of L by

$$\frac{\partial L}{\partial s^{\alpha,q}} = \left[(L^{k/r})_+, L \right], \quad q \geq 0, \quad (162)$$

where $k = m_\alpha + rq$. Denote $\mathbf{s} := (s^{\alpha,q})_{\alpha=1, \dots, l, q \geq 0}$. For $\alpha = l$, the corresponding flows could also be obtained in terms of L via the so-called negative flows [90, 108] or by definition from the original Drinfeld–Sokolov’s matrix Lax system [38]. Since $\partial u^\alpha / \partial s^{1,0} = \partial u^\alpha / \partial x$, we identify $s^{1,0}$ with x . Witten’s ADE conjecture [65, 67, 112] for the D_l case (the Fan–Jarvis–Ruan theorem [65, 67]; see also [72, 78, 89]) can then be stated as follows: the partition function $Z := Z(\mathbf{t}(\mathbf{s}))$ is a particular tau-function for the DS hierarchy of D_l -type, where

$$t^{\alpha,q} = (-1)^{q+1} (\sqrt{-r})^{\frac{3k}{r+1}+1} \left(\frac{m_\alpha}{r} \right)_{q+1} s^{\alpha,q}, \quad q \geq 0; \quad (163)$$

moreover, $Z(\mathbf{t})$ satisfies the string equation (44) with $\eta_{\alpha\beta}$ in (44) given by

$$(\eta_{\alpha\beta}) = \begin{pmatrix} 0 & \dots & 0 & 1 & 0 \\ \vdots & \ddots & \ddots & 0 & 0 \\ 0 & 1 & \ddots & \ddots & \vdots \\ 1 & 0 & \ddots & \ddots & 0 \\ 0 & 0 & \dots & 0 & 1 \end{pmatrix}. \quad (164)$$

One can alternatively write the string equation (44) using the variables $s^{\alpha,q}$ from (163) as

$$\sum_{\alpha=1}^l \sum_{q \geq 0} (m_\alpha + rq + r) s^{\alpha,q+1} \frac{\partial Z}{\partial s^{\alpha,q}} + \sum_{\alpha=1}^{l-1} \frac{m_\alpha(r - m_\alpha)}{2} s^{\alpha,0} s^{l-\alpha,0} Z + \frac{m_l^2}{2} (s^{l,0})^2 Z = \frac{\partial Z}{\partial s^{1,0}}. \quad (165)$$

Denote by $u^\alpha = u^\alpha(\mathbf{s})$ the solution corresponding to the particular tau-function Z , the so-called topological solution. The Lax operator L given by (161) is now also subjected to this solution. Similarly as in the proof for the A case, we note that

$$h_k := \frac{(-1)^{q+1}}{(\sqrt{-r})^{\frac{3k}{r+1}+1} \left(\frac{m_\alpha}{r} \right)_{q+1}} \operatorname{res} L^{\frac{k}{r}} \quad (k > 0 \text{ odd}) \quad (166)$$

are a part of the tau-symmetric Hamiltonian densities for the DS hierarchy of D_l -type. Here $k = m_\alpha + rq$. Therefore,

$$h_k(\mathbf{s}(\mathbf{t})) = \frac{\partial^2 \log Z(\mathbf{t})}{\partial t^{\alpha,q} \partial t^{1,0}}. \quad (167)$$

The initial Lax operator $L|_{s^{\alpha,q} = x \delta^{\alpha,1} \delta^{q,0}}$ will again be denoted by L . We have the following lemma.

Lemma 5. *We have*

$$L = \partial^r + rx - \frac{1}{2} r \partial^{-1}. \quad (168)$$

Proof. For the topological solution, the corresponding normal coordinates r_α satisfy

$$r_\alpha(\mathbf{t}) = \frac{\partial^2 \log Z(\mathbf{t})}{\partial t^{\alpha,0} \partial t^{1,0}}. \quad (169)$$

Dividing the string equation (44) by $Z(\mathbf{t})$, then differentiating it with respect to $t^{\alpha,0}$, and finally taking $t^{\alpha,q} = t^{1,0} \delta^{\alpha,1} \delta^{q,0}$ we obtain that

$$r_\alpha(\mathbf{t})|_{t^{\alpha,q}=t^{1,0} \delta^{\alpha,1} \delta^{q,0}} = \eta_{\alpha 1} t^{1,0} = \delta_{\alpha, l-1} t^{1,0}. \quad (170)$$

Then, as in [13, 14], by using the Miura-type transformation between the normal coordinates and the u -coordinates together with a degree argument we find that

$$u^\alpha|_{s^{\beta,q}=x \delta^{\beta,1} \delta^{q,0}} = \frac{r}{2} x \delta^{\alpha,1}.$$

The lemma is proved. \square

As before, denote $z_k(x) := \text{res } L^{k/r}$. Using (167), (40) and (46) we obtain that

$$\tau_{D_l}(g) = (-1)^{q+g} \frac{z_{2g(r+1)-1}(0)}{r^{3g} \binom{m_\alpha}{r}_{q+2}}. \quad (171)$$

Lemma 6. *The following formula holds true:*

$$L = \partial^{-\frac{1}{2}} \circ (\partial^r + r x) \circ \partial^{\frac{1}{2}}. \quad (172)$$

Proof. RHS = $\partial^r + r \partial^{-\frac{1}{2}} \circ x \circ \partial^{\frac{1}{2}} =$ LHS. \square

Using (73) and (172), we have

$$\begin{aligned} & \left(\partial^r + r x - \frac{1}{2} r \partial^{-1} \right)^\lambda \\ &= \partial^{-\frac{1}{2}} \circ (\partial^r + r x)^\lambda \circ \partial^{\frac{1}{2}} \\ &= \sum_{\substack{0 \leq p \leq j \\ s \geq 0}} r^{j+s} c_{p,j}(r) (\lambda)_{s+j+p}^- \sum_{v=0}^s \binom{-\frac{1}{2}}{v} \frac{x^{s-v}}{(s-v)!} \partial^{r(\lambda-s)-(r+1)j-v}. \end{aligned}$$

Combining with (171) we obtain (156).

To prove (153), let us use the wave-function-pair approach (see Appendix A and Section 4). We first construct a particular pair of wave and dual wave functions for L . Start with solving $L\psi = z^r \psi$, i.e.,

$$\left(\partial^r + (r x - z^r) - \frac{r}{2} \partial^{-1} \right) \psi = 0. \quad (173)$$

Denote $X = z^r/r - x$. We have

$$\left((-\partial_X)^r - r X + \frac{r}{2} \partial_X^{-1} \right) \psi = 0. \quad (174)$$

Similarly as in Section 4 we find that this equation has a unique formal solution of the form

$$P_1(X) = e^{-\frac{1}{r+1} r \frac{r+1}{r} X \frac{r+1}{r}} X^{-\frac{1}{2}} \sum_{m \geq 0} \frac{c_m}{X \frac{(r+1)m}{r}}, \quad c_0 := 1. \quad (175)$$

Therefore, $\psi(x, z) = \alpha_1(z) P_1(X)$ for some $\alpha_1(z)$ to be determined. The choice of $\alpha_1(z)$ is not unique. We will use the particular choice of $\alpha_1(z)$ given by

$$\alpha_1^{\text{bisp}}(z) := \frac{1}{e^{-\frac{r}{r+1} r \frac{1}{r} X \frac{r+1}{r}} X^{-\frac{1}{2}}}|_{x=0} = r^{-\frac{1}{2}} e^{\frac{r}{r+1} C^{-1} z^{r+1}} z^{\frac{r}{2}}. \quad (176)$$

We call this particular choice the *bispectral* one. Namely, we construct

$$\psi = \psi(x, z) := \alpha_1^{\text{bisp}}(z) P_1(X). \quad (177)$$

Similarly, denote by

$$P_2(X) := e^{\frac{r}{r+1} X^{\frac{1}{r}}} X^{-\frac{1}{2} + \frac{1}{r}} \sum_{m \geq 0} \frac{c_m^*}{X^{\frac{(r+1)m}{r}}}, \quad c_0^* := 1 \quad (178)$$

the unique formal solution to the following linear equation:

$$\left(\partial_X^r - rX - \frac{r}{2} \partial_X^{-1} \right) \psi^* = 0. \quad (179)$$

Define

$$\alpha_2^{\text{bisp}}(z) := r^{-\frac{1}{2} + \frac{1}{r}} e^{-\frac{1}{r+1} z^{r+1}} z^{\frac{r}{2} - 1}, \quad (180)$$

and construct

$$\psi^* = \psi^*(x, z) := \alpha_2^{\text{bisp}}(z) P_2(X). \quad (181)$$

Proposition 5. *The ψ, ψ^* form a particular pair of wave and dual wave functions of L .*

Proof. It is easy to verify that ψ is a wave function and ψ^* is a dual wave function for L . Write $\psi = \Phi_1(e^{xz})$ and $\psi^* = \Phi_2(e^{-xz})$, where

$$\Phi_1 = \sum_{k \geq 0} \phi_{1,k}(x) \partial^{-k}, \quad \Phi_2 = \sum_{k \geq 0} \phi_{2,k}(x) \partial^{-k}$$

with $\phi_{1,0} = \phi_{2,0} \equiv 1$. To show ψ and ψ^* form a pair, it suffices to show $\Phi_1 \circ \Phi_2^* = 1$. Similarly as before, we know that this is equivalent to show that for all $i \geq 0$,

$$\text{res}_z \partial^i (\psi(z, x)) \psi^*(z, x) dz = 0. \quad (182)$$

Before continuing the proof of the proposition let us prove two lemmas.

Lemma 7. *Introduce two linear operators R_z and R_z^* :*

$$R_z = \frac{1}{z^{r-1}} \partial_z - \frac{r}{2} z^{-r} - z, \quad R_z^* = -\frac{1}{z^{r-1}} \partial_z + \frac{r-2}{2} z^{-r} - z. \quad (183)$$

Then we have for any $i \geq 0$,

$$\partial_x^i (\psi(x, z)) = (-R_z)^i (\psi(x, z)), \quad \partial_x^i (\psi^*(x, z)) = (R_z^*)^i (\psi^*(x, z)). \quad (184)$$

Proof. By direct calculations. \square

Lemma 8. *The ψ and ψ^* have the following expressions:*

$$\psi(x, z) = \sum_{i \geq 0} \frac{(-1)^i}{i!} R_z^i(c(z)) x^i = \sum_{i \geq 0} \frac{(xz)^i}{i!} f_{i-1/2}\left(\frac{1}{z^{r+1}}\right), \quad (185)$$

$$\psi^*(x, z) = \sum_{i \geq 0} \frac{1}{i!} (R_z^*)^i(c^*(z)) x^i = \sum_{i \geq 0} (-1)^i \frac{(xz)^i}{i!} f_{i+1/2}\left(\frac{-1}{z^{r+1}}\right). \quad (186)$$

Here, $c(z) := \psi(0, z)$, $c^*(z) := \psi^*(0, z)$, and $f_{i \pm 1/2}$ are given by (79).

Proof. Doing the Taylor expansion of ψ with respect to x at $x = 0$ and using Lemma 7 we immediately get the first equality in (185). From (175) we find that $c(z)$ has the form

$$c(z) = \sum_{m \geq 0} r^{\frac{(r+1)m}{r}} \frac{c_m}{z^{(r+1)m}}, \quad c_0 = 1. \quad (187)$$

Define

$$\tilde{f}_i := z^{-i} (-R_z)^i (c(z)).$$

From the definition of S_z one easily finds that $\tilde{f}_i \in \mathbb{C}[[z^{-(r+1)}]]$. Write

$$\tilde{f}_i = \tilde{f}_i(T), \quad T = \frac{1}{z^{r+1}},$$

and we find from the definition of R_z and (173) that

$$\begin{aligned} \tilde{f}_{j+1}(T) &= \left(1 + \left(\frac{r}{2} - j\right)T + (r+1)T^2 \frac{d}{dT}\right) \tilde{f}_j(T), \\ \tilde{f}_{j+r}(T) &= \tilde{f}_j(T) - r\left(j - \frac{1}{2}\right)T \tilde{f}_{j-1}(T). \end{aligned}$$

Comparing with (80)–(81) and using the uniqueness argument that is similar to the one given after the proof of Lemma 1, we conclude that $\tilde{f}_i = f_{i-1/2}$. This proves (185). The proof of (186) is similar. The lemma is proved. \square

End of the proof of Proposition 5. Using Lemma 8 and identity (90) we have

$$\begin{aligned} &\text{res}_z \partial^i (\psi(z, x)) \psi^*(z, x) dz \\ &= \text{res}_z \sum_{m \geq 0} \frac{z^{m+i} x^m}{m!} f_{m+i-1/2} \left(\frac{1}{z^{r+1}}\right) \sum_{\ell \geq 0} (-1)^\ell \frac{(xz)^\ell}{\ell!} f_{\ell+1/2} \left(\frac{-1}{z^{r+1}}\right) dz \\ &= \text{res}_z \sum_{m, \ell, q \geq 0} (-1)^\ell \frac{z^{m+\ell+i-q(r+1)} x^{m+\ell}}{m! \ell!} \left(1 + \frac{q-i-m-\ell-1}{r}\right)_q \tilde{C}_q \left(r, m+i-\frac{1}{2}, \ell+\frac{1}{2}\right) \left(\frac{r}{2}\right)^q dz \\ &= 0. \end{aligned}$$

Proposition 5 is proved. \square

Using formulas (171), (278), (279), Proposition 5, Lemma 8 and Proposition 3 we obtain that

$$\tau_{D_l}(g) = \frac{(-1)^{g+m-1} \tilde{C}_{\frac{1}{2}, -\frac{1}{2}, 2g}^{(r)}}{\left(\left\{\frac{2g-1}{r}\right\}\right)_{m+1}} \frac{1}{2^{2g} r^{3g}}. \quad (188)$$

Lemma 9. *For any $g \geq 0$, we have*

$$n_g = -\frac{\tilde{C}_{2g} \left(r, \frac{1}{2}, -\frac{1}{2}\right)}{2^{2g} (2g-1)}. \quad (189)$$

Proof. From the definition of $\tilde{C}_n(r, i, j)$ and the fact that X is an odd Laurent series in u^{-1} one easily obtains the equality

$$\sum_{g \geq 0} \tilde{C}_{2g} \left(r, \frac{1}{2}, -\frac{1}{2}\right) u^{2g} = -\frac{u^2}{2} \left(\sqrt{\frac{X+1}{X-1}} + \sqrt{\frac{X-1}{X+1}} \right) \frac{dX}{du} = \frac{-u^2 X}{\sqrt{X^2-1}} \frac{dX}{du}. \quad (190)$$

Then it suffices to show that the series P defined by

$$P := \sum_{g \geq 0} \tilde{C}_{2g} \left(r, \frac{1}{2}, -\frac{1}{2}\right) \frac{u^{2g}}{1-2g} = 1 - \frac{r+2}{6} u^2 + \dots \quad (191)$$

satisfies

$$\sum_{j \geq 0} \binom{j+r/2}{2j} \frac{(2u)^{2j} P^{r-2j}}{2j+1} = 1. \quad (192)$$

(Note that since in our case $r = 2l - 2$ is a non-negative even integer, the above sum terminates at $j = l - 1$ and expresses P as an algebraic function of u , but even if r is a complex or formal variable the identity makes sense and will be shown to be true.) Comparing (190) and (191) we find

$$\frac{d}{dX} \left(\frac{P}{u} \right) = \frac{d(P/u)/du}{dX/du} = \frac{X}{\sqrt{X^2 - 1}}.$$

Therefore,

$$P = u\sqrt{X^2 - 1}.$$

Combining with (83) it suffices to show

$$\sum_{j \geq 0} \frac{2^{2j}}{2j+1} \binom{j+r/2}{2j} (X^2 - 1)^{r/2-j} = \frac{(X+1)^{r+1} - (X-1)^{r+1}}{2(r+1)},$$

which is an elementary exercise (also equivalent to (154)). □

Identity (157) follows directly from (188). Formulas (188) and (189) yield (155). Formulas (155) and (134) yield (158).

Finally, the proof of the asymptotic formula (160) is very similar to that of the corresponding statement in Theorem 5, and will only be sketched. In view of the relation (155) we see that it is equivalent to the asymptotic formula

$$n_g(r) \sim \cos\left(\frac{\pi}{r}\right) \frac{(r+1)^{\frac{1}{r}-\frac{1}{2}}}{\sqrt{\pi g^3}} \frac{\sin((2g-1)\frac{\pi}{r})}{(-4(r+1)^{\frac{2}{r}} \sin^2(\frac{\pi}{r}))^g} \quad (l > 4) \quad (193)$$

for the coefficients $n_g(r)$ defined in (152). Writing (153) in the form $(\frac{Z^2}{Z^2-1})^{r+1} - (\frac{1}{Z^2-1})^{r+1} = \frac{r+1}{t^r}$ and setting the derivative of this expression with respect to Z equal to 0, we find that Z^2 is a non-trivial r th root of unity at all singularities of $y = y(t)$, and substituting this back into (153) we find that the singularities are given by $(t, Z^2) = ((r+1)^{1/r}(\alpha - \beta), \alpha/\beta)$ where α and β are distinct r th roots of unity. In particular, the singular points of $y(t)$ of smallest absolute value are equal to $2(r+1)^{1/r} \sin(\pi/r)$ times $(2r)$ th roots of unity. The rest of the proof is exactly along the lines of the proof of part (vii) of Theorem 5 and will be omitted. Just as in the A case for $l = 1$, the proof has to be modified slightly if $l = 4$ because the way that the sheets above the closed disk $\{|t| \leq 2(r+1)^{1/r} \sin(\pi/r)\}$ meet at their boundary is slightly different from what happens for $l > 4$; we leave the details as an exercise.

This completes the proof of all parts of Theorem 6. □

Remark 8. It seems worth observing that the factors 1/2 and 3 appearing in part (vii) of Theorems 5 and 6 respectively are related to the symmetries of the corresponding Dynkin diagrams. Indeed, $|\text{Sym}(A_l)| = 2$ for $l > 1$, 1 for $l = 1$; $|\text{Sym}(D_l)| = 2$ for $l > 4$, 6 for $l = 4$.

We note that the polynomial P on the left-hand side of (153) also has the following expression:

$$P = \frac{2y^{r+1} \sinh\left[(r+1) \operatorname{arcsinh}\left(\frac{t}{2y}\right)\right]}{(r+1)t} - 1. \quad (194)$$

Using the formulas (152)–(155) one can compute the first few values of $\tau_{D_l}(g)$:

$$\tau_{D_l}(1) = \frac{r+2}{24}, \quad (195)$$

$$\tau_{D_l}(2) = \frac{(r+2)(r-6)(2r+1)}{5760r}, \quad (196)$$

$$\tau_{D_l}(3) = \frac{(r+2)(2r+1)(8r^3 - 77r^2 + 196r + 188)}{2903040r^2}. \quad (197)$$

Here we note that $r \geq 6$ (as $l \geq 4$). For $r = 6$, our formulas agree with the explicit computations in [12]. More precisely, the explicit expression of the dual topological ODE of D_4 -type was computed in [12], which reduces to a second order ODE for ϕ_3 (in our current notation):

$$108x^2\phi_3'' - (104x^8 + 108x)\phi_3' - (4x^{14} + 260x^7 + 39)\phi_3 = 0. \quad (198)$$

One could then give an alternative proof of Theorem 6 for D_4 by using (198). We leave this as an exercise because in the next section we will prove the algebricity for E_6 in this way.

Similarly as in the A case, we have the following corollary.

Corollary. *For all $r \geq 2g$, the value of $\tau_{D_l}(g)$ is a Laurent polynomial in r . Moreover, the value of this Laurent polynomial at $r = -1$ is equal to $\frac{(1-2^{1-2g})B_{2g}}{2g}$.*

Proof. If $r \geq 2g$, then $m = 0$, so the right-hand side of (155) reduces to $-n_g(r)/(-r)^{g-1}$. The first statement then follows from the fact that $n_g(r) \in \mathbb{Q}[r]$. By taking the $r \rightarrow -1$ limit in (154) we find the unique solution $y = \frac{t/2}{\sinh(t/2)}$, which yields the second statement of the corollary. \square

It might be of interest to see whether the coefficients or the values of the Laurent polynomials in r occurring in the above corollary and the corollary to Theorem 5 have any topological meaning.

7. THE E_6 CASE

In the proof of Theorems 5 and 6 for the A_l, D_l cases in the previous two sections, we used the equivalent scalar Lax representation of the corresponding DS hierarchy [38]. For the E_6 case, as far as we know, the existence of such a representation is an open question. However, following Proposition 1, we can get the higher-genera one-point invariants by computing the dual topological ODE of E_6 -type. (Of course, for any fixed l , as it was mentioned above (see (198)), we could alternatively have used the corresponding dual topological ODE to give a different proof of Theorems 5–6; this deserves a further study.)

Let us now compute the dual topological ODE of E_6 -type. We use the 27-dimensional representation [49] of $\mathfrak{g} = E_6$. Recall that the Coxeter number and the exponents for this case read as follows:

$$r = 12, \quad m_1 = 1, \quad m_2 = 4, \quad m_3 = 5, \quad m_4 = 7, \quad m_5 = 8, \quad m_6 = 11,$$

and that the dimension of this simple Lie algebra is 78. Denote

$$\begin{aligned} X_1 &= E_{6,7} + E_{8,9} + E_{10,11} + E_{12,14} + E_{15,17} + E_{26,27}, \\ X_2 &= E_{4,5} + E_{6,8} + E_{7,9} - E_{18,20} - E_{21,22} - E_{23,24}, \\ X_3 &= E_{4,6} + E_{5,8} + E_{11,13} + E_{14,16} + E_{17,19} + E_{25,26}, \\ X_4 &= E_{3,4} - E_{8,10} - E_{9,11} - E_{16,18} - E_{19,21} + E_{24,25}, \\ X_5 &= E_{2,3} - E_{10,12} - E_{11,14} - E_{13,16} + E_{21,23} + E_{22,24}, \\ X_6 &= E_{1,2} + E_{12,15} + E_{14,17} + E_{16,19} + E_{18,21} + E_{20,22}. \end{aligned}$$

Kostant's sl_2 -subalgebra of \mathfrak{g} can be given by

$$I^- = 16X_1^T + 22X_2^T + 30X_3^T + 42X_4^T + 30X_5^T + 16X_6^T, \quad (199)$$

$$I^+ = X_1 + X_2 + X_3 + X_4 + X_5 + X_6, \quad (200)$$

$$\rho^\vee = \frac{1}{2}(I^+I^- - I^-I^+). \quad (201)$$

Using the above data one can compute explicitly the dual topological ODE of E_6 -type, i.e., equation (52) for $G \in \mathfrak{g}$, or equivalently (54) for the vector $\phi = (\phi_1, \dots, \phi_6) \in \mathbb{C}^6$. We denote by $(\phi_{\alpha;\beta})_{\beta=1,\dots,6}$ ($\alpha = 1, \dots, 6$) the six linearly independent vector solutions that were introduced in Section 3. (Here β labels the components of the vector.) Recall that each $\phi_{\alpha;6}$ has the form (56). Following the principle of Theorem 4, namely, by reducing the ODE (54) for the vector-valued function ϕ to a scalar ODE for the top component ϕ_6 (the highest weight vector of \mathfrak{g}), we find the following fourth-order one:

$$\begin{aligned} 0 = & 2985984x^4(37x^{39} - 2775x^{26} - 36960x^{13} + 11520)\phi_6'''' \\ & - 466560x^3(2331x^{52} - 162985x^{39} - 2985600x^{26} - 4951296x^{13} + 811008)\phi_6''' \\ & - 27x^2(6545189x^{65} - 1276342935x^{52} + 10115971680x^{39} - 127523831040x^{26} \\ & \quad + 860446310400x^{13} - 52110950400)\phi_6'' \\ & - 27x(23310x^{78} - 8293439x^{65} - 3559160940x^{52} - 153887586840x^{39} \\ & \quad + 1228034776320x^{26} + 236111616000x^{13} + 49235558400)\phi_6' \\ & + (37x^{91} - 3464310x^{78} + 2278737540x^{65} + 114309996390x^{52} \\ & \quad + 10889113435200x^{39} - 60840963615600x^{26} \\ & \quad - 15770999462400x^{13} - 328914432000)\phi_6. \end{aligned} \quad (202)$$

This means that every $\phi_{\alpha;6}$ must satisfy (202). From the discussion in Section 3, we know that each $\phi_{\alpha;6}$ has the form

$$\phi_{\alpha;6} = x^{1+\frac{13}{12}m_\alpha} f_\alpha(x^{13}) \quad (203)$$

for some power series $f_\alpha(u) \in \mathbb{Q}[[u]]$ (resp. $u^{-1}\mathbb{Q}[[u]]$ for $\alpha = 6$). But if we use the Frobenius method, then we find that the indicial equation for (202) at $x = 0$ has only four roots $25/12$, $77/12$, $103/12$, $-1/12$. This implies that $\phi_{2;6}$ and $\phi_{5;6}$ must both vanish (which was not obvious from their original definition), while the expansions of the other $\phi_{\alpha;6}$, if we normalize to make the power series monic, are given by (203) with $1 + \frac{13}{12}m_\alpha = \frac{26\alpha-1}{12}$ and

$$\begin{aligned} f_1(u) &= 1 + \frac{4235}{2^9 3^1 13} u + \frac{23102233}{2^{18} 3^2 13^2} u^2 + \frac{381109489145}{2^{29} 3^3 13^3} u^3 + \dots, \\ f_3(u) &= 1 + \frac{4613}{2^{10} 13} u + \frac{340813583}{2^{19} 3^2 5^1 13^2} u^2 + \frac{1468738987769}{2^{28} 3^2 13^3 17} u^3 + \dots, \\ f_4(u) &= 1 + \frac{34829}{2^8 5^1 7^1 13} u + \frac{112497481}{2^{20} 3^2 13^2} u^2 + \frac{45611422760339}{2^{28} 3^2 5^1 7^1 13^3 19} u^3 + \dots, \\ u f_6(u) &= 1 + \frac{435}{2^8 13} u + \frac{330276383}{2^{19} 3^2 11^1 13^2} u^2 + \frac{7178883185}{2^{27} 3^1 13^3} u^3 + \dots. \end{aligned}$$

According to Proposition 1, the numbers $\tau_{E_6}(g)$ with $g \geq 1$ in the particular normalizations that we are using, can be expressed in terms of $\phi_{\alpha;6}$ as follows: for $g \not\equiv 2 \pmod{3}$,

$$\tau_{E_6}(g) = \frac{c_\alpha}{2^{6m} 3^{4m}} [u^m](f_\alpha), \quad (204)$$

where $\alpha \in \{1, 3, 4, 6\}$ and m are determined by $2g - 1 = m_\alpha + 12m$ and

$$c_1 = \frac{1}{4}, \quad c_3 = \frac{5}{1152}, \quad c_4 = \frac{25}{27648}, \quad c_6 = \frac{1}{2^6 3^4}; \quad (205)$$

otherwise $\tau_{E_6}(g) = 0$. For simplicity we set $\tau_{E_6}(0) = 1$ which also agrees with (204). It should be noted that here the freedom of normalizations is fixed in such a way that it agrees with the explicit Frobenius manifold potential of E_6 -type given by Klemm–Theisen–Schmidt [80] (see (243) below) as well as with the genus 1 formula of Dubrovin–Zhang [55] (cf. [49, 67]). For the reader's convenience, let us list the first few values of $\tau_{E_6}(g)$ in the following table:

g	0	1	2	3	4	5	6	7	8	9	10
$\tau_{E_6}(g)$	1	$\frac{1}{4}$	0	$\frac{5}{1152}$	$\frac{25}{27648}$	0	$\frac{145}{5750784}$	$\frac{4235}{414056448}$	0	$\frac{23065}{79498838016}$	$\frac{174145}{3338951196672}$

One-point FJRW invariants of E_6 -type

Theorem 7. Set $B = 2^{1/12}(3 + 2\sqrt{3})^{1/4}$. Define A_k , $k \geq -1$ from the generating series

$$U(V) = \frac{1}{B} \sum_{k \geq -1} A_k V^{-k/12},$$

where $U(V)$ is the unique solution in

$$\frac{1}{B} V^{1/12} + \mathbb{C}[[V^{-1/12}]]$$

to the polynomial equation

$$\begin{aligned} & U^{24} - 36U^{22} + 540U^{20} - 4488U^{18} + 22992U^{16} - 76032U^{14} \\ & + \left(5V + \frac{2140032}{13}\right)U^{12} - 2\left(19V + \frac{1501248}{13}\right)U^{10} + 108\left(V + \frac{24320}{13}\right)U^8 \\ & - 32\left(5V + \frac{41104}{13}\right)U^6 + 32\left(5V + \frac{9696}{13}\right)U^4 - 96\left(V + \frac{192}{13}\right)U^2 \\ & - \frac{1}{108}\left(V^2 - \frac{34560}{13}V - \frac{442368}{169}\right) = 0, \end{aligned} \quad (206)$$

of degree 12 in U^2 and 2 in V . Then for all $g \geq 0$ with $g \not\equiv 2 \pmod{3}$ we have

$$\tau_{E_6}(g) = \psi_\rho 2^{-\frac{13}{6}g} 3^{-\frac{7}{6}g} \frac{A_{2g-1}}{\left(\left\{\frac{2g-1}{12}\right\}\right)_m}, \quad (207)$$

where $\rho \in \{1, 5, 7, 11\}$ and m are such that $2g - 1 = \rho + 12m$, and ψ_ρ are given by

$$\psi_1 = 2^{\frac{1}{2}} 3^{\frac{5}{12}}, \quad \psi_5 = 2^1 3^{\frac{3}{4}} \varepsilon_3^{-\frac{1}{2}}, \quad \psi_7 = -2^{\frac{3}{2}} 3^{\frac{2}{3}} \varepsilon_3^{-\frac{1}{2}}, \quad \psi_{11} = -12, \quad (208)$$

where $\varepsilon_3 = 2 + \sqrt{3}$.

Proof. The differential equation of ϕ_6 translates into recursions for the coefficients of f_α with $\alpha = 1, 3, 4, 6$. On the other hand, any algebraic function satisfies a linear differential equation and hence its coefficients satisfy a recursion. One verifies by computer that the recursions and the initial conditions agree. \square

The first few values of A_k are given by

$$A_{-1} = 1, \quad A_1 = 2^{-\frac{1}{3}} 3^{\frac{3}{4}}, \quad A_5 = 2^{-\frac{3}{2}} 3^{\frac{3}{4}} 5^1 \varepsilon_3^{\frac{1}{2}}, \quad A_7 = -2^{-\frac{17}{6}} 3^1 5^2 \varepsilon_3^{\frac{1}{2}}. \quad (209)$$

In general, the numbers A_k are (up to fractional powers of 2, 3 and ε_3) algebraic integers belonging to $\mathbb{Q}(\sqrt{3})$, so that Theorem 7 implies an integrality statement for the $\tau_{E_6}(g)$ similar to

the first one in Theorem 1. Another consequence of the theorem is an asymptotic formula for the numbers $\tau_{E_6}(g)$ similar to the ones we found for the A and D cases:

Corollary. *As $g \rightarrow \infty$ with $g \not\equiv 2 \pmod{3}$, we have*

$$\tau_{E_6}(g) \sim 13^{-5/12} \sqrt{\pi} \frac{\theta_\rho}{\Gamma(1 - \{\frac{2g-1}{12}\})} \frac{1}{\Gamma(\frac{2g-1}{12})} g^{-\frac{3}{2}} \left(\frac{\sqrt{3+2\sqrt{3}}}{2^1 3^{\frac{7}{6}} 13^{\frac{1}{6}}} \right)^g, \quad (210)$$

where $\rho \in \{1, 5, 7, 11\}$ and m are such that $2g - 1 = \rho + 12m$, and θ_ρ are constants defined by

$$\theta_1 = 2^1 3^{\frac{5}{12}} (1 + \sqrt{3}), \quad \theta_5 = 2^2 3^{\frac{3}{4}}, \quad \theta_7 = 2^{\frac{5}{2}} 3^{\frac{2}{3}}, \quad \theta_{11} = 2^{\frac{5}{2}} 3^1 (1 + \sqrt{3}). \quad (211)$$

Proof. Using (207) we find that it suffices to show that as $g \rightarrow \infty$ with $g \not\equiv 2 \pmod{3}$,

$$A_{2g-1} \sim \frac{13^{-5/12}}{\sqrt{\pi}} \chi_\rho Q^{g/6} g^{-3/2} \quad (212)$$

with $Q = 2^7 (3 + 2\sqrt{3})^3 / 13$ and with

$$\chi_1 = -\chi_{11} = 1, \quad \chi_5 = -\chi_7 = \varepsilon_3. \quad (213)$$

The corollary can then be proved again by using the similar argument as for the A case. \square

Remark 9. We expect that Theorem 7 (and its corollary) will have an analogue for E_7 and E_8 , and actually also for all non-simply-laced simple Lie algebras.

8. EXPLICIT RELATION FOR FJRW INVARIANTS OF \mathfrak{g} -TYPE

In this section, based on the results in the previous sections and on the theory of Frobenius manifolds (see [43, 46, 47, 74, 93, 109]), we generalize Theorem 3 from A_4 to A_l , D_l and E_6 .

According to Kontsevich–Manin [83, 93] (cf. also Appendix B), a homogeneous CohFT of charge d gives a formal Frobenius manifold of charge d with the formal Frobenius potential $F = F(v)$ given by (31). Denote by B the domain of convergence of F . Let \mathfrak{g} be a simply-laced simple Lie algebra of rank l . As before, denote by $\Omega_{g,n}$ the FJRW CohFT of \mathfrak{g} -type and by r the Coxeter number. It follows from the quasi-homogeneity (42) that the formal Frobenius potential F is a polynomial. Therefore, the convergence domain B is the whole of \mathbb{C}^l . We call the Frobenius manifold B constructed from $\Omega_{g,n}$ *the Frobenius manifold of \mathfrak{g} -type*. Note that the Frobenius structure on B given by (307)–(308) can alternatively be constructed from the orbit space of the Coxeter group of \mathfrak{g} -type ([43, 45, 104, 122]), or else from the miniversal deformation of a simple singularity of \mathfrak{g} -type ([22, 36, 37, 46, 74, 102]). The charge and the spectrum of B are

$$d = \frac{r-2}{r}, \quad \mu_\alpha = \frac{m_\alpha}{r} - \frac{1}{2}, \quad R = 0, \quad (214)$$

and the Euler vector field of B is

$$E = \sum_{\alpha=1}^l \frac{r+1-m_\alpha}{r} v^\alpha \partial_\alpha. \quad (215)$$

Before stating and proving the generalized theorem, we first prove a useful lemma.

Lemma 10. *If \mathfrak{g} is a simply-laced simple Lie algebra, there exists a unique choice of calibration $\{\theta_{\alpha,m}\}_{m \geq 0}$ for the Frobenius manifold of \mathfrak{g} -type.*

Proof. The existence of $\theta_{\alpha,m}$ that satisfy (301)–(306) is known. (For the meaning of calibration and the proof of the existence see Appendix B.) We only need to prove the uniqueness. First of all, observing that set of the differences $\{\mu_\alpha - \mu_\beta\}$ does not contain positive integers, we know from [43, 46] that equations (290)–(291) and (295)–(297) determine $\Theta(v; z)$ uniquely. Therefore, $\theta_{\alpha,m}$ are uniquely determined by (298) and (299) up to constants only, i.e., if $\{\tilde{\theta}_{\alpha,m}\}_{m \geq 0}$ also satisfies (301)–(304), then $\tilde{\theta}_{\alpha,m}(v) - \theta_{\alpha,m}(v) = b_{\alpha,m}$ with $b_{\alpha,m}$ being constants. The uniqueness then follows immediately from (306). The lemma is proved. \square

Lemma 10 is true for the Frobenius manifold associated to any finite Coxeter group with the same proof as above.

The following theorem gives the main result of this section.

Theorem 8. *Let \mathfrak{g} be a simple Lie algebra of type A_l , D_l , or E_6 , and let $(B, \langle \cdot, \cdot \rangle, \cdot)$ denote the Frobenius manifold of \mathfrak{g} -type. Take $v = (v_1, \dots, v_l)$ the flat coordinates on B and $(\theta_{\alpha,m})_{\alpha=1, \dots, l, m \geq 0}$ the associated unique calibration. Denote $v_\alpha^* = \langle \tau_{\alpha, m_\alpha} \rangle$. Then the following identity holds:*

$$\langle \tau_{\alpha, m_\alpha + (r+1)m} \rangle = \theta_{\alpha, m}(v^*), \quad \forall m \geq 0. \quad (216)$$

Here, we recall that $\langle \tau_{\alpha, q} \rangle$ denote the one-point FJRW invariants of \mathfrak{g} -type.

Proof. Let us prove this theorem case by case from A to E .

Start with the A_l case. Consider the Frobenius manifold of A_l -type. In this case, $r = l + 1$ is the Coxeter number. Define

$$\lambda = \lambda(p; s) = p^{l+1} + \sum_{\beta=1}^l s_\beta p^{l-\beta}. \quad (217)$$

Here $s = (s_1, \dots, s_l) \in \mathbb{C}^l$. The reader who is familiar with the theory of Frobenius manifolds recognizes λ as the superpotential, and the Frobenius structure on B could be defined by the standard formulas [43, 46, 47] (cf. also [36, 60, 61, 102, 103]) using the superpotential under some carefully chosen normalization factors, and the flat coordinates v are related to s by

$$v^\alpha = -\frac{(\sqrt{-r})^{\frac{3\alpha}{r+1}-1}}{r-\alpha} \operatorname{res}_{p=\infty} \lambda^{\frac{r-\alpha}{r}} dp. \quad (218)$$

It is not difficult to verify that $\det\left(\frac{\partial v^\alpha}{\partial s_\beta}\right) \neq 0$, so the map $s \mapsto v$ given by (218) is indeed an invertible coordinate transformation. By a degree argument, this coordinate change (218) actually has the triangular nature.

Lemma 11. *For the A_l case, the unique functions $\theta_{\alpha,m}(v)$, $m \geq 0$ have the following expressions:*

$$\theta_{\alpha,m}(v) = -\frac{(-1)^{m+1}}{(\sqrt{-r})^{\frac{3}{r+1} + \frac{\alpha+rm}{r}} \left(\frac{\alpha}{r}\right)_{m+1}} \operatorname{res}_{p=\infty} \lambda^{\frac{\alpha+rm}{r}} dp. \quad (219)$$

Proof. We prove this lemma by showing that the right-hand side of (219), denoted by $\tilde{\theta}_{\alpha,m}$, satisfies the defining conditions (301)–(306). The conditions (304) with $z = 0$ and (301) are known to hold true. Indeed, they follow from the Laplace-type transform between the deformed flat coordinates and λ -periods of the Frobenius manifold, with λ being the corresponding superpotential [4, 36, 37, 43, 46, 61, 62, 102, 103] and with the careful choosing of the rescaling factor to match with Witten's normalization of times [112]. Here, the definitions of the product and the invariant flat metric can be found *ibid.*, and we also normalize the metric such that $\eta_{\alpha\beta} = \delta_{\alpha+\beta, l+1}$. The equality (305) can be easily proved by computing the residues, which also

implies (303). Since equality (304) is a rewritting of (297), we know that (304) is actually a consequence of (301)–(303). To show (302), introduce the following extended Euler operator:

$$\tilde{E} := E + \frac{1}{r} p \frac{\partial}{\partial p}, \quad (220)$$

where E in this case is given by (215) with $m_\alpha = \alpha$. From (217) and (218) we obtain

$$\tilde{E} \lambda = \lambda. \quad (221)$$

The required quasi-homogeneity (302) for the gradients of $\theta_{\alpha,m}$ then holds true for the gradients of $\tilde{\theta}_{\alpha,m}$. It remains to show (306). Observe from the definition (219)–(221) that the $\tilde{\theta}_{\alpha,m}$ themselves satisfy the following quasi-homogeneity condition:

$$E \theta_{\alpha,m} = \left(m + \mu_\alpha + \frac{1}{r} + \frac{1}{2} \right) \theta_{\alpha,m}, \quad \forall m \geq 0. \quad (222)$$

Hence the $\tilde{\theta}_{\alpha,m}$, $m \geq 0$, which are polynomials of v , do not contain constant terms. This gives (306). The lemma is proved. \square

We are ready to show (216). For $m = 0$, the validity of (216) is obvious, because from the definition we know that $\theta_{\alpha,0} = v_\alpha$. Let us now prove the validity of (216) for an arbitrary $m \geq 0$. Following [43, 46, 47], consider the polynomial equation

$$\lambda(p; s) = \xi^r. \quad (223)$$

We know that equation (223) has a unique solution $p = p(\xi; s)$ in $\xi + \mathbb{C}[s][[\xi^{-1}]]$. Write

$$p(\xi; s) = \xi + \sum_{k \geq 1} u_k(s) \xi^{-k}, \quad (224)$$

where $u_k(s) \in \mathbb{Q}[s]$, $k \geq 1$. Using similar arguments to those in the previous sections we have

$$k u_k(s) = \operatorname{res}_{p=\infty} \lambda(p; s)^{\frac{k}{r}} dp. \quad (225)$$

Comparing this equality with (219) we obtain that

$$\theta_{\alpha,m}(v) = - \frac{(-1)^{m+1} (\alpha + mr)}{(\sqrt{-r})^{3 \frac{1+\alpha+rm}{r+1}} \left(\frac{\alpha}{r}\right)_{m+1}} u_{\alpha+mr}(s), \quad \forall m \geq 0. \quad (226)$$

Now define

$$s_\beta^* = \begin{cases} \binom{r}{2i} \frac{a^{2i}}{1+2i}, & \beta = 2i - 1 \ (i = 1, \dots, [r/2]); \\ 0, & \text{otherwise.} \end{cases} \quad (227)$$

And we restrict our discussion to the particular point $s = s^*$ on B . Here

$$a := \sqrt{-r}^{\frac{2-r}{r+1}} / 2. \quad (228)$$

From (217), (223) and (227), we know that the series $p(\xi; s^*)$ satisfies

$$\sum_{i=0}^{[r/2]} \binom{r}{2i} a^{2i} \frac{p(\xi; s^*)^{r-2i}}{1+2i} = \xi^r.$$

Comparing this equation with formula (111)–(113) (cf. (127)) we get

$$\tau_{A_{r-1}}(g) = \frac{(-1)^{m+g} r^{1-g}}{2^{2g} a^{2g} \left(\frac{\alpha}{r}\right)_m} u_{2g-1},$$

where g, α, m are related by $2g - 1 = \alpha + rm$. Substituting (226) in this equality and using (24) and (228), we obtain (216). This completes the proof for the A case.

We continue to prove the statement for the D case. In this case, the Coxeter number $r = 2l - 2$. Similarly as in the A case, introduce the superpotential [36]:

$$\lambda = p^r + \sum_{\beta=1}^{l-1} s_{\beta} p^{r-2\beta} + s_l^2 p^{-2}. \quad (229)$$

The flat coordinates in our normalization are given by

$$v^{\alpha} := -\frac{(\sqrt{-r})^{\frac{3m_{\alpha}-1}{r+1}}}{r-m_{\alpha}} \operatorname{res}_{p=\infty} \lambda^{\frac{r-m_{\alpha}}{r}} dp, \quad \alpha = 1, \dots, l-1 \quad (230)$$

and

$$v^l := -\frac{2\sqrt{l-1}}{\sqrt{-r}^{\frac{3}{2} \frac{r+2}{r+1}}} s_l. \quad (231)$$

It is not difficult to verify again that $\det(\partial v^{\alpha} / \partial s_{\beta}) \neq 0$, and the map $s \mapsto v$ given by (230)–(231) indeed gives an invertible coordinate transformation.

Lemma 12. *The unique functions $\theta_{\alpha,m}(v)$, $m \geq 0$ have the following expressions:*

$$\theta_{\alpha,m}(v) = -\frac{(-1)^{m+1}}{(\sqrt{-r})^{3\frac{1+m_{\alpha}+rm}{r+1}} \left(\frac{m_{\alpha}}{r}\right)_{m+1}} \operatorname{res}_{p=\infty} \lambda^{\frac{m_{\alpha}}{r}+m} dp, \quad \alpha = 1, \dots, l-1, \quad (232)$$

$$\theta_{l,m}(v) = \frac{(-1)^{m+1} \sqrt{l-1}}{(\sqrt{-r})^{3\frac{l+rm}{r+1}} \left(\frac{1}{2}\right)_{m+1}} \operatorname{res}_{p=0} \lambda^{\frac{1}{2}+m} dp. \quad (233)$$

Proof. The proof is almost identical with that of Lemma 11, so we only give some brief indications here. Note that the invariant metric η in our normalization is given by

$$\eta = \begin{pmatrix} 0 & \dots & 0 & 1 & 0 \\ \vdots & \ddots & \ddots & 0 & 0 \\ 0 & \ddots & \ddots & \ddots & \vdots \\ 1 & 0 & \ddots & \ddots & 0 \\ 0 & 0 & \dots & 0 & 1 \end{pmatrix}.$$

The extension of the Euler vector field is again given by (220). The lemma is proved. \square

Following [43, 46, 47], we consider the following polynomial equation:

$$\lambda(p; s) = \xi^r. \quad (234)$$

It is easy to see that (234) has a unique solution $p = p^+(\xi; s)$ in $\xi + \mathbb{C}[s][[\xi^{-1}]]$ as well as a unique solution $p = p^-(\xi; s)$ in $s_l \xi^{-r/2} + \xi^{-(r+2)/2} \mathbb{C}[s][[\xi^{-1}]]$. Write

$$p^+(\xi; s) = \xi + \sum_{k \geq 1} u_k(s) \xi^{-k}, \quad p^-(\xi; s) = \sum_{m \geq 0} v_m(s) \xi^{-m-\frac{r}{2}}. \quad (235)$$

Here $v_0(s) = s_l$. Differentiating both equations in (235) with respect to ξ , we find

$$\frac{dp^+}{d\xi} = 1 - \sum_{k \geq 1} k u_k(s) \xi^{-k-1}, \quad \frac{dp^-}{d\xi} = - \sum_{m \geq 0} \left(m + \frac{r}{2}\right) v_m(s) \xi^{-m-\frac{r}{2}-1}. \quad (236)$$

Therefore,

$$ku_k(s) = \operatorname{res}_{\xi=\infty} \frac{dp^+(\xi; s)}{d\xi} \xi^k d\xi = \operatorname{res}_{p=\infty} \lambda(p; s)^{\frac{k}{r}} dp, \quad (237)$$

$$\left(m + \frac{r}{2}\right)v_m(s) = \operatorname{res}_{\xi=\infty} \frac{dp^-(\xi; s)}{d\xi} \xi^{rm+\frac{r}{2}} d\xi = -\operatorname{res}_{p=0} \lambda(p; s)^{m+\frac{1}{2}} dp. \quad (238)$$

Comparing (237) and (238) with (232) and (233), respectively, we have for $\alpha = 1, \dots, l-1$,

$$\theta_{\alpha, m}(v) = -\frac{(-1)^{m+1}(m_\alpha + mr)}{(\sqrt{-r})^3 \frac{1+m_\alpha+rm}{r+1} \left(\frac{m_\alpha}{r}\right)_{m+1}} u_{m_\alpha+mr}(s), \quad \forall m \geq 0, \quad (239)$$

and for $\alpha = l$,

$$\theta_{l, m}(v) = -\sqrt{l-1} \frac{(-1)^{m+1}(l-1+mr)}{(\sqrt{-r})^3 \frac{l+rm}{r+1} \left(\frac{1}{2}\right)_{m+1}} v_m(s), \quad \forall m \geq 0. \quad (240)$$

Define

$$s_\beta^* = \begin{cases} \binom{l-1+\beta}{2\beta} \frac{a^{2\beta}}{1+2\beta}, & \beta = 1, \dots, l-1, \\ 0, & \text{otherwise,} \end{cases} \quad (241)$$

and we now restrict the discussion to the particular point $s = s^*$ on B . Here

$$a := \sqrt{-r} \frac{2-r}{r+1}. \quad (242)$$

From (229), (234) and (241), we know that the series $p^+(\xi; s^*)$ and $p^-(\xi; s^*)$ satisfy

$$\sum_{i=0}^{l-1} \binom{i+l-1}{2i} \frac{a^{2i}}{1+2i} p^+(\xi; s^*)^{r-2i} = \xi^r, \quad p^-(\xi; s^*) = 0.$$

Comparing with (152)–(155) we get

$$\tau_{D_l}(g) = \frac{(-1)^{m+g} r^{1-g}}{a^{2g} \left(\frac{m_\alpha}{r}\right)_m} u_{2g-1},$$

where g is related with α, m by $2g-1 = m_\alpha + rm$. Substituting (239) in this identity and using (46), and (228), we find (216). This completes the proof for the D case.

Finally, let us prove the statement for the E_6 case. Let $B = \mathbb{C}^6$ be the Frobenius manifold of E_6 -type. The spectrum data (μ, R) are now given by

$$\mu_1 = -\frac{5}{12}, \quad \mu_2 = -\frac{1}{6}, \quad \mu_3 = -\frac{1}{12}, \quad \mu_4 = \frac{1}{12}, \quad \mu_5 = \frac{1}{6}, \quad \mu_6 = \frac{5}{12}, \quad R = 0.$$

According to Klemm–Theisen–Schmidt [80], the Frobenius potential has the explicit expression:

$$\begin{aligned}
F = & \frac{1}{2} (v^1)^2 v^6 + v^1 v^2 v^5 + v^1 v^3 v^4 + \frac{1}{2} (v^2)^2 v^3 \\
& + \frac{1}{2} (v^2)^2 v^4 v^6 + \frac{1}{2} v^2 (v^4)^2 v^5 + \frac{1}{6} (v^3)^3 v^6 + \frac{1}{4} (v^3)^2 (v^5)^2 \\
& + \frac{1}{2} v^2 v^3 v^5 (v^6)^2 + \frac{1}{6} v^2 (v^5)^3 v^6 + \frac{1}{2} v^3 v^4 (v^5)^2 v^6 + \frac{1}{12} v^4 (v^5)^4 + \frac{1}{12} (v^4)^4 v^6 \\
& + \frac{1}{6} v^2 v^4 v^5 (v^6)^3 + \frac{1}{24} (v^2)^2 (v^6)^4 + \frac{1}{4} (v^4)^2 (v^5)^2 (v^6)^2 + \frac{1}{6} v^3 (v^4)^2 (v^6)^3 \\
& + \frac{1}{60} (v^3)^2 (v^6)^5 + \frac{1}{24} v^4 (v^5)^2 (v^6)^5 + \frac{1}{24} v^3 (v^5)^2 (v^6)^4 + \frac{1}{24} (v^5)^4 (v^6)^3 \\
& + \frac{1}{252} (v^4)^2 (v^6)^7 + \frac{1}{576} (v^5)^2 (v^6)^8 + \frac{(v^6)^{13}}{185328}. \tag{243}
\end{aligned}$$

Here, $(v^1, \dots, v^6) \in B$ is a system of flat coordinates satisfying that ∂_1 is the identity vector field. The invariant flat metric and the Euler vector field in this coordinate system are given by $\eta_{\alpha\beta} = \delta_{\alpha+\beta,7}$, and

$$E = v^1 \partial_1 + \frac{3}{4} v^2 \partial_2 + \frac{2}{3} v^3 \partial_3 + \frac{1}{2} v^4 \partial_4 + \frac{5}{12} v^5 \partial_5 + \frac{1}{6} v^6 \partial_6. \tag{244}$$

The first few values of the unique calibration $\theta_{\alpha,p}$ for B are

$$\begin{aligned}
\theta_{\alpha,0} &= v_\alpha, \\
\theta_{1,1} &= v^1 v^6 + v^3 v^4 + v^2 v^5, \\
\theta_{2,1} &= v^1 v^5 + v^2 v^3 + v^2 v^4 v^6 + \frac{1}{2} (v^4)^2 v^5 \\
&\quad + \frac{1}{2} v^3 v^5 (v^6)^2 + \frac{1}{6} v^4 v^5 (v^6)^3 + \frac{1}{6} (v^5)^3 v^6 + \frac{1}{12} v^2 (v^6)^4.
\end{aligned}$$

To prove (216), similarly as in the A and D cases, we will use the corresponding superpotential. Following Eguchi–Yang [63] (cf. [19, 81, 85]), introduce three polynomials Q_1, P_1, P_2 :

$$\begin{aligned}
Q_1 = & 270 p^{15} + (171 + 57\sqrt{3}) t_{10} p^{13} + (54 + 27\sqrt{3}) t_{10}^2 p^{11} \\
& + (126 + 84\sqrt{3}) t_7 p^{10} + \left(\left(\frac{35}{4} + \frac{175}{36}\sqrt{3} \right) t_{10}^3 + (144 + 72\sqrt{3}) t_6 \right) p^9 \\
& + \left(\frac{135}{2} + \frac{81}{2}\sqrt{3} \right) t_7 t_{10} p^8 \\
& + \left(\left(\frac{225}{4} + \frac{125}{4}\sqrt{3} \right) t_6 t_{10} + \left(\frac{345}{384} + \frac{35}{96}\sqrt{3} \right) t_{10}^4 + (135 + 81\sqrt{3}) t_4 \right) p^7 \\
& + \left((126 + 72\sqrt{3}) t_3 + \left(10 + \frac{35}{6}\sqrt{3} \right) t_7 t_{10}^2 \right) p^6 \\
& + \left(\left(\frac{63}{4} + 9\sqrt{3} \right) t_7^2 + (36 + 21\sqrt{3}) t_4 t_{10} + \left(\frac{11}{768} + \frac{19\sqrt{3}}{2304} \right) t_{10}^5 + \left(\frac{21}{4} + 3\sqrt{3} \right) t_6 t_{10}^2 \right) p^5 \\
& + \left(\left(\frac{33}{2} + \frac{19}{2}\sqrt{3} \right) t_3 t_{10} + \left(\frac{19}{48} + \frac{11}{48}\sqrt{3} \right) t_7 t_{10}^3 + (24 + 14\sqrt{3}) t_6 t_7 \right) p^4 \\
& - \left(\frac{11}{8} + \frac{19}{24}\sqrt{3} \right) t_7^2 t_{10} p^3 + \left(\frac{45}{4} + \frac{13}{2}\sqrt{3} \right) t_3 t_7 p + \left(\frac{5}{8} + \frac{13}{36}\sqrt{3} \right) t_7^3, \tag{245}
\end{aligned}$$

$$P_1 = 78 p^{10} + (30 + 10\sqrt{3}) t_{10} p^8 + \left(\frac{14}{3} + \frac{7}{3}\sqrt{3} \right) t_{10}^2 p^6 + \left(\frac{33}{2} + 11\sqrt{3} \right) t_7 p^5$$

$$\begin{aligned}
 & + \left(\left(\frac{1}{4} + \frac{5}{36}\sqrt{3} \right) t_{10}^3 + (16 + 8\sqrt{3}) t_6 \right) p^4 + \left(\frac{25}{12} + \frac{5}{4}\sqrt{3} \right) t_7 t_{10} p^3 \\
 & + \left((5 + 3\sqrt{3}) t_4 + \left(\frac{7}{3456} + \frac{1}{864}\sqrt{3} \right) t_{10}^4 + \left(\frac{3}{4} + \frac{5}{12}\sqrt{3} \right) t_6 t_{10} \right) p^2 \\
 & - \left(\frac{7}{2} + 2\sqrt{3} \right) t_3 p - \left(\frac{7}{12} + \frac{1}{3}\sqrt{3} \right) t_7^2, \tag{246}
 \end{aligned}$$

$$\begin{aligned}
 P_2 & = 12p^{10} + (6 + 2\sqrt{3}) t_{10} p^8 + \left(\frac{4}{3} + \frac{2}{3}\sqrt{3} \right) t_{10}^2 p^6 + (6 + 4\sqrt{3}) t_7 p^5 \\
 & + \left((8 + 4\sqrt{3}) t_6 + \left(\frac{1}{8} + \frac{5}{72}\sqrt{3} \right) t_{10}^3 \right) p^4 + \left(\frac{5}{3} + \sqrt{3} \right) t_7 t_{10} p^3 \\
 & + \left((10 + 6\sqrt{3}) t_4 + \left(\frac{7}{1728} + \frac{1}{432}\sqrt{3} \right) t_{10}^4 + \left(\frac{3}{2} + \frac{5}{6}\sqrt{3} \right) t_6 t_{10} \right) p^2 \\
 & + (14 + 8\sqrt{3}) t_3 p + \left(\frac{7}{12} + \frac{1}{3}\sqrt{3} \right) t_7^2, \tag{247}
 \end{aligned}$$

and define

$$\lambda(p; t) = \frac{1}{270 + 156\sqrt{3}} \left(-u_0 + \frac{Q_1 + P_1\sqrt{P_2}}{p^3} \right), \tag{248}$$

where

$$\begin{aligned}
 u_0 & = -(270 + 156\sqrt{3}) t_0 - \frac{1}{16} (19 + 11\sqrt{3}) t_4 t_{10}^2 - \frac{1}{576} (33 + 19\sqrt{3}) t_6 t_{10}^3 \\
 & - \frac{1}{4} (21 + 12\sqrt{3}) t_6^2 - \frac{1}{16} (33 + 19\sqrt{3}) t_{10} t_7^2. \tag{249}
 \end{aligned}$$

Here, $(t_0, t_3, t_4, t_6, t_7, t_{10})$ gives the flat coordinates of Eguchi–Yang, which relate to the flat coordinates (v^1, \dots, v^6) of Klemm–Theisen–Schmidt via a rescaling:

$$v^1 = \kappa_6 t_0, \quad v^2 = \kappa_5 t_3, \quad v^3 = \kappa_4 t_4, \quad v^4 = \kappa_3 t_6, \quad v^5 = \kappa_2 t_7, \quad v^6 = \kappa_1 t_{10}, \tag{250}$$

where

$$\begin{aligned}
 \kappa_1 & = -\frac{1}{2\sqrt{3}}, \quad \kappa_2 = (\sqrt{3} - 1)^{\frac{1}{2}}, \quad \kappa_3 = -(\sqrt{3} - 1), \\
 \kappa_4 & = 2(\sqrt{3} - 1), \quad \kappa_5 = -2\sqrt{3}(\sqrt{3} - 1)^{\frac{1}{2}}, \quad \kappa_6 = 8\sqrt{3}.
 \end{aligned}$$

Lemma 13. *We have*

$$\theta_{\alpha, m}(v) = -\kappa_\alpha \frac{(8\sqrt{3})^m}{\left(\frac{m_\alpha}{12}\right)_{m+1}} \operatorname{res}_{p=\infty} \lambda^{\frac{m_\alpha+12m}{12}} dp, \quad \alpha = 1, \dots, 6, m \geq 0. \tag{251}$$

Proof. The conditions (304) with $z = 0$ and (301) have been verified by Eguchi–Yang [63], where we use different normalization constants κ_α from the ones in [63] to agree with Klemm–Theisen–Schmidt’s normalization for the product and the metric. The proof is again similar to that of Lemma 11. For this case, the extended Euler operator is again defined by

$$\tilde{E} := E + \frac{1}{r} p \frac{\partial}{\partial p}, \tag{252}$$

where E is given by (244). We have $\tilde{E}\lambda = \lambda$ and the quasi-homogeneity (302) for the gradients of $\theta_{\alpha, m}$ follows. The condition (306) follows from the quasi-homogeneity (just as (222)) of $\theta_{\alpha, m}$. The lemma is proved. \square

Following [43, 46, 47], we consider the polynomial equation

$$\lambda(p; t) = \xi^{12}. \tag{253}$$

It is easy to see that (253) has a unique solution $p = p(\xi; t)$ in $\xi + \mathbb{C}[t][[\xi^{-1}]]$. Write

$$p(\xi; t) = \xi + \sum_{k \geq 1} u_k(t) \xi^{-k}. \quad (254)$$

By using the same arguments as before we have

$$k u_k(t) = \operatorname{res}_{p=\infty} \lambda(p; t) \frac{k}{i^2} dp. \quad (255)$$

Comparing (255) with (251) we obtain

$$\theta_{\alpha, m}(v) = -\kappa_{\alpha} (8\sqrt{3})^m \frac{m_{\alpha} + 12m}{\binom{m_{\alpha}}{12}_{m+1}} u_{m_{\alpha}+12m}(t), \quad \forall m \geq 0. \quad (256)$$

Now define $(v^2)^* = (v^5)^* = 0$ and

$$(v^1)^* = \frac{145}{5750784}, \quad (v^3)^* = \frac{25}{27648}, \quad (v^4)^* = \frac{5}{1152}, \quad (v^6)^* = \frac{1}{4}, \quad (257)$$

and restrict the discussion to the particular point $v = v^*$ on B , which corresponds $t = t^*$. From (248), (253) we know that the series $p = p(\xi; t^*)$ satisfies

$$\frac{1}{270 + 156\sqrt{3}} \left(-u_0^* + \frac{Q_1^* + P_1^* \sqrt{P_2^*}}{p^3} \right) = \xi^{12}, \quad (258)$$

where $Q_1^*, P_1^*, P_2^*, u_0^*$ are respectively Q_1, P_1, P_2, u_0 evaluated at $t = t^*$. Theorem 8 is proved by simplifying (258). \square

Observe that on the left-hand side of (216), $g = \frac{m_{\alpha} + rm + 1}{2}$. So if $m_{\alpha} + rm + 1$ is an odd number, then as a direct consequence of (216) the value $\theta_{\alpha, m}(v^*)$ must vanish.

We also have the following corollary.

Corollary. *For a FJRW CohFT of \mathfrak{g} -type with \mathfrak{g} being A_l, D_l , or E_6 , we have*

$$\langle \tau_{\alpha, m_{\alpha} + (r+1)m} \rangle = \sum_{n \geq 0} \sum_{\substack{1 \leq \alpha_1, \dots, \alpha_n \leq l \\ 1 \leq \beta_1, \dots, \beta_n \leq l}} \frac{\langle \tau_{\alpha, m-1} \tau_{\alpha_1, 0} \dots \tau_{\alpha_n, 0} \rangle_0}{n!} \prod_{i=1}^n \eta^{\alpha_i \beta_i} \langle \tau_{\beta_i, m_{\beta_i}} \rangle, \quad \forall m \geq 1. \quad (259)$$

Proof. Following [43], consider the principal integrable hierarchy associated to B :

$$\frac{\partial v^{\alpha}}{\partial t^{\beta, q}} = \eta^{\alpha \gamma} \partial_{t^{1,0}} \left(\frac{\partial \theta_{\beta, q+1}}{\partial v^{\gamma}} \right), \quad q \geq 0 \quad (260)$$

and its initial value problem (viewing $t^{1,0}$ as the space variable) with the initial data:

$$v^{\alpha} |_{t^{\beta, q} = t^{1,0} \delta^{\beta, 1} \delta^{q, 0}} = \delta^{\alpha, 1} t^{1,0}. \quad (261)$$

The defining equations (301)–(306) for $\theta_{\beta, q}$ and the axioms of Frobenius manifolds imply that the equations in (260) all commute, and thus the above initial value problem has a unique solution in $\mathbb{C}[[\mathbf{t}]]^{\otimes l}$, called the topological solution to the principal hierarchy, denoted by $v^{\text{top}}(\mathbf{t}) = (v^{\text{top}, 1}(\mathbf{t}), \dots, v^{\text{top}, l}(\mathbf{t}))$. It is shown in [43] that the following identity holds true:

$$\frac{\partial^2 \mathcal{F}_0(\mathbf{t})}{\partial t^{1,0} \partial t^{\beta, m}} = \theta_{\beta, m}(v^{\text{top}}(\mathbf{t})). \quad (262)$$

Taking $t^{\gamma, p} = t^{\gamma, 0} \delta^{p, 0}$ on both sides of this identity, and noticing that

$$v^{\text{top}, \alpha}(\mathbf{t}) |_{t^{\gamma, p} = t^{\gamma, 0} \delta^{p, 0}} = t^{\alpha, 0}, \quad (263)$$

we obtain

$$\frac{\partial^2 \mathcal{F}_0(\mathbf{t})}{\partial t^{1,0} \partial t^{\beta,m}} \Big|_{t^{\gamma,p}=t^{\gamma,0} \delta_{p,0}} = \theta_{\beta,m}(t^{1,0}, \dots, t^{l,0}). \quad (264)$$

On the other hand, the genus zero part of the string equation (44) reads as follows:

$$\sum_{p \geq 1} t^{\alpha,p} \frac{\partial \mathcal{F}_0(\mathbf{t})}{\partial t^{\alpha,p-1}} + \frac{1}{2} \eta_{\rho\sigma} t^{\rho,0} t^{\sigma,0} = \frac{\partial \mathcal{F}_0(\mathbf{t})}{\partial t^{1,0}}. \quad (265)$$

For $m \geq 1$, by differentiating (265) with respect to $t^{\beta,m}$ and taking $t^{\gamma,q} = t^{\gamma,0} \delta^{q,0}$ we get

$$\frac{\partial \mathcal{F}_0(\mathbf{t})}{\partial t^{\beta,m-1}} \Big|_{t^{\gamma,p}=t^{\gamma,0} \delta_{p,0}} = \frac{\partial^2 \mathcal{F}_0(\mathbf{t})}{\partial t^{1,0} \partial t^{\beta,m}} \Big|_{t^{\gamma,p}=t^{\gamma,0} \delta_{p,0}}. \quad (266)$$

Using the definition (37) for $\mathcal{F}_0(\mathbf{t})$, identities (264) and (266), as well as identity (216), we obtain (259). \square

Note that due to (42) the sum “ $\sum_{n \geq 0}$ ” in the right-hand side of (259) is actually a finite sum. We also note that identity (259) can be written alternatively as follows: for all $g \geq (r+2)/2$,

$$\tau_{\mathfrak{g}}(g) = \sum_{n \geq 0} \sum_{\substack{1 \leq \alpha_1, \dots, \alpha_n \leq l \\ 1 \leq \beta_1, \dots, \beta_n \leq l}} \frac{\langle \tau_{\alpha, m-1} \tau_{\alpha_1, 0} \dots \tau_{\alpha_n, 0} \rangle_0}{n!} \prod_{i=1}^n \eta^{\alpha_i \beta_i} \langle \tau_{\beta_i, m_{\beta_i}} \rangle, \quad (267)$$

where α, m are such that $2g - 1 = m_{\alpha} + rm$.

Remark 10. For $\mathfrak{g} = E_7$ or E_8 , we also expect the validity of identity (216) (or (259), (267)). It might be possible to get proofs in these cases by applying the constructions of the λ -periods for the orbit space of the Coxeter group of \mathfrak{g} -type ([43, 44, 45, 104]) (or for the simple singularity of \mathfrak{g} -type [4, 10, 46, 72, 74, 92, 102, 103, 105]) as well as their Laplace-type transforms [43, 46, 47] to compute the $\theta_{\alpha,m}$, and then matching these with the left-hand side $\tau_{\mathfrak{g}}(g)$ of (216), which can be read off from the coefficients of the top components $\phi_{\alpha;l}$ of the fundamental solutions to the dual topological ODE of \mathfrak{g} -type. Moreover, since we know that the λ -periods for the Frobenius manifold of \mathfrak{g} -type are algebraic, this method of proof, if it works, would also lead to algebraicity (as already mentioned in Remark 9) and therefore integrality of the renormalized numbers of $\tau_{\mathfrak{g}}(g)$. When \mathfrak{g} is a non-simply-laced simple Lie algebra, in order for identity (216) (or (259), (267)) to remain valid, one may need to use the notion of the *partial CohFT* [89]. We hope to study these cases and other more general situations (semisimple or nonsemisimple CohFTs including the cases with the Novikov ring mentioned in Remark 5; cf. e.g. [19, 43, 51, 56, 121]) in later work.

Remark 11. We also observe that the particular point v^* on the Frobenius manifold of A -type that we use for $l \geq 2$ is different from the particular semisimple point used by Pandharipande, Pixton and Zvonkine [97, 98] for obtaining relations in the cohomology ring of $\overline{\mathcal{M}}_{g,n}$. It might be interesting to see whether their method can be applied also to our v^* to give further information connected with the topology of $\overline{\mathcal{M}}_{g,n}$.

We provide a few examples to illustrate the results of this section.

Example 1 (A_1). The superpotential reads as follows:

$$\lambda = p^2 + s_1, \quad v^1 = \frac{s_1}{2}.$$

We have $\eta = 1$ and $F = (v^1)^3/6$. For $m \geq 0$, we know that $\theta_{1,m}(v) = (v^1)^{1+m}/(1+m)!$. Recall again the well-known formula for the one-point invariants:

$$\langle \tau_{1,1+3m} \rangle_{g=1+m} = \frac{1}{24^{1+m}(1+m)!}. \quad (268)$$

From these explicit expressions we immediately see the validity of identity (216), where the particular point of the Frobenius manifold is given by $v_1^* = 1/24$.

Example 2 (A_2). The superpotential reads

$$\lambda = p^3 + s_1 p + s_2, \quad v^1 = \frac{s_2}{3\sqrt{-3}^{1/4}}, \quad v^2 = \frac{\sqrt{-3}^{1/2}}{3} s_1.$$

We have $F = (v^1)^2 v^2 / 2 + (v^2)^4 / 72$ and $\eta_{\alpha\beta} = \delta_{\alpha+\beta,3}$. The first few terms of the unique calibration can be read from

$$\begin{aligned} \theta_2(v; z) &= v^1 + \left(\frac{(v^2)^3}{18} + \frac{(v^1)^2}{2} \right) z + \left(\frac{(v^1)^3}{6} + \frac{1}{18} (v^2)^3 v^1 \right) z^2 + \dots, \\ \theta_1(v; z) &= v^2 + v^1 v^2 z + \left(\frac{(v^2)^4}{36} + \frac{1}{2} (v^1)^2 v^2 \right) z^2 + \dots. \end{aligned}$$

The particular point v^* of the Frobenius manifold is given by $v_1^* = 1/12$ and $v_2^* = 0$.

Example 3 (A_4). We have

$$\begin{aligned} \lambda &= p^5 + s_1 p^3 + s_2 p^2 + s_3 p + s_4, \\ v^1 &= -\frac{s_1 s_2 - 5s_4}{25\sqrt{-5}^{1/2}}, \quad v^2 = -\frac{s_1 - 5s_3}{25}, \quad v^3 = \frac{1}{5}\sqrt{-5}^{1/2} s_2, \quad v^4 = \frac{\sqrt{-5}}{5} s_1, \end{aligned}$$

$$\begin{aligned} F &= \frac{1}{2}(v^1)^2(v^4) + v^1 v^2 v^3 + \frac{(v^2)^3}{6} + \frac{(v^4)^6}{15000} + \frac{1}{150}(v^3)^2(v^4)^3 \\ &\quad + \frac{1}{20}(v^2)^2(v^4)^2 + \frac{1}{10}v^2(v^3)^2 v^4 + \frac{(v^3)^4}{60}. \end{aligned}$$

and $\eta_{\alpha\beta} = \delta_{\alpha+\beta,5}$. The first few terms of the unique calibration and the particular point v^* of the Frobenius manifold are already given in Theorem 3.

Example 4 (D_4). The superpotential reads

$$\lambda = p^6 + s_1 p^4 + s_2 p^2 + s_3 + \frac{s_4^2}{p^2},$$

and

$$\begin{aligned} p^+(\xi; s) &= \xi - \frac{s_1}{6\xi} + \frac{s_1^2 - 4s_2}{24\xi^3} + \frac{-7s_1^3 + 36s_2 s_1 - 216s_3}{1296\xi^5} \\ &\quad + \frac{-55s_1^4 + 360s_2 s_1^2 - 864s_3 s_1 - 432s_2^2 - 5184s_4^2}{31104\xi^7} + \dots, \\ p^-(\xi; s) &= \frac{s_4}{\xi^3} + \frac{s_3 s_4}{2\xi^9} + \frac{(3s_3^2 + 4s_2 s_4^2)s_4}{8\xi^{15}} + \frac{(8s_1 s_4^4 + 20s_2 s_3 s_4^2 + 5s_3^3)s_4}{16\xi^{21}} + \dots. \end{aligned}$$

We have $\eta = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ and

$$\begin{aligned} F &= \frac{1}{2}(v^1)^2 v^3 + \frac{1}{2} v^1 (v^2)^2 + \frac{1}{36} (v^2)^3 v^3 + \frac{1}{216} (v^2)^2 (v^3)^3 + \frac{(v^3)^7}{272160} \\ &\quad + \left(\frac{v^1}{2} - \frac{1}{12} v^2 v^3 + \frac{(v^3)^3}{216} \right) (v^4)^2. \end{aligned} \tag{269}$$

We have $\theta_{\alpha,0} = v_\alpha = \eta_{\alpha\beta} v^\beta$, and

$$\begin{aligned}\theta_{1,1} &= \frac{(v^2)^2}{2} + \frac{(v^4)^2}{2} + v^1 v^3, \\ \theta_{2,1} &= \frac{1}{108} v^2 (v^3)^3 + \frac{1}{12} (v^2)^2 v^3 - \frac{1}{12} v^3 (v^4)^2 + v^1 v^2, \\ \theta_{3,1} &= \frac{(v^3)^6}{38880} + \frac{1}{72} (v^2)^2 (v^3)^2 + \frac{1}{72} (v^3)^2 (v^4)^2 + \frac{(v^2)^3}{36} - \frac{1}{12} v^2 (v^4)^2 + \frac{(v^1)^2}{2}, \\ \theta_{4,1} &= \frac{1}{108} v^4 (v^3)^3 - \frac{1}{6} v^2 v^3 v^4 + v^1 v^4.\end{aligned}$$

The particular point v^* of the Frobenius manifold is given by

$$v_1^* = \frac{1}{3}, \quad v_2^* = 0, \quad v_3^* = \frac{13}{40824}, \quad v_4^* = 0. \quad (270)$$

We list in the following table the first few of the numbers $\tau_{D_4}(g)$, again putting $\tau_{D_4}(0) = 1$.

g	0	1	2	3	4	5	6	7	8	9
$\tau_{D_4}(g)$	1	$\frac{1}{3}$	0	$\frac{13}{40824}$	$\frac{13}{122472}$	0	$\frac{1433}{16665989760}$	$\frac{253}{9999593856}$	0	$\frac{33917}{2041117097886720}$

One-point FJRW invariants of D_4 -type

APPENDIX A. WAVE FUNCTIONS

In this appendix we give a method of computing residues of pseudo-differential operators by means of wave functions. Denote by r a positive integer. Let $q_m(x)$ ($m \leq r-1$) be arbitrarily given power series of x , and L the pseudodifferential operator

$$L := \partial^r + \sum_{m=-\infty}^{r-1} q_m(x) \partial^m. \quad (271)$$

An element $\psi = \psi(x, z) \in \mathbb{C}[[x]][[z^{-1}]] \otimes e^{xz}$ of the form $\psi = \sum_{i=0}^{\infty} \phi_i(x) z^{-i} e^{xz}$ with $\phi_0(x) \equiv 1$ is called a *wave function* of L if

$$L\psi = z^r \psi. \quad (272)$$

Here we recall that $\partial^{-i}(e^{xz}) := e^{xz} z^{-i}$, $i \geq 0$, and that for all $f(x) \in \mathbb{C}[[x]]$,

$$\partial^m \circ f(x) := \sum_{\ell \geq 0} \binom{m}{\ell} f^{(\ell)}(x) \partial^{m-\ell}, \quad \forall m \in \mathbb{Z}.$$

Multiplying by e^{-xz} on both sides of (272) and comparing the coefficients of powers of z give

$$\sum_{\substack{j, \ell \geq 0 \\ j+\ell \leq i}} q_{\ell+j+r-i}(x) \binom{\ell+j+r-i}{\ell} \phi_j^{(\ell)}(x) = \phi_i(x), \quad i \geq 0. \quad (273)$$

This leads to recursive relations for the power series $\phi_i(x)$, $i \geq 0$. The solution ψ depends on a sequence of arbitrary constants C_1, C_2, \dots . Alternatively, we observe that if $\psi(x, z)$ is a wave function, then for an arbitrary power series $g(z)$ of z^{-1} with constant coefficients $g(z) = \sum_{i \geq 0} g_i z^{-i}$, $g_0 = 1$, the product $g(z) \psi(z, x)$ is again a wave function.

A pseudo-differential operator Φ of the form

$$\Phi = \sum_{i \geq 0} \tilde{\phi}_i(x) \partial^{-i}, \quad \tilde{\phi}_i(x) \in \mathbb{C}[[x]], \quad \tilde{\phi}_0(x) \equiv 1 \quad (274)$$

is called a *dressing operator of L* if

$$\Phi \circ \partial^r \circ \Phi^{-1} = L. \quad (275)$$

For the given $q_m(x)$ ($m \leq r-1$), the dressing operator is not unique, its freedom being in one-to-one correspondence with the coefficients of $g(z)$ above. Indeed, there is a one-to-one correspondence between wave functions ψ and dressing operators by $\psi = \sum_i \phi_i(x) z^{-i} e^{xz} \leftrightarrow \Phi = \sum_i \phi_i(x) \partial^{-i}$. Define the formal adjoint operator Φ^* of Φ by

$$\Phi^* := \sum_{i \geq 0} (-1)^i \partial^{-i} \circ \tilde{\phi}_i(x).$$

Fix ψ a wave function of L , and take Φ to be the corresponding dressing operator. Define ψ^* by

$$\psi^* = \psi^*(x, z) := (\Phi^{-1})^* (e^{-xz}). \quad (276)$$

Clearly, $\psi^* \in \mathbb{C}[[x]][[z^{-1}]] \otimes e^{-xz}$, the product $\psi^* e^{xz}$ has leading term 1, and

$$L^* \psi^* = z^r \psi^*. \quad (277)$$

We call ψ^* a *dual wave function of L* associated to ψ , and we call (ψ, ψ^*) a *pair of wave and dual wave functions*.

Lemma 14. *Let (ψ, ψ^*) and $(\tilde{\psi}, \tilde{\psi}^*)$ be two pairs of wave and dual wave functions of L . Then*

$$\psi(x, z) \psi^*(x, z) = \tilde{\psi}(x, z) \tilde{\psi}^*(x, z).$$

Proof. Let $g = g(x, z) := \tilde{\psi}(x, z)/\psi(x, z)$, which must have the form

$$g = \sum_{i \geq 0} g_i(x) z^{-i} \in \mathbb{C}[[x]][[z^{-1}]], \quad g_0(x) \equiv 1.$$

It follows from (273) that $g'_i(x) = 0$, $i \geq 1$. Therefore, for $i \geq 1$, g_i are all constants. Let $\Phi, \tilde{\Phi}$ be the dressing operators corresponding to $\psi, \tilde{\psi}$, respectively. We have $\tilde{\psi} = g(z) \Phi(e^{xz}) = \tilde{\Phi} \circ G(e^{xz})$, where $G := \sum_{i \geq 0} g_i \partial_x^{-i}$. Therefore, $\tilde{\Phi} = \Phi \circ G$. It follows that $(\tilde{\Phi}^{-1})^* \circ G^* = (\Phi^{-1})^*$. So $(\tilde{\Phi}^{-1})^*(g(z) e^{-xz}) = (\Phi^{-1})^*(e^{-xz})$. Namely, $g(z) \tilde{\psi}^*(x, z) = \psi^*(x, z)$. The lemma is proved. \square

Let ψ and ψ^* be a pair of wave and dual wave functions of L . Define

$$c(z) := (e^{-xz} \psi(z, x))|_{x=0}, \quad c^*(z) := (e^{xz} \psi^*(z, x))|_{x=0}, \quad H(z) := c(z) c^*(z). \quad (278)$$

It follows from Lemma 14 that the product $H(z) \in \mathbb{C}[[z^{-1}]]$ is uniquely determined by L , where we recall that L is defined by (271). We have the following lemma.

Lemma 15. *Define $z_k(x) = \text{res } L^{\frac{k}{r}}$ for all $k \geq 0$. Then*

$$H(z) = 1 + \sum_{k \geq 1} (-1)^k z_{k-1}(0) z^{-k}. \quad (279)$$

Proof. Following Liu–Vakil–Xu [86], define $\Phi_- = \sum_{i \geq 0} \phi_{-,i}(x) \partial^{-i}$ with $\phi_{-,i}(x) \in \mathbb{C}[[x]]$ and $\tilde{\phi}_{-,0}(x) \equiv 1$ as the particular dressing operator of L fixed by the additional conditions:

$$\phi_{-,i}(0) = 0, \quad i \geq 1.$$

Denote by (ψ_-, ψ_-^*) the corresponding pair of wave and dual wave functions, and define c_-, c_-^* as ψ_-, ψ_-^* evaluated at $x = 0$, respectively. It is clear from the definition that $c_-(z) \equiv 1$. Therefore, $H(z) \equiv c_-^*(z)$. Now, on one hand, noticing that

$$(\Phi_-^{-1})^* = \sum_{i \geq 0} (-\partial)^{-i} \circ \phi_{-,i}(x) = \sum_{k \geq 0} (-1)^k \sum_{\substack{i, \ell \geq 0 \\ i + \ell = k}} \binom{k-1}{\ell} \phi_{-,i}^{(\ell)}(x) \partial^{-k},$$

we find

$$\psi_-^*(x, z) = (\Phi_-^{-1})^*(e^{-xz}) = \sum_{k \geq 0} (-1)^k \sum_{\substack{i, \ell \geq 0 \\ i + \ell = k}} \binom{k-1}{\ell} \phi_{-,i}^{(\ell)}(x) z^{-k} e^{-xz}.$$

Hence

$$c_-^*(z) = \sum_{k \geq 0} (-1)^k \sum_{i=0}^k \binom{k-1}{i-1} \phi_{-,i}^{(k-i)}(0) z^{-k}. \quad (280)$$

On the other hand, from the definition of the dressing operator Φ_- we know that

$$z_k(x) = \text{res} (\Phi_- \circ \partial^r \circ \Phi_-^{-1})^{\frac{k}{r}} = \text{res} \Phi_- \circ \partial^k \circ \Phi_-^{-1}.$$

Taking $x = 0$ in this formula we find that

$$z_k(0) = (\text{res} \partial^k \circ \Phi_-^{-1})|_{x=0} = \left(\text{res} \sum_{i \geq 0} \sum_{\ell=0}^k \binom{k}{\ell} \phi_{-,i}^{(k-\ell)}(x) \partial^{\ell-i} \right) \Big|_{x=0} = \sum_{\ell=0}^k \binom{k}{\ell} \phi_{-, \ell+1}^{k-\ell}(0).$$

The lemma is proved by comparing this expression with (280) and using $c_-(z) \equiv 1$. \square

We note that, in the above proof, the function $c_-(z)$ is very simple, being just the constant function 1, but the formula (280) for $c_-^*(z)$ might be very complicated. The main point of the wave-function-pair approach is the following: using Lemma 14 it is sometimes possible to find a particular choice of pair of wave functions such that $c(z)$ and $c^*(z)$ both have relatively simple expressions so that their product can be given in closed form, and this is the case in particular both for this paper and for [52, 53], where *bispectrality* [52, 53, 58] is used for fixing the particular choice. In other situations, the Sato tau-function, theta-functions, and etc. could also be used to construct a pair of wave functions; instead of giving the details we refer to [5, 11, 14, 20, 34, 40] for specific constructions.

APPENDIX B. FROBENIUS MANIFOLDS

In this appendix, we give a brief review of the theory of Frobenius manifolds [43, 46, 47, 57, 93, 109]. Recall that a *Frobenius algebra* is a triple $(V, \mathbb{1}, \langle, \rangle)$, where V is a commutative associative algebra over \mathbb{C} with unity $\mathbb{1}$, and $\langle, \rangle : V \times V \rightarrow \mathbb{C}$ is a symmetric non-degenerate bilinear form satisfying $\langle x \cdot y, z \rangle = \langle x, y \cdot z \rangle, \forall x, y, z \in V$.

Definition 2 ([41, 43]). A *Frobenius structure* of charge d on a complex manifold M^d is a family of Frobenius algebras $(T_p M, \mathbf{1}_p, \langle, \rangle_p), p \in M$ depending holomorphically on p and satisfying:

- FM1** The metric \langle, \rangle on M is flat. Let ∇ denote the Levi-Civita connection of \langle, \rangle . Then the unit vector field $\mathbf{1}$ must satisfy $\nabla \mathbf{1} = 0$.
- FM2** Define a 3-tensor field c by $c(X, Y, Z) := \langle X \cdot Y, Z \rangle$, for X, Y, Z being holomorphic vector fields on M . The 4-tensor field ∇c must be symmetric.

FM3 There exists a holomorphic vector field E on M , called the Euler vector field, satisfying

$$\nabla \nabla E = 0, \quad (281)$$

$$[E, X \cdot Y] - [E, X] \cdot Y - X \cdot [E, Y] = X \cdot Y, \quad (282)$$

$$E \langle X, Y \rangle - \langle [E, X], Y \rangle - \langle X, [E, Y] \rangle = (2 - d) \langle X, Y \rangle. \quad (283)$$

A complex manifold endowed with a Frobenius structure is called a *Frobenius manifold*, with $\langle \cdot, \cdot \rangle$ being called the *invariant flat metric*.

Let M be a Frobenius manifold of complex dimension l . Following [43], define a one-parameter family of affine connections on M :

$$\tilde{\nabla}_X Y := \nabla_X Y + z X \cdot Y, \quad z \in \mathbb{C}. \quad (284)$$

This family of affine connections $\tilde{\nabla}$ are all flat [43], and is called the *deformed flat connection*. Moreover, it can be extended to a flat affine connection [43] on $M \times \mathbb{C}^*$, still denoted by $\tilde{\nabla}$, whose definition along the z -direction is given as follows:

$$\tilde{\nabla}_{\partial_z} X := \frac{\partial X}{\partial z} + E \cdot X - \frac{1}{z} \mu X, \quad \tilde{\nabla}_{\partial_z} \partial_z := 0, \quad \tilde{\nabla}_X \partial_z := 0, \quad (285)$$

for X being an arbitrary holomorphic vector field on $M \times \mathbb{C}^*$ with zero component along the z direction. Here, $\mu := \frac{2-d}{2} - \nabla E$. We call $\tilde{\nabla}$ the *extended deformed flat connection*. A holomorphic function $f(v; z)$ on some open subset of $M \times \mathbb{C}^*$ is called $\tilde{\nabla}$ -flat if

$$\tilde{\nabla} df = 0. \quad (286)$$

To understand the flat coordinates for the extended deformed flat connection $\tilde{\nabla}$, let us take $v = (v^1, \dots, v^l)$ a system of flat coordinates with respect to $\langle \cdot, \cdot \rangle$. Denote $\partial_\alpha = \frac{\partial}{\partial v^\alpha}$, $\eta_{\alpha\beta} := \langle \partial_\alpha, \partial_\beta \rangle$, $\eta = (\eta_{\alpha\beta})$, and $(\eta^{\alpha\beta}) := \eta^{-1}$. By FM1 we choose v^1 satisfying $\partial_1 = \mathbf{1}$. For simplicity we will assume that ∇E is diagonalizable, so the flat coordinates are chosen such that $\mu = \text{diag}(\mu_1, \dots, \mu_l)$, and we have

$$E = \sum_{1 \leq \alpha \leq l} \left(\left(1 - \frac{d}{2} - \mu_\alpha\right) v^\alpha \partial_\alpha + r^\alpha \partial_\alpha \right). \quad (287)$$

The axioms FM1–FM3 imply the local existence of a holomorphic function F , called the *Frobenius potential* of M , satisfying

$$\frac{\partial^3 F}{\partial v^\alpha \partial v^\beta \partial v^\gamma} = c_{\alpha\beta\gamma} := c(\partial_\alpha, \partial_\beta, \partial_\gamma), \quad (288)$$

$$EF = (3 - d)F + \text{a quadratic function of } v. \quad (289)$$

Clearly, F is uniquely determined by the Frobenius structure up to a quadratic function of v . The flatness of $\tilde{\nabla}$ implies that there locally exist l independent $\tilde{\nabla}$ -flat holomorphic functions

$$\tilde{v}_1, \dots, \tilde{v}_l$$

on $M \times \mathbb{C}^*$, called the *deformed flat coordinates* for the Frobenius manifold. Let us give their construction. For a $\tilde{\nabla}$ -flat holomorphic function f on $M \times \mathbb{C}^*$, denote $y^\alpha = \eta^{\alpha\beta} \partial f / \partial v^\beta$ and $y = (y^1, \dots, y^l)^T$. By definition the column-vector-valued function y satisfies the following system of linear differential equations:

$$\frac{\partial y}{\partial v^\alpha} = z C_\alpha y, \quad (290)$$

$$\frac{dy}{dz} = \left(\mathcal{U} + \frac{\mu}{z} \right) y. \quad (291)$$

Here, $(C_\alpha)_\gamma^\beta := c_{\alpha\gamma}^\beta$ and $\mathcal{U}_\beta^\alpha := E^\rho c_{\rho\beta}^\alpha$. To fix a system of deformed flat coordinates we need to choose a basis of the solution space to (290)–(291). Observe that the ODE system (291) possesses a Fuchsian singular point at $z = 0$ and an irregular singular point of Poincaré rank 1 at $z = \infty$. The axioms of Frobenius manifolds imply the isomonodromicity of (290)–(291). The monodromy data around $z = 0$ is then given by two constant matrices μ, R , satisfying

$$R^* = -e^{\pi i \mu} R e^{-\pi i \mu} \quad (292)$$

$$z^\mu R z^{-\mu} = R_0 + R_1 z + R_2 z^2 + \dots \quad (293)$$

for some matrices R_k , $k \geq 0$ satisfying

$$[\mu, R_k] = k R_k, \quad k \geq 0. \quad (294)$$

Let us fix a choice of R . It is shown in [43, 46] (cf. also [70]) that locally there exists a fundamental solution matrix $\mathcal{Y} = \mathcal{Y}(v; z)$ around $z = 0$ to (290)–(291) of the form

$$\mathcal{Y}(v; z) = \Theta(v; z) z^\mu z^R = \sum_{m \geq 0} \Theta_m(v) z^{m+\mu} z^R, \quad (295)$$

where $\Theta(v; z)$ is a matrix-valued analytic function on $M \times \mathbb{C}$ satisfying

$$\Theta(v; 0) \equiv I, \quad (296)$$

$$\eta^{-1} \Theta(v; -z)^T \eta \Theta(v; z) \equiv I. \quad (297)$$

It can be easily checked that the matrix-valued function $\Theta(v; z)$ satisfies

$$\partial^\beta \Theta_\gamma^\alpha(v; z) = \partial^\alpha \Theta_\gamma^\beta(v; z).$$

Then by the Poincaré lemma, there locally exist holomorphic functions $\theta_\gamma(v; z)$ of the form

$$\theta_\gamma(v; z) = \sum_{m \geq 0} \theta_{\gamma, m}(v) z^m, \quad (298)$$

such that

$$\eta^{\alpha\beta} \frac{\partial \theta_\gamma(v; z)}{\partial v^\beta} = \Theta_\gamma^\alpha(v; z). \quad (299)$$

Hence the functions $\tilde{v}_1(v; z), \dots, \tilde{v}_n(v; z)$ defined by

$$(\tilde{v}_1(v; z), \dots, \tilde{v}_n(v; z)) = (\theta_1(v; z), \dots, \theta_l(v; z)) z^\mu z^R \quad (300)$$

give a system of deformed flat coordinates on $M \times \mathbb{C}^*$ (see [43, 46, 47, 57] for more details).

From the above construction we know that $\theta_{\alpha, m}$, $m \geq 0$ satisfy the following conditions:

$$\partial_\alpha \partial_\beta (\theta_{\gamma, m+1}) = c_{\alpha\beta}^\sigma \partial_\sigma (\theta_{\gamma, m}), \quad p \geq 0, \quad (301)$$

$$E (\partial_\beta \theta_{\alpha, m}) = (p + \mu_\alpha + \mu_\beta) \partial_\beta (\theta_{\alpha, m}) + \sum_{1 \leq k \leq m} (R_k)_\alpha^\gamma \partial_\beta (\theta_{\gamma, m-k}), \quad p \geq 0, \quad (302)$$

$$\frac{\partial \theta_{\alpha, 0}}{\partial v^\beta} = \eta_{\alpha\beta}, \quad (303)$$

$$\langle \nabla \theta_\alpha(v; z), \nabla \theta_\beta(v; -z) \rangle = \eta_{\alpha\beta}. \quad (304)$$

One can further normalize $\theta_{\alpha, m}(v)$ by requiring

$$\theta_{\alpha, 0} = v_\alpha, \quad (305)$$

$$\partial_1 (\theta_{\alpha, m+1}) = \theta_{\alpha, m}, \quad m \geq 0. \quad (306)$$

Indeed, (305) is obviously compatible with (301)–(304), and we leave the verification of the compatibility between (306) and (301)–(304) as an exercise to the reader. As in [50], we call a choice of $\{\theta_{\alpha,m}\}_{m \geq 0}$ satisfying (301)–(306) a *calibration* on M .

According to Kontsevich–Manin [83, 93], the genus zero part of a CohFT

$$(V^l, \eta, \mathbb{1}, \{\Omega_{g,n}\}_{2g-2+n>0})$$

gives a pre-Frobenius structure on the convergence domain B under the assumption (see Section 2). Here “pre” means the axiom FM3 is not required. (In some literature a pre-Frobenius structure is called a Frobenius structure.) Let us recall Kontsevich–Manin’s construction of the Frobenius structure from the CohFT. As in Section 2, take $e_1 = \mathbb{1}, e_2, \dots, e_l$ a basis of V . Define a metric $\langle \cdot, \cdot \rangle$ on B and a multiplication “ \cdot ” on the tangent spaces of B by

$$\langle \partial_\alpha, \partial_\beta \rangle := \eta_{\alpha\beta}, \quad (307)$$

$$\langle \partial_\alpha \cdot \partial_\beta, \partial_\gamma \rangle := \frac{\partial^3 F}{\partial v^\alpha \partial v^\beta \partial v^\gamma}, \quad (308)$$

where F is defined in (31). From (307), it is obvious that the metric $\langle \cdot, \cdot \rangle$ is flat and in fact (v^1, \dots, v^n) is a system of flat coordinates for $\langle \cdot, \cdot \rangle$. The invariance of $\langle \cdot, \cdot \rangle$ with respect to the multiplication “ \cdot ” on each tangent space as well as the FM2 are also obviously true from the construction. The axioms of the CohFT imply [83] that the multiplication “ \cdot ” is associative and ∂_1 is a unit vector field. So $(B, \langle \cdot, \cdot \rangle, \cdot, \partial_1)$ is a pre-Frobenius manifold. Using (35) we see further that a *homogeneous* CohFT $(V, \eta, \mathbb{1}, \{\Omega_{g,n}\}_{2g-2+n>0})$ of charge d endows B with a Frobenius structure of charge d with E given by (32) being the Euler vector field.

REFERENCES

- [1] Adler, M., van Moerbeke, P., A matrix integral solution to two-dimensional W_p -gravity. *Comm. Math. Phys.* **147** (1992), 25–56.
- [2] Alexandrov, A., Milanov, T., Matrix model for the total descendant potential of a simple singularity of type D . arXiv: 2011.00837.
- [3] Arnold, V.I., Normal forms of functions near degenerate critical points, Weyl groups A_k, D_k, E_k and Lagrange singularities. *Funct. Anal. Appl.* **6** (1972), 254–272.
- [4] Arnold, V.I., Varchenko, A.N., Gusein-Zade, S.M., *Singularities of Differentiable Maps: Volume II Monodromy and Asymptotic Integrals*. Springer Science Business Media, 2012.
- [5] Babelon, O., Bernard, D., Talon, M., *Introduction to classical integrable systems*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 2003.
- [6] Bakalov, B., Milanov, T., \mathcal{W} -constraints for the total descendant potential of a simple singularity. *Compos. Math.* **149** (2013), 840–888.
- [7] Balog, J., Fehér, L., O’Raifeartaigh, L., Forgacs, P., Wipf, A., Toda theory and W-algebra from a gauged WZNW point of view. *Annals of Physics* **203** (1990), 76–136.
- [8] Balogh, F., Yang, D., Geometric interpretation of Zhou’s explicit formula for the Witten-Kontsevich tau function. *Lett. Math. Phys.* **107** (2017), 1837–1857.
- [9] Balogh, F., Yang, D., Zhou, J., Explicit formula for Witten’s r -spin partition function. In preparation.
- [10] Basalaev, A., Buryak, A., Open Saito Theory for A and D Singularities. arXiv:1909.00598.
- [11] Bertola, M., Dubrovin, B., Yang, D., Correlation functions of the KdV hierarchy and applications to intersection numbers over $\overline{\mathcal{M}}_{g,n}$. *Physica D: Nonlinear Phenomena* **327** (2016), 30–57.
- [12] Bertola, M., Dubrovin, B., Yang, D., Simple Lie algebras and topological ODEs. *IMRN* (2018), 1368–1410.
- [13] Bertola, M., Dubrovin, B., Yang, D., Simple Lie algebras, Drinfeld–Sokolov hierarchies, and multi-point correlation functions. *Mosc. Math. J.* (to appear). arXiv:1610.07534.
- [14] Bertola, M., Yang, D., The partition function of the extended r -reduced Kadomtsev–Petviashvili hierarchy. *Journal of Physics A: Mathematical and Theoretical* **48** (2015), 195205.
- [15] Bouchard, V., Eynard, B., Reconstructing WKB from topological recursion. *J. Éc. polytech. Math.* **4** (2017), 845–908.

- [16] Brézin, E., Hikami, S., Intersection numbers of Riemann surfaces from Gaussian matrix models. *J. High Energy Phys.* (2007), 096.
- [17] Brézin, E., Hikami, S., Computing topological invariants with one and two-matrix models. *J. High Energy Phys.* (2009), 110.
- [18] Brézin, E., Hikami, S., Random matrix theory with an external source. *SpringerBriefs in Mathematical Physics* **19**. Springer, Singapore, 2016.
- [19] Brini, A., E_8 spectral curves. *Proc. Lond. Math. Soc.* **121** (2020), 954–1032.
- [20] Buryak, A., Open intersection numbers and the wave function of the KdV hierarchy. *Mosc. Math. J.* **16** (2016), 27–44.
- [21] Buryak, A., Double ramification cycles and the n -point function for the moduli space of curves. *Mosc. Math. J.* **17** (2017), 1–13.
- [22] Buryak, A., Extended r -spin theory and the mirror symmetry for the A_{r-1} -singularity. *Mosc. Math. J.* **20** (2020), 475–493.
- [23] Buryak, A., Clader, E., Tessler, R., Closed extended r -spin theory and the Gelfand-Dickey wave function. *J. Geom. Phys.* **137** (2019), 132–153.
- [24] Buryak, A., Posthuma, H., Shadrin, S., A polynomial bracket for the Dubrovin-Zhang hierarchies. *J. Differential Geom.* **92** (2012), 153–185.
- [25] Cafasso, M., Wu, C.-Z., Borodin-Okounkov formula, string equation and topological solutions of Drinfeld-Sokolov hierarchies. *Lett. Math. Phys.* **109** (2019), 2681–2722.
- [26] Chang, H.-L., Li, J., Li, W.-P., Wittens top Chern class via cosection localization. *Inventiones Mathematicae* **200** (2015), 1015–1063.
- [27] Chiodo, A., The Witten top Chern class via K-theory. *Journal of Algebraic Geometry*, **15** (2006), 681–707.
- [28] Chiodo, A., Stable twisted curves and their r -spin structures. *Ann. Inst. Fourier (Grenoble)* **58** (2008), 1635–1689.
- [29] Cohen, P.B., Manin, Y., Zagier, D., Automorphic pseudodifferential operators. *Algebraic aspects of integrable systems*, 17–47, *Progr. Nonlinear Differential Equations Appl.*, **26**, Birkhäuser Boston, Boston, MA, 1997.
- [30] Date, E., Jimbo, M., Kashiwara, M., Miwa, T., Transformation groups for soliton equations. In *Publications of RIMS* **18** (1981), 1077–1110.
- [31] De Sole, A., Kac, V.G., Valeri, D., Classical W -algebras and generalized Drinfeld-Sokolov bi-Hamiltonian systems within the theory of Poisson vertex algebras. *Comm. Math. Phys.* **323** (2013), 663–711.
- [32] Deligne, P., Mumford, D., The irreducibility of the space of curves of given genus. *Inst. Hautes Études Sci. Publ. Math. No.* **36** (1969), 75–109.
- [33] Deift, P., *Orthogonal Polynomials and Random Matrices: A Riemann–Hilbert Approach*. Courant Lecture Notes in Mathematics **3**. American Mathematical Society, Providence, 1999.
- [34] Dickey, L.A., *Soliton Equations and Hamiltonian Systems* (Vol. **26**). 2nd edition. World Scientific, 2003.
- [35] Dijkgraaf, R., Verlinde, H., Verlinde, E., Notes on topological string theory and 2D quantum gravity. *String theory and quantum gravity (Trieste, 1990)*, 91–156, World Sci. Publ., River Edge, NJ, 1991.
- [36] Dijkgraaf, R., Verlinde, H., Verlinde, E., Topological strings in $d < 1$. *Nuclear Physics B* **352** (1991), 59–86.
- [37] Dijkgraaf, R., Witten, E., Mean field theory, topological field theory, and multi-matrix models. *Nuclear Phys. B* **342** (1990), 486–522.
- [38] Drinfeld, V.G., Sokolov, V.V., Lie algebras and equations of Korteweg–de Vries type, *J. Math. Sci.* **30** (1985), 1975–2036. Translated from *Itogi Nauki i Tekhniki, Seriya Sovremennye Problemy Matematiki (Noveishie Dostizheniya)* **24** (1984), 81–180.
- [39] Dubrovin, B., Periodic problems for the Korteweg–de Vries equation in the class of finite band potentials. *Funct. Anal. Appl.* **9** (1975), 215–223.
- [40] Dubrovin, B.A., Theta-functions and nonlinear equations. (Russian) With an appendix by I.M. Krichever. *Uspekhi Mat. Nauk* **36** (1981), no. 2 (218), 11–80.
- [41] Dubrovin, B., Integrable systems in topological field theory. *Nuclear Phys. B* **379** (1992), 627–689.
- [42] Dubrovin, B., Integrable systems and classification of 2-dimensional topological field theories. *Integrable systems (Luminy, 1991)*, *Progr. Math.*, **115**, 313–359, Birkhäuser Boston, 1993.
- [43] Dubrovin, B., Geometry of 2D topological field theories. In “Integrable Systems and Quantum Groups” (Montecatini Terme, 1993), Editors: Francaviglia, M., Greco, S.. *Springer Lecture Notes in Math.* **1620**, 120–348, 1996.
- [44] Dubrovin, B., Flat pencils of metrics and Frobenius manifolds. *Integrable systems and algebraic geometry (Kobe/Kyoto, 1997)*, 47–72, World Sci. Publ., River Edge, NJ, 1998.
- [45] Dubrovin, B., Geometry of the space of orbits of a Coxeter Group. *Surv. Diff. Geom.* **4** (1998), 181–211.

- [46] Dubrovin, B., Painlevé transcendents in two-dimensional topological field theory. In “The Painlevé Property, One Century later”, Editor: R. Conte, Springer, 287–412, 1999.
- [47] Dubrovin, B., On almost duality for Frobenius manifolds. In “Geometry, Topology, and Mathematical Physics” (Amer. Math. Soc. Transl. Ser. 2, Vol. 212, 75–132), Editors: Buchstaber, V.M., Krichever, I.M., AMS, Prov., R.I., 2004.
- [48] Dubrovin, B., Gromov-Witten invariants and integrable hierarchies of topological type. *Topology, geometry, integrable systems, and mathematical physics*, 141–171, Amer. Math. Soc. Transl. Ser. **2**, 234, Adv. Math. Sci., **67**, Amer. Math. Soc., Providence, RI, 2014.
- [49] Dubrovin, B., Liu, S.-Q., Zhang, Y., Frobenius manifolds and central invariants for the Drinfeld–Sokolov biHamiltonian structures. *Advances in Mathematics* **219** (2008), 780–837.
- [50] Dubrovin, B., Liu, S.-Q., Yang, D., Zhang, Y., Hodge integrals and tau-symmetric integrable hierarchies of Hamiltonian evolutionary PDEs. *Adv. Math.* **293** (2016), 382–435.
- [51] Dubrovin, B., Strachan, I.A.B., Zhang, Y., Zuo, D., Extended affine Weyl groups of BCD-type: their Frobenius manifolds and Landau-Ginzburg superpotentials. *Adv. Math.* **351** (2019), 897–946.
- [52] Dubrovin, B., Yang, D., Zagier, D., Gromov-Witten invariants of the Riemann sphere. *Pure Appl. Math. Q.* **16** (2020), 153–190.
- [53] Dubrovin, B., Yang, D., Zagier, D., On tau-functions for the KdV hierarchy. *Selecta Math.* (to appear). arXiv:1812.08488.
- [54] Dubrovin, B., Yang, D., Zagier, D., Geometry and arithmetic of integrable hierarchies of KdV type. II. Topological ODEs of the second kind. In preparation.
- [55] Dubrovin, B., Zhang, Y., Bi-Hamiltonian hierarchies in 2D topological field theory at one-loop approximation. *Comm. Math. Phys.* **198** (1998), 311–361.
- [56] Dubrovin, B., Zhang, Y., Extended affine Weyl groups and Frobenius manifolds. *Compositio Math.* **111** (1998), 167–219.
- [57] Dubrovin, B., Zhang, Y., Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov-Witten invariants. arXiv:math/0108160.
- [58] Duistermaat, J.J., Grünbaum, F.A., Differential equations in the spectral parameter. *Comm. Math. Phys.* **103** (1986), 177–240.
- [59] Dunin-Barkowski, P., Norbury, P., Orantin, N., Popolitov, A., Shadrin, S., Dubrovin’s superpotential as a global spectral curve. *J. Inst. Math. Jussieu* **18** (2019), 449–497.
- [60] Eguchi, T., Kanno, H., Yamada, Y., Yang, S.-K., Topological strings, flat coordinates and gravitational descendants. *Phys. Lett. B* **305** (1993), 235–241.
- [61] Eguchi, T., Yamada, Y., Yang, S.-K., Topological field theories and the period integrals. *Modern Phys. Lett. A* **8** (1993), 1627–1637.
- [62] Eguchi, T., Yamada, Y., Yang, S.-K., On the genus expansion in the topological string theory. *Rev. Math. Phys.* **7** (1995), 279–309.
- [63] Eguchi, T., Yang, S.-K., A new description of the E_6 singularity. *Phys. Lett. B* **394** (1997), 315–322.
- [64] Faber, C., Shadrin, S., Zvonkine, D., Tautological relations and the r -spin Witten conjecture. *Annales scientifiques de l’École Normale Supérieure* **43** (2010), 621–658.
- [65] Fan, H., Francis, A., Jarvis, T., Merrell, E., Ruan, Y., Witten’s D_4 Integrable Hierarchies Conjecture. *Chinese Annals of Mathematics* **37** (2016), 175–192.
- [66] Fan, H., Jarvis, T., Ruan, Y., The Witten equation and its virtual fundamental cycle. arXiv:0712.4025.
- [67] Fan, H., Jarvis, T., Ruan, Y., The Witten equation, mirror symmetry, and quantum singularity theory. *Annals of Mathematics* **178** (2013), 1–106.
- [68] Frenkel, E., Givental, A., Milanov, T., Soliton equations, vertex operators, and simple singularities. *Funct. Anal. Other Math.* **3** (2010), 47–63.
- [69] Givental, A., Semisimple Frobenius structures at higher genus. *IMRN* **2001**, 1265–1286.
- [70] Givental, A.B., Gromov–Witten invariants and quantization of quadratic Hamiltonians. *Mosc. Math. J.* **1**, 551–568.
- [71] Givental, A., A_{n-1} singularities and nKdV hierarchies. Dedicated to Vladimir I. Arnold on the occasion of his 65th birthday. *Mosc. Math. J.* **3** (2003), 475–505.
- [72] Givental, A.B., Milanov, T.E., Simple singularities and integrable hierarchies. The breadth of symplectic and Poisson geometry, 173–201, *Progr. Math.*, **232**, Birkhäuser Boston, Boston, MA, 2005.
- [73] Harer, J., Zagier, D., The Euler characteristic of the moduli space of curves. *Invent. Math.* **85** (1986), 457–485.

- [74] Hertling, C., *Frobenius manifolds and moduli spaces for singularities*. Cambridge Tracts in Mathematics **151**, Cambridge University Press, Cambridge, 2002.
- [75] Jarvis, T.J., Kimura, T., Vaintrob, A., Moduli spaces of higher spin curves and integrable hierarchies. *Compositio Mathematica* **126** (2001), 157–212.
- [76] Kac, V., Infinite-dimensional algebras, Dedekinds η -function, classical Möbius function and the very strange formula. *Advances in Mathematics* **30** (1978), 85–136.
- [77] Kac, V., Schwarz, A., Geometric interpretation of the partition function of 2D gravity. *Phys. Lett. B* **257** (1991), 329–334.
- [78] Kac, V., Wakimoto, M., Exceptional hierarchies of Soliton Equations. *Proc. Symp. Pure Math.* **49** (1989), 138–177.
- [79] Khesin, B., Zakharevich, I., Poisson-Lie group of pseudodifferential symbols. *Comm. Math. Phys.* **171** (1995), 475–530.
- [80] Klemm, A., Theisen, S., Schmidt, M.G., Correlation functions for topological Landau-Ginzburg models with $c \leq 3$, *Internat. J. Modern Phys. A* **7** (1992), 6215–6244.
- [81] Knapp, J., Omer, H., Matrix factorizations, minimal models and Massey products, *J. High Energy Phys.* (2006), 064, 33pp.
- [82] Kontsevich, M., Intersection theory on the moduli space of curves and the matrix Airy function. *Comm. Math. Phys.* **147** (1992), 1–23.
- [83] Kontsevich, M., Manin, Yu., Gromov-Witten classes, quantum cohomology, and enumerative geometry. *Comm. Math. Phys.* **164** (1994), 525–562.
- [84] Kostant, B., The Principal Three-Dimensional Subgroup and the Betti Numbers of a Complex Simple Lie Group. *American Journal of Mathematics* **81** (1959), 973–1032.
- [85] Lerche, W., Warner, N.P., Exceptional SW geometry from ALE fibrations. *Phys. Lett. B* **423** (1998), 79–86.
- [86] Liu, K., Vakil, R., Xu, H., Formal pseudodifferential operators and Witten’s r -spin numbers. *J. Reine Angew. Math.* **728** (2017), 1–33.
- [87] Liu, K., Xu, H., New results of intersection numbers on moduli spaces of curves. *Proc. Natl. Acad. Sci. USA* **104** (2007), 13896–13900.
- [88] Liu, K., Xu, H., The n -point functions for intersection numbers on moduli spaces of curves. *Adv. Theor. Math. Phys.* **15** (2011), 1201–1236.
- [89] Liu, S.-Q., Ruan, Y., Zhang, Y., BCFG Drinfeld–Sokolov hierarchies and FJRW-theory. *Inventiones mathematicae* **201** (2015), 711–772.
- [90] Liu, S.-Q., Wu, C.-Z., Zhang, Y., On the Drinfeld-Sokolov hierarchies of D type. *IMRN* (2011), 1952–1996.
- [91] Liu, S.-Q., Yang, D., Zhang, Y., Uniqueness Theorem of \mathcal{W} -Constraints for Simple Singularities. *Lett. Math. Phys.* **103** (2013), 1329–1345.
- [92] Looijenga, E., A period mapping for certain semi-universal deformations. *Compositio Math.* **30** (1975), 299–316.
- [93] Manin, Yu.I., *Frobenius manifolds, quantum cohomology, and moduli spaces*. AMS Colloquium Publ., **47**, Providence, RI, 1999.
- [94] Mehta, M.L., *Random Matrices*, 2nd edition. Academic Press, Cambridge, 1991.
- [95] Mochizuki, T., The virtual class of the moduli stack of stable r -spin curves. *Comm. Math. Phys.* **264** (2006), 1–40.
- [96] Okounkov, A., Generating functions for intersection numbers on moduli spaces of curves. *IMRN* (2002), 933–957.
- [97] Pandharipande, R., Pixton, A., Zvonkine, D., Relations on $\overline{\mathcal{M}}_{g,n}$ via 3-spin structures. *J. Amer. Math. Soc.* **28** (2015), 279–309.
- [98] Pandharipande, R., Pixton, A., Zvonkine, D., Tautological relations via r -spin structures. With an appendix by F. Janda and the authors. *J. Algebraic Geom.* **28** (2019), 439–496.
- [99] Penner, R.C., Perturbative series and the moduli space of Riemann surfaces. *J. Differential Geom.* **27** (1988), 35–53.
- [100] Polishchuk, A., Witten’s top Chern class on the moduli space of higher spin curves. In *Frobenius manifolds* (pp. 253–264). Vieweg + Teubner Verlag, 2004.
- [101] Polishchuk, A., Vaintrob, A., Algebraic construction of Witten’s top Chern class. In *Advances in Algebraic Geometry motivated by Physics*, E. Previato, ed., 229–250. AMS, 2001.
- [102] Saito, K., Primitive forms for a universal unfolding of a function with an isolated critical point. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **28** (1981), 775–792 (1982).
- [103] Saito, K., Period mapping associated to a primitive form. *Publ. Res. Inst. Math. Sci.* **19** (1983), 1231–1264.

- [104] Saito, K., On a linear structure of the quotient variety by a finite reflection group. *Publ. Res. Inst. Math. Sci.* **29** (1993), 535–579.
- [105] Saito, M., On the structure of Brieskorn lattices. *Ann. Inst. Fourier (Grenoble)* **39** (1989), 27–72.
- [106] Sato, M., Soliton Equations as Dynamical Systems on a Infinite Dimensional Grassmann Manifolds (Random Systems and Dynamical Systems). *RIMS Kokyuroku* **439** (1981), 30–46.
- [107] Segal, G., Wilson, G., Loop groups and equations of KdV type. *Inst. Hautes Études Sci. Publ. Math. No.* **61** (1985), 5–65.
- [108] Takasaki, K., Integrable hierarchy underlying topological Landau-Ginzburg models of D -type. *Lett. Math. Phys.* **29** (1993), 111–121.
- [109] Teleman, C., The structure of 2D semi-simple field theories. *Inventiones mathematicae* **188** (2012), 525–588.
- [110] Witten, E., Two-dimensional gravity and intersection theory on moduli space. *Surveys in Differential Geometry* (Cambridge, MA, 1990), 243–310, Lehigh Univ., Bethlehem, PA, 1991.
- [111] Witten, E., The N matrix model and gauged WZW models. *Nuclear Phys. B* **371** (1992), 191–245.
- [112] Witten, E., Algebraic geometry associated with matrix models of two-dimensional gravity (pp. 235–269). *Topological methods in modern mathematics* (Stony Brook, NY, 1991), Publish or Perish, Houston, TX, 1993.
- [113] Wu, C.-Z., A remark on Kac-Wakimoto hierarchies of D -type. *J. Phys. A* **43** (2010), 035201.
- [114] Yang, D., Zagier, D., Arithmetic properties of coefficients of Picard–Fuchs differential equations, In preparation.
- [115] Zagier, D., The Arithmetic and Topology of Differential Equations. In “Proceedings of the European Congress of Mathematics” (Berlin, 18–22 July, 2016), Mehrmann, V.; Skutella, M. (Eds.), European Mathematical Society, 717–776, 2018.
- [116] Zhou, J., Topological recursions of Eynard-Orantin type for intersection numbers on moduli spaces of curves. *Lett. Math. Phys.* **103** (2013), 1191–1206.
- [117] Zhou, J., Solution of W-constraints for R-spin intersection numbers. arXiv:1305.6991.
- [118] Zhou, J., Quantum deformation theory of the Airy curve and mirror symmetry of a point. arXiv:1405.5296.
- [119] Zhou, J., Emergent geometry and mirror symmetry of a point. arXiv:1507.01679.
- [120] Zhou, J., Fermionic computations for integrable hierarchies. arXiv:1508.01999.
- [121] Zhou, J., On emergent geometry of the Gromov-Witten theory of quintic Calabi-Yau threefold. arXiv:2008.03407.
- [122] Zuber, J.-B., On Dubrovin topological field theories. *Modern Physics Letters A* **9** (1994), 749–760.

Di Yang

School of Mathematical Sciences, USTC, Jinzhai Road 96, Hefei 230026, P.R. China
 diyang@ustc.edu.cn

Don Zagier

Max-Planck-Institut für Mathematik, Vivatsgasse 7, Bonn 53111, Germany
 and International Centre for Theoretical Physics, Strada Costiera 11, Trieste 34014, Italy
 dbz@mpim-bonn.mpg.de