

# Quantum Black Holes, Wall Crossing, and Mock Modular Forms

---

Atish Dabholkar<sup>1,2</sup>, Sameer Murthy<sup>3</sup>, and Don Zagier<sup>4,5</sup>

<sup>1</sup>*Theory Division, CERN, PH-TH Case C01600, CH-1211, 23 Geneva, Switzerland*

<sup>2</sup>*Laboratoire de Physique Théorique et Hautes Energies (LPTHE)*

*Université Pierre et Marie Curie-Paris 6; CNRS UMR 7589*

*Tour 24-25, 5<sup>ème</sup> étage, Boite 126, 4 Place Jussieu, 75252 Paris Cedex 05, France*

<sup>3</sup>*NIKHEF theory group, Science Park 105, 1098 XG Amsterdam, The Netherlands*

<sup>4</sup>*Max-Planck-Institut für Mathematik, Vivatsgasse 7, 53111 Bonn, Germany*

<sup>5</sup>*Collège de France, 3 Rue d'Ulm, 75005 Paris, France*

*Emails:* atish@lpthe.jussieu.fr, murthy.sameer@gmail.com, don.zagier@mpim-bonn.mpg.de

**ABSTRACT:** We show that the meromorphic Jacobi form that counts the quarter-BPS states in  $\mathcal{N} = 4$  string theories can be canonically decomposed as a sum of a *mock Jacobi form* and an *Appell-Lerch sum*. The quantum degeneracies of single-centered black holes are Fourier coefficients of this mock Jacobi form, while the Appell-Lerch sum captures the degeneracies of multi-centered black holes which decay upon wall-crossing. The *completion* of the mock Jacobi form restores the modular symmetries expected from  $AdS_3/CFT_2$  holography but has a holomorphic anomaly reflecting the non-compactness of the microscopic CFT. For every positive integral value  $m$  of the magnetic charge invariant of the black hole, our analysis leads to a special mock Jacobi form of weight two and index  $m$ , which we characterize uniquely up to a Jacobi cusp form. This family of special forms and another closely related family of weight-one forms contain almost all the known mock modular forms including the mock theta functions of Ramanujan, the generating function of Hurwitz-Kronecker class numbers, the mock modular forms appearing in the Mathieu and Umbral moonshine, as well as an infinite number of new examples.

**KEYWORDS:** [black holes](#), [mock modular forms](#), [superstrings](#), [wall-crossing](#).

---

## Contents

<b>1. Introduction</b>	<b>3</b>
1.1 Introduction for mathematicians	3
1.2 Introduction for physicists	6
1.3 Organization of the paper	12
<b>2. Review of Type-II superstring theory on <math>K3 \times T^2</math></b>	<b>12</b>
<b>3. Modular forms in one variable</b>	<b>17</b>
3.1 Basic definitions and properties	18
3.2 Quantum black holes and modular forms	21
<b>4. Jacobi forms</b>	<b>23</b>
4.1 Basic definitions	23
4.2 Theta expansion and Taylor expansion	24
4.3 Example: Jacobi forms of index 1	27
4.4 Hecke-like operators	29
4.5 Quantum black holes and Jacobi forms	32
<b>5. Siegel modular forms</b>	<b>33</b>
5.1 Definitions and examples of Siegel modular forms	33
5.2 Quantum black holes and Siegel modular forms	36
<b>6. Walls and contours</b>	<b>37</b>
<b>7. Mock modular forms</b>	<b>40</b>
7.1 Pure mock modular forms	40
7.2 Mock Jacobi forms	44
7.3 Mock modular forms: the general case	48
7.4 Superconformal characters and mock modular forms	50
<b>8. From meromorphic Jacobi forms to mock modular forms</b>	<b>51</b>
8.1 The Fourier coefficients of a meromorphic Jacobi form	52
8.2 The polar part of $\varphi$ (case of simple poles)	53
8.3 Mock modularity of the Fourier coefficients	57
8.4 The case of double poles	61
8.5 Examples	64

<b>9. Special mock Jacobi forms</b>	<b>67</b>
9.1 The simplest meromorphic Jacobi forms	67
9.2 Choosing the function $\varphi$ for weight 2 and small index	69
9.3 Further data and formulation of main results for the weight 2 family	75
9.4 Optimal choice of the function $\Phi_{2,m}$	81
9.5 Observations on the weight one family, integrality, and positivity	86
9.6 Higher weights	95
<b>10. Structure theorems for Jacobi and mock Jacobi forms</b>	<b>97</b>
10.1 Relations among the polar coefficients of weak Jacobi forms	98
10.2 Choosing optimal versions of weak elliptic forms	102
10.3 The residues of mock Jacobi forms	107
10.4 Remaining proofs of theorems from §9	109
<b>11. Quantum black holes and mock modular forms</b>	<b>114</b>
11.1 Mock Jacobi forms, immortal black holes, and $AdS_2$ holography	115
11.2 Meromorphy, mock modularity, and noncompactness	120
11.3 The M-Theory limit and $AdS_3$ holography	122
11.4 Open problems and comments	126
<b>A. Appendix: Tables of special mock Jacobi forms</b>	<b>130</b>
A.1 Table of $\mathcal{Q}_M$ (weight 2 case)	130
A.2 Table of $\Phi_{1,m}^0$ of optimal growth	134
A.3 Table of $\mathcal{Q}_M$ (weight 1 case)	138
<b>B. Bibliography</b>	<b>143</b>

*“My dream is that I will live to see the day when our young physicists, struggling to bring the predictions of superstring theory into correspondence with the facts of nature, will be led to enlarge their analytic machinery to include not only theta-functions but mock theta-functions . . . But before this can happen, the purely mathematical exploration of the mock-modular forms and their mock-symmetries must be carried a great deal further.”*

*Freeman Dyson (1987 Ramanujan Centenary Conference)*

# 1. Introduction

The quantum degeneracies associated with a black hole horizon are of central importance in quantum gravity. The counting function for these degeneracies for a large class of black holes in string theory is expected to be modular from the perspective of holography. However, in situations with wall-crossing, there is an apparent loss of modularity and it is far from clear if and how such a counting function can be modular. In the context of quarter-BPS black holes in  $\mathcal{N} = 4$  supersymmetric theories, we develop the required analytic machinery that provides a complete answer to this question which leads naturally to the mathematics of mock modular forms. We present a number of new mathematical results motivated by but independent of these physical considerations.

Since this paper is of possible interest to both theoretical physicists (especially string theorists) and theoretical mathematicians (especially number theorists), we give two introductions in their respective dialects.

## 1.1 Introduction for mathematicians

In the quantum theory of black holes in the context of string theory, the physical problem of counting the dimensions of certain eigenspaces (“the number of quarter-BPS dyonic states of a given charge”) has led to the study of Fourier coefficients of certain *meromorphic* Siegel modular forms and to the question of the modular nature of the corresponding generating functions. Using and refining results of S. Zwegers [119], we show that these generating functions belong to the recently discovered class of functions called *mock modular forms*.

Since this notion is still not widely known, it will be reviewed in some detail (in §7.1). Very roughly, a mock modular form of weight  $k$  (more precisely, “pure” mock modular forms; we will also introduce a somewhat more general notion of “mixed” mock modular forms in §7.3) is a holomorphic function  $f$  in the upper half plane to which is associated a holomorphic modular form  $g$  of weight  $2 - k$ , called the “shadow” of  $f$ , such that the sum of  $f$  and a suitable non-holomorphic integral of  $g$  transforms like a holomorphic modular form of weight  $k$ . Functions of this type occur in several contexts in mathematics and mathematical physics: as certain  $q$ -hypergeometric series (like Ramanujan’s original mock theta functions), as generating functions of class numbers of imaginary quadratic fields [115], or as characters of extended superconformal algebras [88], with special cases of the last class being conjecturally related to the Mathieu group  $M_{24}$  [47]. They also arise, as was shown by Zwegers in his thesis [119], as the Fourier coefficients of meromorphic Jacobi forms. It is this last occurrence which is at the origin of the connection to black hole physics, because the Fourier coefficients of meromorphic Jacobi forms have the same wall-crossing behavior as that exhibited by the degeneracies of BPS states.

The specific meromorphic Jacobi forms that are of interest for the black hole counting problem are the Fourier-Jacobi coefficients  $\psi_m(\tau, z)$  of the meromorphic Siegel modular form

$$\frac{1}{\Phi_{10}(\Omega)} = \sum_{m=-1}^{\infty} \psi_m(\tau, z) p^m, \quad \left( \Omega = \begin{pmatrix} \tau & z \\ z & \sigma \end{pmatrix}, p = e^{2\pi i \sigma} \right), \quad (1.1)$$

the reciprocal of the Igusa cusp form of weight 10, which arises as the partition function of quarter-BPS dyons in the type II compactification on the product of a  $K3$  surface and an elliptic curve [46, 98, 37]. These coefficients, after multiplication by the discriminant function  $\Delta(\tau)$ , are meromorphic Jacobi forms of weight 2 with a double pole at  $z = 0$  and no others (up to translation by the period lattice).

The new mathematical results of the paper are contained in Sections 8, 9 and 10. In §8, extending the results of [119], we show that any meromorphic Jacobi form  $\varphi(\tau, z)$  having poles only at torsion points  $z = \alpha\tau + \beta$  ( $\alpha, \beta \in \mathbb{Q}$ ) has a canonical decomposition into two pieces  $\varphi^F(\tau, z)$  and  $\varphi^P(\tau, z)$ , called its “finite” part and “polar” part, respectively, of which the first is a finite linear combination of classical theta series with mock modular forms as coefficients, and the second is an elementary expression that is determined completely by the poles of  $\varphi$ . Again using the results of Zwegers, we give explicit formulas for the polar part  $\varphi^P$  for all  $\varphi$  having only simple or double poles. In the particular case of  $\psi_m$ , the polar part is given by the formula

$$\psi_m^P(\tau, z) = \frac{p_{24}(m+1)}{\Delta(\tau)} \mathcal{A}_{2,m}(\tau, z) \quad (q = e^{2\pi i \tau}, y = e^{2\pi i z}), \quad (1.2)$$

where  $\Delta(\tau)$  is the Ramanujan discriminant function,  $p_{24}(m+1)$  the coefficient of  $q^m$  in  $\Delta(\tau)^{-1}$ , and  $\mathcal{A}_{2,m}(\tau, z)$  the elementary function (Appell-Lerch sum)

$$\mathcal{A}_{2,m}(\tau, z) = \sum_{s \in \mathbb{Z}} \frac{q^{ms^2+s} y^{2ms+1}}{(1 - q^s y)^2}. \quad (1.3)$$

Note that  $\mathcal{A}_{2,m}$  exhibits wall-crossing: for  $0 < \text{Im}(z) < \text{Im}(\tau)$  it has the Fourier expansion

$$\mathcal{A}_{2,m}(\tau, z) = \sum_{\substack{r \geq \ell > 0 \\ r \equiv \ell \pmod{2m}}} \ell q^{\frac{r^2 - \ell^2}{4m}} y^r, \quad (1.4)$$

but the Fourier expansions are different in other strips  $n < \text{Im}(z)/\text{Im}(\tau) < n + 1$ . This wall-crossing is at the heart of both the mathematical and the physical theories. On the mathematical side it explains the “mockness” of the finite part  $\psi_m^F$  of  $\psi_m$ . On the physical side it has an interpretation in terms of counting two-centered black holes, with the integer  $\ell$  in (1.4) being the dimension of the  $SU(2)$  multiplet with angular momentum  $(\ell - 1)/2$  contained in the electromagnetic field produced by the two centers.

Starting in §9 we focus attention on two particular classes of functions  $\{\varphi_{1,m}\}$  and  $\{\varphi_{2,m}\}$ , where  $\varphi_{k,m}$  is a meromorphic Jacobi form of weight  $k \in \{1, 2\}$  and index  $m \in \mathbb{N}$  having singular part  $(2\pi iz)^{-k} + O(1)$  as  $z \rightarrow 0$  and no other poles except the translates of this one by the period lattice. These functions in the case  $k = 2$  are related to the Fourier coefficients  $\psi_m$  defined in (1.1) by

$$\psi_m(\tau, z) = \frac{p_{24}(m+1)}{\Delta(\tau)} \varphi_{2,m}(\tau, z) + (\text{weakly holomorphic Jacobi form}). \quad (1.5)$$

The polar part of  $\varphi_{2,m}$  is the function  $\mathcal{A}_{2,m}$  defined above, and that of  $\varphi_{1,m}$  an Appell-Lerch sum  $\mathcal{A}_{1,m}$  with a similar definition (eq. (9.6)), but the functions themselves are not unique, since we can add to them any (weak) Jacobi forms of weight  $k$  and index  $m$  without changing the defining property. This will change the mock Jacobi form  $\Phi_{k,m} = \varphi_{k,m}^F$  in the corresponding way. Much of our analysis concerns finding “optimal” choices, meaning choices for which the functions  $\Phi_{k,m}$  have poles of as small an order as possible at infinity and consequently Fourier coefficients whose asymptotic growth is as small as possible. Our main results concern the case  $k = 2$ , and are as follows:

1. If  $m$  is a prime power, then the functions  $\Phi_{2,m}$  can be chosen to be strongly holomorphic (i.e., with Fourier expansions containing only monomials  $q^n y^r$  with  $4nm - r^2 \geq 0$ ). Their Fourier coefficients then have polynomial growth and are given by explicit linear combinations of class numbers of imaginary quadratic fields.
2. More generally, for arbitrary  $m$ , the function  $\Phi_{2,m}$  can be chosen to be the sum of the images under appropriate “Hecke-like operators” of special mock Jacobi forms  $\mathcal{Q}_M$ , where  $M$  ranges over all divisors of  $m$  having an even number of distinct prime factors. These summands are the eigencomponents of  $\Phi_{2,m}$  with respect to the Atkin-Lehner-like operators  $W_{m_1}$  ( $m_1|m$ ,  $(m_1, m/m_1) = 1$ ) acting on the space of mock Jacobi forms of index  $m$ .
3. The mock Jacobi form  $\mathcal{Q}_1$  is the generating function of class numbers of imaginary quadratic fields, and is strongly holomorphic. The other special mock Jacobi forms  $\mathcal{Q}_M$  ( $M = 6, 10, 14, \dots$ ) can be chosen to have “optimal growth” (meaning that their Fourier expansions contain only monomials  $q^n y^r$  with  $4nm - r^2 \geq -1$ ). Their Fourier coefficients then grow like  $e^{\pi\sqrt{\Delta}/m}$  as  $\Delta := 4nm - r^2$  tends to infinity.
4. One can also choose  $\Phi_{2,m}(\tau, z)$  for arbitrary  $m \geq 1$  to be of the form  $\sum \Phi_{2,m/d^2}^0(\tau, dz)$ , where  $d$  ranges over positive integers with  $d^2|m$  and each  $\Phi_{2,m/d^2}^0$  has optimal growth.
5. There are explicit formulas for the polar coefficients (those with  $\Delta < 0$ ) of  $\Phi_{2,m}^0$  and  $\mathcal{Q}_M$ . In particular, the coefficient of  $q^n y^r$  in  $\mathcal{Q}_M$  vanishes if  $4nM - r^2 = 0$  and equals  $\pm\varphi(M)/12$  if  $4nM - r^2 = -1$ , where  $\varphi(M) = \prod_{p|M} (p-1)$  is the Euler  $\varphi$ -function of  $M$ .

The proofs of these properties are contained in §10, which gives a detailed description (based on difficult results on Jacobi forms proved in [103, 104]) of the way that the space of holomorphic

Jacobi forms is contained in the spaces of weak or of weakly holomorphic Jacobi forms. This analysis contains several surprises, such as the result (Theorem 9.4) that for all  $m \geq 1$  the space of holomorphic Jacobi forms of index  $m$  has codimension exactly 1 in the space of Jacobi forms of index  $m$  with optimal growth.

For  $\varphi_{1,m}$  the description is much less complete. There is still a result like 2., with  $M$  now ranging over the divisors of  $m$  with an *odd* number of distinct prime factors, but it is no longer possible in general to choose the  $\mathcal{Q}_M$  to have optimal growth. The few cases where this is possible turn out to be related to mock theta functions that have played a prominent role in the past. Thus  $\mathcal{Q}_{30}$  and  $\mathcal{Q}_{42}$  are essentially equal to the mock theta functions of “order 5” and “order 7” in Ramanujan’s original letter to Hardy, and  $\mathcal{Q}_2$  and several of the other  $\mathcal{Q}_M$  are related to the “Mathieu moonshine” and “Umbral moonshine” story [47, 21]. The mock Jacobi forms  $\mathcal{Q}_M$  for the  $k = 2$  case are also related to special mock theta functions, now of weight  $3/2$ , e.g.  $\mathcal{Q}_6$  is the mock theta function with shadow  $\eta(\tau)$  given in [118].

The applications of the results (for the  $k = 2$  case, via the fact that  $\psi_m$  is the sum of  $p_{24}(m+1)\varphi_{2,m}/\Delta$  and a weakly holomorphic Jacobi form of weight  $-10$  and index  $m$ ) to the original physics problem are explained in §11 and in the following “second introduction.”

## 1.2 Introduction for physicists

The microscopic quantum description of supersymmetric black holes in string theory usually starts with a brane configuration of given charges and mass at weak coupling, which is localized at a single point in the noncompact spacetime. One then computes an appropriate indexed partition function in the world-volume theory of the branes, which from the perspective of enumerative geometry computes topological invariants such as the Donaldson-Thomas invariants. At strong coupling, the brane configuration gravitates and the indexed partition function is expected to count the microstates of the corresponding macroscopic gravitational configurations. Assuming that the gravitational configuration is a single-centered black hole then gives a way to obtain a statistical understanding of the entropy of the black hole in terms of its microstates, in accordance with the Boltzmann relation<sup>1</sup>.

One problem that one often encounters is that the macroscopic configurations are no longer localized at a point and include not only a single-centered black hole of interest but also several multi-centered ones [41, 42, 11, 43]. Moreover, the indexed degeneracy of the multi-centered configurations typically jumps upon crossing walls of marginal stability in the moduli space where the multi-centered configuration breaks up into its single-centered constituents. These jumps are referred to as the ‘wall-crossing phenomenon’.

If one is interested in the physics of the horizon or the microstates of a single black hole, the multi-centered configurations and the associated wall-crossings are thus something of a nuisance.

---

<sup>1</sup>Under certain conditions the index equals the absolute number [95, 96, 32].



It is desirable to have a mathematical characterization that isolates the single-centered black holes directly at the microscopic level. One distinguishing feature of single-centered black holes is that they are ‘*immortal*’ in that they exist as stable quantum states for all values of the moduli and hence their degeneracy does not exhibit the wall-crossing phenomenon. We will use this property later to define the counting function for the immortal black holes.

The wall-crossing phenomenon raises important conceptual questions regarding the proper holographic formulation in this context. In many cases, the black hole can be viewed as an excitation of a black string. The near horizon geometry of a black string is  $AdS_3$  which is expected to be holographically dual to a two-dimensional conformal field theory  $CFT_2$ . The conformal boundary of Euclidean  $AdS_3$  is a 2-torus with a complex structure parameter  $\tau$ , and the physical partition function of  $AdS_3$  and of the boundary  $CFT_2$  is a function of  $\tau$ . The  $SL(2, \mathbb{Z})$  transformations of  $\tau$  can be identified geometrically with global diffeomorphisms of the boundary of  $AdS_3$  space. The partition function is expected to have good modular properties under this geometric symmetry. This symmetry is crucial for the Rademacher-type expansions of the black hole degeneracies for understanding the quantum entropy of these black holes via holography [95, 96, 31, 44, 39, 81, 106, 86]. Implementing the modular symmetries presents several subtleties in situations when there is wall-crossing.

The wall-crossing phenomenon has another important physical implication for the invariance of the spectrum under large gauge transformations of the antisymmetric tensor field. Large gauge transformations lead to the ‘spectral flow symmetry’ of the partition function of the black string [39]. Since these transformations act both on the charges and the moduli, degeneracies of states with a charge vector  $\Gamma$  at some point  $\phi$  in the moduli space get mapped to the degeneracies of states with charge vector  $\Gamma'$  at some other point  $\phi'$  in the moduli space. Typically, there are many walls separating the point  $\phi'$  and the original point  $\phi$ . As a result, the degeneracies extracted from the black string at a *given* point  $\phi$  in the moduli space do not exhibit the spectral-flow symmetry. On the other hand, the spectrum of immortal black holes is independent of asymptotic moduli and hence must exhibit the spectral-flow symmetry. This raises the question as to how to make the spectral-flow symmetry manifest for the degeneracies of immortal black holes in the generic situation when there is wall-crossing.

With these motivations, our objective will be to isolate the partition functions of the black string associated with immortal black holes and investigate their transformation properties under the boundary modular group and large gauge transformations. More precisely, we would like to investigate the following four questions.

1. Can one define a *microscopic* counting function that cleanly isolates the microstates of immortal black holes from those of the multi-centered black configurations?
2. What are the modular properties of the counting function of immortal black holes when the asymptotic spectrum exhibits the wall-crossing phenomenon?



3. Can this counting function be related to a quantity that is properly modular as might be expected from the perspective of near-horizon  $AdS_3/CFT_2$  holography?
4. Can one define a partition function of the immortal black holes that manifestly exhibits the spectral-flow symmetry resulting from large gauge transformations?

The main difficulties in answering these questions stem from the complicated moduli dependence of the black hole spectrum which is often extremely hard to compute. To address the central conceptual issues in a tractable context, we consider the compactification of Type-II on  $K3 \times T^2$  with  $\mathcal{N} = 4$  supersymmetry in four dimensions. The spectrum of quarter-BPS dyonic states in this model is exactly computable [46, 55, 98, 99, 77, 37] and by now is well understood at *all* points in the moduli space [94, 27, 22] and for *all* possible duality orbits [27, 7, 8, 6, 29]. Moreover, as we will see, this particular model exhibits almost all of the essential issues that we wish to address. The  $\mathcal{N} = 4$  black holes have the remarkable property that even though their spectrum is moduli-dependent, the partition function itself is moduli-independent. The entire moduli dependence of the black hole degeneracy is captured by the moduli dependence of the choice of the Fourier contour [94, 27, 22]. Moreover, the only multi-centered configurations that contribute to the supersymmetric index of quarter-BPS states are the ones with only two centers, each of which is half-BPS [33]. Consequently, the only way the index can jump at a wall is by the decay of a two-centered configuration into its half-BPS constituents [94, 27, 22]; this is a considerable simplification compared to the general  $\mathcal{N} = 2$  case where more complicated multi-centered decays are possible. These features make the  $\mathcal{N} = 4$  case much more tractable.

The number of microstates of quarter-BPS dyonic states for the above-mentioned compactification is given by a Fourier coefficient of a meromorphic Jacobi form  $\psi_m(\tau, z)$  with a moduli-dependent contour. The partition function (1.1) referred to earlier is the generating function for these meromorphic Jacobi forms. Using this simplicity of the moduli dependence and the knowledge of the exact spectrum, it is possible to give very precise answers to the above questions in the  $\mathcal{N} = 4$  framework, which turn out to naturally involve mock modular forms.

1. One can define a holomorphic function for counting the microstates of immortal black holes<sup>2</sup> as a Fourier coefficient of the partition function of the black string for a specific choice of the Fourier contour [94, 27, 22]. The contour corresponds to choosing the asymptotic moduli of the theory in the attractor region of the single-centered black hole.
2. Because asymptotic counting function is a *meromorphic* Jacobi form, the near horizon counting function of immortal black holes is a *mock* modular form in that it fails to be

---

<sup>2</sup>We will use the terms ‘immortal’ and ‘single-centered’ interchangeably. In general, the moduli-independent ‘immortal’ degeneracies can receive contributions not only from single black holes but also from scaling solutions [43]. They are not expected to contribute to the  $\mathcal{N} = 4$  index that we consider [33]. In addition, there can be ‘hair’ degrees of freedom [10, 67], which are degrees of freedom localized outside the black hole horizon that carry part of the charge of the black hole. In frames where the black hole is represented entirely in terms of D-branes, such hair modes are expected to be absent.

modular but in a very specific way. The failure of modularity is governed by a *shadow*, which is given in terms of another holomorphic modular form.

3. Given a mock modular form and its shadow, one can define its *completion* which is a non-holomorphic modular form. The failure of holomorphy can be viewed as a ‘holomorphic anomaly’ which is also governed by the shadow.
4. The partition function of immortal black holes with manifest spectral-flow invariance is a mock Jacobi form – a new mathematical object defined and elaborated upon in §7.2.

The main physical payoff of the mathematics of mock modular forms in this context is the guarantee that one can still define a completion as in (3) which *is* modular albeit non-holomorphic. As mentioned earlier, the modular transformations on the  $\tau$  parameter can be identified with global diffeomorphisms of the boundary of the near horizon  $AdS_3$ . This connection makes the mathematics of mock modular forms physically very relevant for  $AdS_3/CFT_2$  holography in the presence of wall-crossing and holomorphic anomalies. The required mathematical results concerning mock modular forms are developed in sections §7, §8, §9, and §10.

To orient the physics reader, we summarize the essential conclusions of this mathematical investigation from the perspective of the questions posed above.

1. **Decomposition:** Given an asymptotic counting function, the degeneracies of single-centered black holes can be isolated using the results in §8. Applying Theorem 8.3 to the meromorphic Jacobi form  $\psi_m(\tau, z)$  gives a unique decomposition

$$\psi_m(\tau, z) = \psi_m^F(\tau, z) + \psi_m^P(\tau, z), \tag{1.6}$$

such that  $\psi_m^P(\tau, z)$  is a simple function (1.2) with the same pole structure in  $z$  as  $\psi_m(\tau, z)$  and  $\psi_m^F(\tau, z)$  has no poles. The elegant decomposition (1.6) is motivated partly by the choice of ‘attractor contour’ for single-centered black holes and has a direct physical interpretation:  $\psi_m(\tau, z)$  is the counting function of all asymptotic states including both single and multi-centered configurations,  $\psi_m^F(\tau, z)$  is the counting function of immortal black holes, whereas  $\psi_m^P(\tau, z)$  is the counting function of multi-centered black holes.

Since both  $\psi_m(\tau, z)$  and  $\psi_m^P(\tau, z)$  have poles in  $z$ , their Fourier coefficients depend on the choice of the contour which in turn depends on the moduli. On the other hand, the Fourier coefficients of  $\psi_m^F(\tau, z)$  are unambiguously defined without any contour or moduli dependence. This is what is expected for immortal black holes.

2. **Modular Completion:** The immortal counting function  $\psi_m^F(\tau, z)$  defined by the decomposition (1.6) is not modular. However, theorem 8.3 ensures that by adding a specific nonholomorphic function to  $\psi_m^F(\tau, z)$ , one can obtain its nonholomorphic completion

$\widehat{\psi}_m^{\text{F}}(\tau, z)$  which is modular and transforms as a Jacobi form. The failure of holomorphy of the completion  $\widehat{\psi}_m^{\text{F}}(\tau, z)$  is given by the equation

$$\tau_2^{3/2} \frac{\partial}{\partial \bar{\tau}} \widehat{\psi}_m^{\text{F}}(\tau, z) = \sqrt{\frac{m}{8\pi i}} \frac{p_{24}(m+1)}{\Delta(\tau)} \sum_{\ell \bmod (2m)} \overline{\vartheta_{m,\ell}(\tau, 0)} \vartheta_{m,\ell}(\tau, z). \quad (1.7)$$

Hence the counting function of immortal black holes has a hidden modular symmetry and more specifically is a mock Jacobi form as defined in §7.2. This is one of our main physics results and is described in more detail in §11.

3. **Holomorphic anomaly:** The completion is a natural object to be identified with the indexed partition function of the superconformal field theory  $SCFT_2$  dual to a single-centered  $AdS_3$ , which is expected to be modular.

From this perspective, the equation (1.7) can be viewed as a holomorphic anomaly and can in fact be taken as a defining property of a mock Jacobi form for physics applications. Naively, an indexed partition function is expected to be holomorphic because of a cancellation between right-moving bosons and fermions as for the elliptic genus [112]. However, if the  $SCFT_2$  is noncompact, then the spectrum is continuous and this naive reasoning may fail leading to an ‘anomaly’. The holomorphic anomaly can then arise as a consequence of the fact that for the right-movers in a noncompact  $SCFT$ , the density of states of bosons and fermions may be slightly different [108, 4] and may not precisely cancel. The detailed connection between the holomorphic anomaly and the noncompactness of the  $SCFT_2$  in this context needs to be understood better from a path-integral perspective.

4. **Optimality:** The Fourier coefficients of  $\psi_m^{\text{F}}(\tau, z)$  grow exponentially rapidly as expected for a counting function of black hole degeneracies. It is clear from the anomaly equation (1.7) that if we add a holomorphic true Jacobi form to  $\psi_m^{\text{F}}(\tau, z)$  with the same weight and index, it will still admit a modular completion satisfying the same anomaly equation. This raises the question whether for a given holomorphic anomaly there is an ‘optimal’ mock Jacobi form whose Fourier coefficients grow as slowly as possible. The answer to this question (for the functions  $\varphi_{2,m}$  related to  $\psi_m$  by (1.5)) is in the affirmative but is subtle and requires several new results in the theory of Jacobi forms developed in §10, motivated by and explaining the numerical experiments and observations described in §9.

A practical implication of such an optimal choice is that the leading contribution to the black hole entropy is then determined essentially by a Fourier coefficient of a true Jacobi form. One can thus apply the familiar Cardy formula and the Rademacher expansion of the Fourier coefficients of true modular forms for the leading answer. There will be

exponentially subleading corrections to this leading answer coming from the optimal mock Jacobi form. A Rademacher expansion for these corrections requires a generalization [13, 16] applicable for mock rather than true modular forms.

5. **Examples:** Modular forms with slowly growing Fourier coefficients are mathematically particularly interesting and many of the best-known examples of mock modular forms share this property. An ‘optimal’ choice thus enables us in many cases to obtain a very explicit expression for the immortal counting function  $\psi_m^F(\tau, z)$  in terms of these known mock modular forms. For example, for all  $m$  prime, the optimal mock part of  $\psi_m^F(\tau, z)$  can be expressed in terms of the generating function of Hurwitz-Kronecker class numbers (see §9.2). On the other hand, for a nonprime  $m$  our analysis leads to new mock modular forms with very slowly growing Fourier coefficients.

The functions  $\psi_m(\tau, z)$  that arise in the black hole problem have a double pole at  $z = 0$  and its translates, but our methods can also be applied to a related class of functions with just a single pole at  $z = 0$ . This leads to a second infinite family of mock modular forms, this time of weight  $1/2$ . (In the double pole case, the mock Jacobi forms had weight 2 and their coefficients were mock modular forms of weight  $3/2$ . In both cases, the coefficients are in fact mock theta functions, i.e., mock modular forms whose shadows are unary theta series.) Unlike the first case, where we found that the mock modular forms occurring can always be chosen to have at most simple poles at the cusps, the pole order here is much greater in general, and there are only a handful of examples having only simple poles. It is remarkable that these include essentially all the most prominent examples of mock theta functions, including the original ones of Ramanujan, the mock theta function conjecturally related to the Mathieu group  $M_{24}$  and the functions arising in the umbral moonshine conjecture [20].

Modular symmetries are very powerful in physics applications because they relate strong coupling to weak coupling, or high temperature to low temperature. The hidden modular symmetry of mock modular forms is therefore expected to be useful in diverse physics contexts. As mentioned above, mock modularity of the counting function in the present context of black holes is a consequence of meromorphy. In the previous two subsections and in this one, we have encountered several examples of  $\mathcal{Q}_M$  which, after of the asymptotic counting function which in turn is a consequence of noncompactness of the target space of the microscopic SCFT. Now, conformal field theories with a noncompact target space occur naturally in several physics contexts. For example, a general class of four-dimensional BPS black holes is obtained as a supersymmetric D-brane configuration in Type-II compactification on a Calabi-Yau three-fold  $X_6$ . In the M-theory limit, these black holes can be viewed as excitations of the MSW black string [79, 84]. The microscopic theory describing the low energy excitations of the MSW

string is the  $(0, 4)$  MSW SCFT. The target space of this SCFT does not necessarily have to be compact in which case the considerations of this paper will apply. Very similar objects [115, 65] have already made their appearance in the context of topological supersymmetric Yang-Mills theory on  $\mathbb{CP}^2$  [109]. Other examples include the theory of multiple M5-branes [2] quantum Liouville theory and E-strings [83], and the  $SL(2, \mathbb{R})/U(1)$  SCFT [49, 108, 4] where the CFT is noncompact.

We expect that the general framework of mock modular forms and Jacobi forms developed in this paper is likely to have varied physical applications in the context of non-compact conformal field theories, wall-crossings in enumerative geometry, and recently formulated Mathieu and umbral moonshine conjectures [47, 20].

### 1.3 Organization of the paper

In §2, we review the physics background concerning the string compactification on  $K3 \times T^2$  and the classification of BPS states corresponding to the supersymmetric black holes in this theory. In sections §3, §4, and §5, we review the basic mathematical definitions of various types of classical modular forms (elliptic, Jacobi, Siegel) and illustrate an application to the physics of quantum black holes in each case by means of an example. In §6, we review the moduli dependence of the Fourier contour prescription for extracting the degeneracies of quarter-BPS black holes in the  $\mathcal{N} = 4$  theory from the partition function which is a meromorphic Siegel modular form. In §8, we refine results due to Zwegers to show that any meromorphic Jacobi form with poles only at the sub-multiples of the period lattice can be decomposed canonically into two pieces, one of which is a mock Jacobi form. We give explicit formulas for this decomposition in the case when the poles have at most order 2, and again give several examples. In §9 we give a detailed description of the experimental results for the Fourier coefficients of the two families of mock Jacobi forms  $\{\Phi_{2,m}\}$  and  $\{\Phi_{1,m}\}$ , and formulate the main mathematical results concerning them. The proofs of these results are given in §10, after we have formulated and proved a number of structural results about holomorphic and weakly holomorphic Jacobi forms that are new and may be of independent interest. In §11, we apply these results in the physical context to determine the mock Jacobi form that counts the degeneracies of single-centered black holes and discuss the implications for  $AdS_2/CFT_1$  and  $AdS_3/CFT_2$  holography.

## 2. Review of Type-II superstring theory on $K3 \times T^2$

Superstring theories are naturally formulated in ten-dimensional Lorentzian spacetime  $\mathcal{M}_{10}$ . A ‘compactification’ to four-dimensions is obtained by taking  $\mathcal{M}_{10}$  to be a product manifold  $\mathbb{R}^{1,3} \times X_6$  where  $X_6$  is a compact Calabi-Yau threefold and  $\mathbb{R}^{1,3}$  is the noncompact Minkowski spacetime. We will focus in this paper on a compactification of Type-II superstring theory when  $X_6$  is itself the product  $X_6 = K3 \times T^2$ . A highly nontrivial and surprising result from the 90s is

the statement that this compactification is quantum equivalent or ‘dual’ to a compactification of heterotic string theory on  $T^4 \times T^2$  where  $T^4$  is a four-dimensional torus [66, 113]. One can thus describe the theory either in the Type-II frame or the heterotic frame.

The four-dimensional theory in  $\mathbb{R}^{1,3}$  resulting from this compactification has  $\mathcal{N} = 4$  supersymmetry<sup>3</sup>. The massless fields in the theory consist of 22 vector multiplets in addition to the supergravity multiplet. The massless moduli fields consist of the  $S$ -modulus  $\lambda$  taking values in the coset

$$SL(2, \mathbb{Z}) \backslash SL(2, \mathbb{R}) / O(2, \mathbb{R}), \quad (2.1)$$

and the  $T$ -moduli  $\mu$  taking values in the coset

$$O(22, 6, \mathbb{Z}) \backslash O(22, 6, \mathbb{R}) / O(22, \mathbb{R}) \times O(6, \mathbb{R}). \quad (2.2)$$

The group of discrete identifications  $SL(2, \mathbb{Z})$  is called the  $S$ -duality group. In the heterotic frame, it is the electro-magnetic duality group [90, 91], whereas in the type-II frame, it is simply the group of area-preserving global diffeomorphisms of the  $T^2$  factor. The group of discrete identifications  $O(22, 6, \mathbb{Z})$  is called the  $T$ -duality group. Part of the  $T$ -duality group  $O(19, 3, \mathbb{Z})$  can be recognized as the group of geometric identifications on the moduli space of K3; the other elements are stringy in origin and have to do with mirror symmetry.

At each point in the moduli space of the internal manifold  $K3 \times T^2$ , one has a distinct four-dimensional theory. One would like to know the spectrum of particle states in this theory. Particle states are unitary irreducible representations, or supermultiplets, of the  $\mathcal{N} = 4$  superalgebra. The supermultiplets are of three types which have different dimensions in the rest frame. A long multiplet is 256-dimensional, an intermediate multiplet is 64-dimensional, and a short multiplet is 16-dimensional. A short multiplet preserves half of the eight supersymmetries (*i.e.* it is annihilated by four supercharges) and is called a half-BPS state; an intermediate multiplet preserves one quarter of the supersymmetry (*i.e.* it is annihilated by two supercharges), and is called a quarter-BPS state; and a long multiplet does not preserve any supersymmetry and is called a non-BPS state. One consequence of the BPS property is that the spectrum of these states is ‘topological’ in that it does not change as the moduli are varied, except for jumps at certain walls in the moduli space [114].

An important property of a BPS states that follows from the superalgebra is that its mass is determined by its charges and the moduli [114]. Thus, to specify a BPS state at a given point in the moduli space, it suffices to specify its charges. The charge vector in this theory transforms in the vector representation of the  $T$ -duality group  $O(22, 6, \mathbb{Z})$  and in the

---

<sup>3</sup>This supersymmetry is a super Lie algebra containing  $ISO(1, 3) \times SU(4)$  as the even subalgebra where  $ISO(1, 3)$  is the Poincaré symmetry of the  $\mathbb{R}^{1,3}$  spacetime and  $SU(4)$  is an internal symmetry usually referred to as R-symmetry. The odd generators of the superalgebra are called supercharges. With  $\mathcal{N} = 4$  supersymmetry, there are eight complex supercharges which transform as a spinor of  $ISO(1, 3)$  and a fundamental of  $SU(4)$ .

fundamental representation of the  $S$ -duality group  $SL(2, \mathbb{Z})$ . It is thus given by a vector  $\Gamma^{\alpha I}$  with integer entries

$$\Gamma^{\alpha I} = \begin{pmatrix} N^I \\ M^I \end{pmatrix} \quad \text{where} \quad \alpha = 1, 2; \quad I = 1, 2, \dots, 28 \quad (2.3)$$

transforming in the  $(2, 28)$  representation of  $SL(2, \mathbb{Z}) \times O(22, 6, \mathbb{Z})$ . The vectors  $N$  and  $M$  can be regarded as the quantized electric and magnetic charge vectors of the state respectively. They both belong to an even, integral, self-dual lattice  $\Pi^{22,6}$ . We will assume in what follows that  $\Gamma = (N, M)$  in (2.3) is primitive in that it cannot be written as an integer multiple of  $(N_0, M_0)$  for  $N_0$  and  $M_0$  belonging to  $\Pi^{22,6}$ . A state is called purely electric if only  $N$  is non-zero, purely magnetic if only  $M$  is non-zero, and dyonic if both  $M$  and  $N$  are non-zero.

To define  $S$ -duality transformations, it is convenient to represent the  $S$ -modulus as a complex field  $S$  taking values in the upper half plane. An  $S$ -duality transformation

$$\gamma \equiv \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2; \mathbb{Z}) \quad (2.4)$$

acts simultaneously on the charges and the  $S$ -modulus by

$$\begin{pmatrix} N \\ M \end{pmatrix} \rightarrow \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} N \\ M \end{pmatrix}, \quad S \rightarrow \frac{aS + b}{cS + d}. \quad (2.5)$$

To define  $T$ -duality transformations, it is convenient to represent the  $T$ -moduli by a  $28 \times 28$  matrix  $\mu_I^A$  satisfying

$$\mu^t L \mu = L \quad (2.6)$$

with the identification that  $\mu \sim k\mu$  for every  $k \in O(22; \mathbb{R}) \times O(6; \mathbb{R})$ . Here  $L$  is the  $28 \times 28$  matrix

$$L_{IJ} = \begin{pmatrix} -\mathbf{C}_{16} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{I}_6 \\ \mathbf{0} & \mathbf{I}_6 & \mathbf{0} \end{pmatrix}, \quad (2.7)$$

with  $\mathbf{I}_s$  the  $s \times s$  identity matrix and  $\mathbf{C}_{16}$  is the Cartan matrix of  $E_8 \times E_8$ . The  $T$ -moduli are then represented by the matrix

$$\mathcal{M} = \mu^t \mu \quad (2.8)$$

which satisfies

$$\mathcal{M}^t = \mathcal{M}, \quad \mathcal{M}^t L \mathcal{M} = L. \quad (2.9)$$

In this basis, a  $T$ -duality transformation can then be represented by a  $28 \times 28$  matrix  $R$  with integer entries satisfying

$$R^t L R = L, \quad (2.10)$$



which acts simultaneously on the charges and the  $T$ -moduli by

$$N \rightarrow RN; \quad M \rightarrow RM; \quad \mu \rightarrow \mu R^{-1} \quad (2.11)$$

Given the matrix  $\mu_I^A$ , one obtains an embedding  $\Lambda^{22,6} \subset \mathbb{R}^{22,6}$  of  $\Pi^{22,6}$  which allows us to define the moduli-dependent charge vectors  $Q$  and  $P$  by

$$Q^A = \mu_I^A N_I, \quad P^A = \mu_I^A M_I. \quad (2.12)$$

The matrix  $L$  has a 22-dimensional eigensubspace with eigenvalue  $-1$  and a 6-dimensional eigensubspace with eigenvalue  $+1$ . Given  $Q$  and  $P$ , one can define the ‘right-moving’ and ‘left-moving’ charges<sup>4</sup>  $Q_{R,L}$  and  $P_{L,R}$  as the projections

$$Q_{R,L} = \frac{(1 \pm L)}{2} Q; \quad P_{R,L} = \frac{(1 \pm L)}{2} P. \quad (2.13)$$

If the vectors  $N$  and  $M$  are nonparallel, then the state is quarter-BPS. On the other hand, if  $N = pN_0$  and  $M = qN_0$  for some  $N_0 \in \Pi^{22,6}$  with  $p$  and  $q$  relatively prime integers, then the state is half-BPS.

An important piece of nonperturbative information about the dynamics of the theory is the exact spectrum of all possible dyonic BPS-states at all points in the moduli space. More specifically, one would like to compute the number  $d(\Gamma)|_{S,\mu}$  of dyons of a given charge  $\Gamma$  at a specific point  $(S, \mu)$  in the moduli space. Computation of these numbers is of course a very complicated dynamical problem. In fact, for a string compactification on a general Calabi-Yau threefold, the answer is not known. One main reason for focusing on this particular compactification on  $K3 \times T^2$  is that in this case the dynamical problem has been essentially solved and the exact spectrum of dyons is now known. Furthermore, the results are easy to summarize and the numbers  $d(\Gamma)|_{S,\mu}$  are given in terms of Fourier coefficients of various modular forms.

In view of the duality symmetries, it is useful to classify the inequivalent duality orbits labeled by various duality invariants. This leads to an interesting problem in number theory of classification of inequivalent duality orbits of various duality groups such as  $SL(2, \mathbb{Z}) \times O(22, 6; \mathbb{Z})$  in our case and more exotic groups like  $E_{7,7}(\mathbb{Z})$  for other choices of compactification manifold  $X_6$ . It is important to remember though that a duality transformation acts simultaneously on charges and the moduli. Thus, it maps a state with charge  $\Gamma$  at a point in the moduli space  $(S, \mu)$  to a state with charge  $\Gamma'$  but at some other point in the moduli space  $(S', \mu')$ . In this respect, the half-BPS and quarter-BPS dyons behave differently.

---

<sup>4</sup>The right-moving charges couple to the graviphoton vector fields associated with the right-moving chiral currents in the conformal field theory of the dual heterotic string.

- For half-BPS states, the spectrum does not depend on the moduli. Hence  $d(\Gamma)|_{S',\mu'} = d(\Gamma)|_{S,\mu}$ . Furthermore, by an  $S$ -duality transformation one can choose a frame where the charges are purely electric with  $M = 0$  and  $N \neq 0$ . Single-particle states have  $N$  primitive and the number of states depends only on the  $T$ -duality invariant integer  $n \equiv N^2/2$ . We can thus denote the degeneracy of half-BPS states  $d(\Gamma)|_{S',\mu'}$  simply by  $d(n)$ .
- For quarter-BPS states, the spectrum does depend on the moduli, and  $d(\Gamma)|_{S',\mu'} \neq d(\Gamma)|_{S,\mu}$ . However, the partition function turns out to be independent of moduli and hence it is enough to classify the inequivalent duality orbits to label the partition functions. For the specific duality group  $SL(2, \mathbb{Z}) \times O(22, 6; \mathbb{Z})$  the partition functions are essentially labeled by a single discrete invariant [27, 5, 6].

$$I = \text{gcd}(N \wedge M), \quad (2.14)$$

The degeneracies themselves are Fourier coefficients of the partition function. For a given value of  $I$ , they depend only on<sup>5</sup> the moduli and the three  $T$ -duality invariants  $(m, n, \ell) \equiv (M^2/2, N^2/2, N \cdot M)$ . Integrality of  $(m, n, \ell)$  follows from the fact that both  $N$  and  $M$  belong to  $\Pi^{22,6}$ . We can thus denote the degeneracy of these quarter-BPS states  $d(\Gamma)|_{S,\mu}$  simply by  $d(m, n, \ell)|_{S,\mu}$ . For simplicity, we consider only  $I = 1$  in this paper.

Given this classification, it is useful to choose a representative set of charges that can sample all possible values of the three  $T$ -duality invariants. For this purpose, we choose a point in the moduli space where the torus  $T^2$  is a product of two circles  $S^1 \times \tilde{S}^1$  and choose the following charges in a Type-IIB frame.

- For electric charges, we take  $n$  units of momentum along the circle  $S^1$ , and  $\tilde{K}$  Kaluza-Klein monopoles associated with the circle  $\tilde{S}^1$ .
- For magnetic charges, we take  $Q_1$  units of D1-brane charge wrapping  $S^1$ ,  $Q_5$  D5-brane wrapping  $K3 \times S^1$  and  $l$  units of momentum along the  $\tilde{S}^1$  circle.

We can thus write

$$\Gamma = \begin{bmatrix} N \\ M \end{bmatrix} = \begin{bmatrix} 0, n; 0, \tilde{K} \\ Q_1, \tilde{n}; Q_5, 0 \end{bmatrix}. \quad (2.15)$$

The  $T$ -duality quadratic invariants can be computed using a restriction of the matrix (2.7) to a  $\Lambda^{(2,2)}$  Narain lattice of the form

$$L = \begin{pmatrix} \mathbf{0} & \mathbf{I}_2 \\ \mathbf{I}_2 & \mathbf{0} \end{pmatrix}, \quad (2.16)$$

---

<sup>5</sup>There is an additional dependence on arithmetic  $T$ -duality invariants but the degeneracies for states with nontrivial values of these  $T$ -duality invariants can be obtained from the degeneracies discussed here by demanding  $S$ -duality invariance [6].

to obtain

$$M^2/2 = Q_1 Q_5, \quad N^2/2 = n\tilde{K}, \quad N \cdot M = \tilde{n}\tilde{K}. \quad (2.17)$$

We can simplify the notation further by choosing  $\tilde{K} = Q_5 = 1$ ,  $Q_1 = m$ ,  $\tilde{n} = l$  to obtain

$$M^2/2 = m, \quad N^2/2 = n, \quad N \cdot M = l. \quad (2.18)$$

For this set of charges, we can focus our attention on a subset of  $T$ -moduli associated with the torus  $T^2$  parametrized by

$$\mathcal{M} = \begin{pmatrix} G^{-1} & G^{-1}B \\ -BG^{-1} & G - BG^{-1}B \end{pmatrix}, \quad (2.19)$$

where  $G_{ij}$  is the metric on the torus and  $B_{ij}$  is the antisymmetric tensor field. Let  $U = U_1 + iU_2$  be the complex structure parameter,  $A$  be the area, and  $\epsilon_{ij}$  be the Levi-Civita symbol with  $\epsilon_{12} = -\epsilon_{21} = 1$ , then

$$G_{ij} = \frac{A}{U_2} \begin{pmatrix} 1 & U_1 \\ U_1 & |U|^2 \end{pmatrix} \quad \text{and} \quad B_{ij} = AB\epsilon_{ij}, \quad (2.20)$$

and the complexified Kähler modulus  $U = U_1 + iU_2$  is defined as  $U := B + iA$ . The  $S$ -modulus  $S = S_1 + S_2$  is defined as

$$S := a + i \exp(-2\phi) \quad (2.21)$$

where  $a$  is the axion and  $\phi$  is the dilaton field in the four dimensional heterotic frame. the relevant moduli can be parametrized by three complex scalars  $S, T, U$  which define the so-called ‘STU’ model in  $\mathcal{N} = 2$  supergravity. Note that these moduli are labeled naturally in the heterotic frame which are related to the  $S_B, T_B$ , and  $U_B$  moduli in the Type-IIB frame by

$$S = U_B, \quad T = S_B, \quad U = T_B. \quad (2.22)$$

### 3. Modular forms in one variable

Before discussing mock modular forms, it is useful to recall the variety of modular objects that have already made their appearance in the context of counting black holes. In the following sections we give the basic definitions of modular forms, Jacobi forms, and Siegel forms, using the notations that are standard in the mathematics literature, and then in each case illustrate a physics application to counting quantum black holes by means of an example.

In the physics context, these modular forms arise as generating functions for counting various quantum black holes in string theory. The structure of poles of the counting function is of particular importance in physics, since it determines the asymptotic growth of the Fourier coefficients as well as the contour dependence of the Fourier coefficients which corresponds to the wall crossing phenomenon. These examples will also be relevant later in §11 in connection with mock modular forms. We suggest chapters I and III of [17] respectively as a good general reference for classical and Siegel modular forms and [52] for Jacobi modular forms.

### 3.1 Basic definitions and properties

Let  $\mathbb{H}$  be the upper half plane, *i.e.*, the set of complex numbers  $\tau$  whose imaginary part satisfies  $\text{Im}(\tau) > 0$ . Let  $SL(2, \mathbb{Z})$  be the group of matrices  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  with integer entries such that  $ad - bc = 1$ .

A *modular form*  $f(\tau)$  of weight  $k$  on  $SL(2, \mathbb{Z})$  is a holomorphic function on  $\mathbb{H}$ , that transforms as

$$f\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^k f(\tau) \quad \forall \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}), \quad (3.1)$$

for an integer  $k$  (necessarily even if  $f(0) \neq 0$ ). It follows from the definition that  $f(\tau)$  is periodic under  $\tau \rightarrow \tau + 1$  and can be written as a Fourier series

$$f(\tau) = \sum_{n=-\infty}^{\infty} a(n) q^n \quad (q := e^{2\pi i \tau}), \quad (3.2)$$

and is bounded as  $\text{Im}(\tau) \rightarrow \infty$ . If  $a(0) = 0$ , then the modular form vanishes at infinity and is called a *cusp form*. Conversely, one may weaken the growth condition at  $\infty$  to  $f(\tau) = O(q^{-N})$  rather than  $O(1)$  for some  $N \geq 0$ ; then the Fourier coefficients of  $f$  have the behavior  $a(n) = 0$  for  $n < -N$ . Such a function is called a *weakly holomorphic modular form*.

The vector space over  $\mathbb{C}$  of holomorphic modular forms of weight  $k$  is usually denoted by  $M_k$ . Similarly, the space of cusp forms of weight  $k$  and the space of weakly holomorphic modular forms of weight  $k$  are denoted by  $S_k$  and  $M_k^!$  respectively. We thus have the inclusion

$$S_k \subset M_k \subset M_k^!. \quad (3.3)$$

The Fourier coefficients of the modular forms in these spaces have different growth properties:

1.  $f \in S_k \Rightarrow a_n = O(n^{k/2})$  as  $n \rightarrow \infty$ ;
2.  $f \in M_k \Rightarrow a_n = O(n^{k-1})$  as  $n \rightarrow \infty$ ;
3.  $f \in M_k^! \Rightarrow a_n = O(e^{C\sqrt{n}})$  as  $n \rightarrow \infty$  for some  $C > 0$ .

Some important modular forms on  $SL(2, \mathbb{Z})$  are:

1. The *Eisenstein series*  $E_k \in M_k$  ( $k \geq 4$ ). The first two of these are

$$E_4(\tau) = 1 + 240 \sum_{n=1}^{\infty} \frac{n^3 q^n}{1 - q^n} = 1 + 240q + 2160q^2 + \dots, \quad (3.4)$$

$$E_6(\tau) = 1 - 504 \sum_{n=1}^{\infty} \frac{n^5 q^n}{1 - q^n} = 1 - 504q - 16632q^2 - \dots. \quad (3.5)$$

2. The *discriminant function*  $\Delta$ . It is given by the product expansion

$$\Delta(\tau) = q \prod_{n=1}^{\infty} (1 - q^n)^{24} = q - 24q^2 + 252q^3 + \dots \quad (3.6)$$

or by the formula  $\Delta = (E_4^3 - E_6^2)/1728$ . We mention for later use that the function  $E_2(\tau) = \frac{1}{2\pi i} \frac{\Delta'(\tau)}{\Delta(\tau)} = 1 - 24 \sum_{n=1}^{\infty} \frac{nq^n}{1 - q^n}$  is also an Eisenstein series, but is not modular. (It is a so-called *quasimodular form*, meaning in this case that the non-holomorphic function  $\widehat{E}_2(\tau) = E_2(\tau) - \frac{3}{\pi \operatorname{Im}(\tau)}$  transforms like a modular form of weight 2.) This function can be used to form the *Ramanujan-Serre derivative*

$$\partial_k^{\text{RS}} : M_k \rightarrow M_{k+2}, \quad \partial_k^{\text{RS}} f(\tau) := \frac{1}{2\pi i} f'(\tau) - \frac{k}{12} E_2(\tau) f(\tau). \quad (3.7)$$

The ring of modular forms on  $SL(2, \mathbb{Z})$  is generated freely by  $E_4$  and  $E_6$ , so any modular form of weight  $k$  can be written (uniquely) as a sum of monomials  $E_4^\alpha E_6^\beta$  with  $4\alpha + 6\beta = k$ . We also have  $M_k = \mathbb{C} \cdot E_k \oplus S_k$  and  $S_k = \Delta \cdot M_{k-12}$ , so any  $f \in M_k$  also has a unique expansion as  $\sum_{n=0}^{\lfloor k/12 \rfloor} \alpha_n E_{k-12n} \Delta^n$  (with  $E_0 = 1$  and  $E_2$  replaced by 0). From either representation, we see that a modular form is uniquely determined by its weight and first few Fourier coefficients.

Given two modular forms  $(f, g)$  of weight  $(k, l)$ , one can produce a sequence of modular forms of weight  $k + l + 2n$ ,  $n \geq 0$  using the *Rankin-Cohen bracket*

$$[f, g]_n = [f, g]_n^{(k,l)} = \sum_{r+s=n} (-1)^s \binom{k+n-1}{r} \binom{\ell+n-1}{s} f^{(s)}(\tau) g^{(r)}(\tau) \quad (3.8)$$

where  $f^{(m)} := \left(\frac{1}{2\pi i} \frac{d}{d\tau}\right)^m f$ . For  $n = 0$ , this is simply the product of the two forms, while for  $n > 0$  we always have  $[f, g]_n \in S_{k+l+2n}$ . The first two non-trivial examples are

$$[E_4, E_6]_1 = -3456 \Delta, \quad [E_4, E_4]_2 = 4800 \Delta. \quad (3.9)$$

As well as modular forms on the full modular group  $SL(2, \mathbb{Z})$ , one can also consider modular forms on subgroups of finite index, with the same transformation law (3.1) and suitable conditions on the Fourier coefficients to define the notions of holomorphic, weakly holomorphic and cusp forms. The weight  $k$  now need no longer be even, but can be odd or even half integral, the easiest way to state the transformation property when  $k \in \mathbb{Z} + \frac{1}{2}$  being to say that  $f(\tau)/\theta(\tau)^{2k}$  is invariant under some congruence subgroup of  $SL(2, \mathbb{Z})$ , where  $\theta(\tau) = \sum_{n \in \mathbb{Z}} q^{n^2}$ . The graded vector space of modular forms on a fixed subgroup  $\Gamma \subset SL(2, \mathbb{Z})$  is finite dimensional in each weight, finitely generated as an algebra, and closed under Rankin-Cohen brackets. Important

examples of modular forms of half-integral weight are the *unary theta series*, i.e., theta series associated to a quadratic form in one variable. They come in two types:

$$\sum_{n \in \mathbb{Z}} \varepsilon(n) q^{\lambda n^2} \quad \text{for some } \lambda \in \mathbb{Q}_+ \text{ and some even periodic function } \varepsilon \quad (3.10)$$

and

$$\sum_{n \in \mathbb{Z}} n \varepsilon(n) q^{\lambda n^2} \quad \text{for some } \lambda \in \mathbb{Q}_+ \text{ and some odd periodic function } \varepsilon, \quad (3.11)$$

the former being a modular form of weight  $1/2$  and the latter a cusp form of weight  $3/2$ . A theorem of Serre and Stark says that in fact every modular form of weight  $1/2$  is a linear combination of form of the type (3.10), a simple example being the identity

$$\eta(\tau) := q^{1/24} \prod_{n=1}^{\infty} (1 - q^n) = \sum_{n=1}^{\infty} \chi_{12}(n) q^{n^2/24}, \quad (3.12)$$

proved by Euler for the so-called *Dedekind eta function*  $\eta(t) = \Delta(\tau)^{1/24}$ . Here  $\chi_{12}$  is the function of period 12 defined by

$$\chi_{12}(n) = \begin{cases} +1 & \text{if } n \equiv \pm 1 \pmod{12} \\ -1 & \text{if } n \equiv \pm 5 \pmod{12} \\ 0 & \text{if } (n, 12) > 1. \end{cases} \quad (3.13)$$

Finally, we recall the definition of the Petersson scalar product. If  $f(\tau)$  and  $g(\tau)$  are two modular forms of the same weight  $k$  and the same “multiplier system” on some subgroup  $\Gamma$  of finite index of  $SL_2(\mathbb{Z})$  (this means that the quotient  $f/g$  is invariant under  $\Gamma$ ), and if either  $k < 1$  or else at least one of  $f$  and  $g$  is a cusp form, then we can define the (normalized) *Petersson scalar product* of  $f$  and  $g$  by

$$(f, g) = \int_{\mathcal{F}_\Gamma} f(\tau) \overline{g(\tau)} \tau_2^k d\mu(\tau) \Big/ \int_{\mathcal{F}_\Gamma} d\mu(\tau) \quad \left( d\mu(\tau) := \frac{d\tau_1 d\tau_2}{\tau_2^2} \right), \quad (3.14)$$

where  $\mathcal{F}_\Gamma$  is a fundamental domain for  $\Gamma$ . This definition is independent of the choice of the subgroup  $\Gamma$  and the fundamental domain  $\mathcal{F}_\Gamma$ . By the Rankin-Selberg formula (see, for example, [116]), we have that if  $f = \sum_{\lambda} a_{\lambda} q^{\lambda}$  and  $g = \sum_{\lambda} b_{\lambda} q^{\lambda}$  (where  $\lambda \geq 0$  may have a denominator), then

$$(f, g) = \frac{\Gamma(k)}{(4\pi)^k} \operatorname{Res}_{s=k} \left( \sum_{\lambda > 0} \frac{a_{\lambda} \overline{b_{\lambda}}}{\lambda^s} \right), \quad (3.15)$$

a formula that will be used later. For instance, for  $f = g = \eta$  we have  $k = 1/2$  and  $\sum_{\lambda} a_{\lambda} b_{\lambda} \lambda^{-s} = 24^s (1 - 2^{-2s})(1 - 3^{-2s}) \zeta(2s)$ , and hence  $(\eta, \eta) = 1/\sqrt{6}$ .

### 3.2 Quantum black holes and modular forms

Modular forms occur naturally in the context of counting the Dabholkar-Harvey (DH) states [34, 28], which are states in the string Hilbert space that are dual to perturbative BPS states. The spacetime helicity supertrace counting the degeneracies reduces to the partition function of a chiral conformal field theory on a genus-one worldsheet. The  $\tau$  parameter above becomes the modular parameter of the genus one Riemann surface. The degeneracies are given by the Fourier coefficients of the partition function.

A well-known simple example is the partition function  $Z(\tau)$  which counts the half-BPS DH states for the Type-II compactification on  $K3 \times T^2$  considered here. In the notation of (2.3) these states have zero magnetic charge  $M = 0$ , but nonzero electric charge  $N$  with the  $T$ -duality invariant  $N^2 = 2n$ , which can be realized for example by setting  $Q_1 = Q_5 = l = 0$  in (2.15). They are thus purely electric and perturbative in the heterotic frame<sup>6</sup>. The partition function is given by the partition function of the chiral conformal field theory of 24 left-moving transverse bosons of the heterotic string. The Hilbert space  $\mathcal{H}$  of this theory is a unitary Fock space representation of the commutation algebra

$$[a_{in}, a_{jm}^\dagger] = \delta_{ij} \delta_{n+m,0} \quad (i, j = 1, \dots, 24, \quad n, m = 1, 2, \dots, \infty) \quad (3.16)$$

of harmonic modes of oscillations of the string in 24 different directions. The Hamiltonian is

$$H = \sum_{i=1}^{24} n a_{in}^\dagger a_{in} - 1, \quad (3.17)$$

and the partition function is

$$Z(\tau) = \text{Tr}_{\mathcal{H}}(q^H). \quad (3.18)$$

This can be readily evaluated since each oscillator mode of energy  $n$  contributes to the trace

$$1 + q^n + q^{2n} + \dots = \frac{1}{1 - q^n}. \quad (3.19)$$

The partition function then becomes

$$Z(\tau) = \frac{1}{\Delta(\tau)}, \quad (3.20)$$

where  $\Delta$  is the cusp form (3.6). Since  $\Delta$  has a simple zero at  $q = 0$ , the partition function itself has a pole at  $q = 0$ , but has no other poles in  $\mathbb{H}$ . Hence,  $Z(\tau)$  is a weakly holomorphic

---

<sup>6</sup>Not all DH states are half-BPS. For example, the states that are perturbative in the Type-II frame correspond to a Type-II string winding and carrying momentum along a cycle in  $T^2$ . For such states both  $M$  and  $N$  are nonzero and nonparallel, and hence the state is quarter-BPS.



modular form of weight  $-12$ . This property is essential in the present physical context since it determines the asymptotic growth of the Fourier coefficients.

The degeneracy  $d(n)$  of the state with electric charge  $N$  depends only on the  $T$ -duality invariant integer  $n$  and is given by

$$Z(\tau) = \sum_{n=-1}^{\infty} d(n) q^n. \quad (3.21)$$

For the Fourier integral

$$d(n) = \int_{\mathcal{C}} e^{-2\pi i \tau n} Z(\tau) d\tau, \quad (3.22)$$

one can choose the contour  $\mathcal{C}$  in  $\mathbb{H}$  to be

$$0 \leq \text{Re}(\tau) < 1, \quad (3.23)$$

for a fixed imaginary part  $\text{Im}(\tau)$ . Since the partition function has no poles in  $\mathbb{H}$  except at  $q = 0$ , smooth deformations of the contour do not change the Fourier coefficients and consequently the degeneracies  $d(n)$  are uniquely determined from the partition function. This reflects the fact that the half-BPS states are immortal and do not decay anywhere in the moduli space. As a result, there is no wall crossing phenomenon, and no jumps in the degeneracy.

In number theory, the partition function above is well-known in the context of the problem of partitions of integers. We can therefore identify

$$d(n) = p_{24}(n+1) \quad (n \geq 0). \quad (3.24)$$

where  $p_{24}(I)$  is the number of colored partitions of a positive integer  $I$  using integers of 24 different colors.

These states have a dual description in the Type-II frame where they can be viewed as bound states of  $Q_1$  number of D1-branes and  $Q_5$  number of D5-branes with  $M^2/2 = Q_1 Q_5 \equiv m$ . This corresponds to setting  $n = \tilde{K} = l = 0$  in (2.15). In this description, the number of such bound states  $d(m)$  equals the orbifold Euler character  $\chi(\text{Sym}^{m+1}(K3))$  of the symmetric product of  $(m+1)$  copies of  $K3$ -surface [109]. The generating function for the orbifold Euler character

$$\hat{Z}(\sigma) = \sum_{m=-1}^{\infty} \chi(\text{Sym}^{m+1}(K3)) p^m \quad (p := e^{2\pi i \sigma}) \quad (3.25)$$

can be evaluated [59] to obtain

$$\hat{Z}(\sigma) = \frac{1}{\Delta(\sigma)}. \quad (3.26)$$

Duality requires that the number of immortal BPS-states of a given charge must equal the number of BPS-states with the dual charge. The equality of the two partition functions (3.20)

and (3.26) coming from two very different counting problems is consistent with this expectation. This fact was indeed one of the early indications of a possible duality between heterotic and Type-II strings [109].

The DH-states correspond to the microstates of a small black hole [92, 24, 35] for large  $n$ . The macroscopic entropy  $S(n)$  of these black holes should equal the asymptotic growth of the degeneracy by the Boltzmann relation

$$S(n) = \log d(n); \quad n \gg 1. \quad (3.27)$$

In the present context, the macroscopic entropy can be evaluated from the supergravity solution of small black holes [73, 76, 75, 74, 24, 35]. The asymptotic growth of the microscopic degeneracy can be evaluated using the Hardy-Ramanujan expansion (Cardy formula). There is a beautiful agreement between the two results [24, 71]

$$S(n) = \log d(n) \sim 4\pi\sqrt{n} \quad n \gg 1. \quad (3.28)$$

Given the growth properties of the Fourier coefficients mentioned above, it is clear that, for a black hole whose entropy scales as a power of  $n$  and not as  $\log(n)$ , the partition function counting its microstates can be only weakly holomorphic and not holomorphic.

These considerations generalize in a straightforward way to congruence subgroups of  $SL(2, \mathbb{Z})$  which are relevant for counting the DH-states in various orbifold compactifications with  $\mathcal{N} = 4$  or  $\mathcal{N} = 2$  supersymmetry [25, 93, 26].

## 4. Jacobi forms

### 4.1 Basic definitions

Consider a holomorphic function  $\varphi(\tau, z)$  from  $\mathbb{H} \times \mathbb{C}$  to  $\mathbb{C}$  which is “modular in  $\tau$  and elliptic in  $z$ ” in the sense that it transforms under the modular group as

$$\varphi\left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}\right) = (c\tau + d)^k e^{\frac{2\pi imcz^2}{c\tau + d}} \varphi(\tau, z) \quad \forall \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2; \mathbb{Z}) \quad (4.1)$$

and under the translations of  $z$  by  $\mathbb{Z}\tau + \mathbb{Z}$  as

$$\varphi(\tau, z + \lambda\tau + \mu) = e^{-2\pi im(\lambda^2\tau + 2\lambda z)} \varphi(\tau, z) \quad \forall \quad \lambda, \mu \in \mathbb{Z}, \quad (4.2)$$

where  $k$  is an integer and  $m$  is a positive integer.

These equations include the periodicities  $\varphi(\tau + 1, z) = \varphi(\tau, z)$  and  $\varphi(\tau, z + 1) = \varphi(\tau, z)$ , so  $\varphi$  has a Fourier expansion

$$\varphi(\tau, z) = \sum_{n,r} c(n, r) q^n y^r, \quad (q := e^{2\pi i\tau}, y := e^{2\pi iz}). \quad (4.3)$$

Equation (4.2) is then equivalent to the periodicity property

$$c(n, r) = C(4nm - r^2, r), \quad \text{where } C(\Delta, r) \text{ depends only on } r \pmod{2m}. \quad (4.4)$$

(We will sometimes denote  $c(n, r)$  and  $C(\Delta, r)$  by  $c_\varphi(n, r)$  and  $C_\varphi(\Delta, r)$  or by  $c(\varphi; n, r)$  and  $C(\varphi; \Delta, r)$  when this is required for emphasis or clarity.) The function  $\varphi(\tau, z)$  is called a *holomorphic Jacobi form* (or simply a *Jacobi form*) of weight  $k$  and index  $m$  if the coefficients  $C(\Delta, r)$  vanish for  $\Delta < 0$ , *i.e.* if

$$c(n, r) = 0 \quad \text{unless} \quad 4mn \geq r^2. \quad (4.5)$$

It is called a *Jacobi cusp form* if it satisfies the stronger condition that  $C(\Delta, r)$  vanishes unless  $\Delta$  is strictly positive, *i.e.*

$$c(n, r) = 0 \quad \text{unless} \quad 4mn > r^2, \quad (4.6)$$

and it is called a *weak Jacobi form* if it satisfies the weaker condition

$$c(n, r) = 0 \quad \text{unless} \quad n \geq 0 \quad (4.7)$$

rather than (4.5), whereas a merely *weakly holomorphic Jacobi form* satisfies only the yet weaker condition that  $c(n, r) = 0$  unless  $n \geq n_0$  for some possibly negative integer  $n_0$  (or equivalently  $C(\Delta, r) = 0$  unless  $\Delta \geq \Delta_0$  for some possibly negative integer  $\Delta_0$ ). The space of all holomorphic (resp. cuspidal, weak, or weakly holomorphic) Jacobi forms of weight  $k$  and index  $m$  will be denoted by  $J_{k,m}$  (resp.  $J_{k,m}^0$ ,  $\tilde{J}_{k,m}$ , or  $\tilde{J}_{k,m}^1$ ).

Finally, the quantity  $\Delta = 4mn - r^2$ , which by virtue of the above discussion is the crucial invariant of a monomial  $q^n y^r$  occurring in the Fourier expansion of  $\varphi$ , will be referred to as its *discriminant*. (It would be mathematically more correct to use this word for the quantity  $-\Delta$ , but  $\Delta$  is usually positive and it is more convenient to work with positive numbers.)

## 4.2 Theta expansion and Taylor expansion

A Jacobi form has two important representations, the *theta expansion* and the *Taylor expansion*. In this subsection, we explain both of these and the relation between them.

If  $\varphi(\tau, z)$  is a Jacobi form, then the transformation property (4.2) implies its Fourier expansion with respect to  $z$  has the form

$$\varphi(\tau, z) = \sum_{\ell \in \mathbb{Z}} q^{\ell^2/4m} h_\ell(\tau) e^{2\pi i \ell z} \quad (4.8)$$

where  $h_\ell(\tau)$  is periodic in  $\ell$  with period  $2m$ . In terms of the coefficients (4.4) we have

$$h_\ell(\tau) = \sum_{\Delta} C(\Delta, \ell) q^{\Delta/4m} \quad (\ell \in \mathbb{Z}/2m\mathbb{Z}). \quad (4.9)$$

Because of the periodicity property, equation (4.8) can be rewritten in the form

$$\varphi(\tau, z) = \sum_{\ell \in \mathbb{Z}/2m\mathbb{Z}} h_\ell(\tau) \vartheta_{m,\ell}(\tau, z), \quad (4.10)$$

where  $\vartheta_{m,\ell}(\tau, z)$  denotes the standard index  $m$  theta function

$$\vartheta_{m,\ell}(\tau, z) := \sum_{\substack{r \in \mathbb{Z} \\ r \equiv \ell \pmod{2m}}} q^{r^2/4m} y^r = \sum_{n \in \mathbb{Z}} q^{(\ell+2mn)^2/4m} y^{\ell+2mn} \quad (4.11)$$

(which is a Jacobi form of weight  $\frac{1}{2}$  and index  $m$  on some subgroup of  $SL(2, \mathbb{Z})$ ). This is the theta expansion of  $\varphi$ . The coefficients  $h_\ell(\tau)$  are modular forms of weight  $k - \frac{1}{2}$  and are weakly holomorphic, holomorphic or cuspidal if  $\varphi$  is a weak Jacobi form, a Jacobi form or a Jacobi cusp form, respectively. More precisely, the vector  $h := (h_1, \dots, h_{2m})$  transforms like a modular form of weight  $k - \frac{1}{2}$  under  $SL(2, \mathbb{Z})$ .

The theta decomposition (4.10) leads to the definition of a differential operator on Jacobi forms as follows. Let

$$\mathcal{L}_m = \frac{4m}{2\pi i} \frac{\partial}{\partial \tau} - \frac{1}{(2\pi i)^2} \frac{\partial^2}{\partial z^2}$$

be the index  $m$  heat operator, which sends  $\varphi = \sum c(n, r) q^n y^r$  to  $\sum (4nm - r^2) c(n, r) q^n y^r$ . Here the Fourier coefficients have the same periodicity property (4.4) as for  $\varphi$ , so  $\mathcal{L}_m \varphi$  has the same elliptic transformation properties as a Jacobi form of index  $m$ . Moreover, since  $\mathcal{L}_m$  annihilates all the  $\vartheta_{m,\ell}$ , we have  $\mathcal{L}_m(\sum h_\ell \vartheta_{m,\ell}) = 4m \sum h'_\ell \vartheta_{m,\ell}$ , so the modified heat operator

$$\mathcal{L}_{k,m} = \mathcal{L}_m - \frac{m(k - \frac{1}{2})}{3} E_2 : \sum_{\ell} h_\ell(\tau) \vartheta_{m,\ell}(\tau, z) \mapsto 4m \sum_{\ell} \partial_{k-\frac{1}{2}}^{\text{RS}} h_\ell(\tau) \vartheta_{m,\ell}(\tau, z), \quad (4.12)$$

where  $\partial_{*}^{\text{RS}}$  is the Ramanujan-Serre derivative defined in (3.7), sends  $J_{k,m}$  to  $J_{k+2,m}$  (and also  $\tilde{J}_{k,m}$  to  $\tilde{J}_{k+2,m}$  and  $J_{k,m}^!$  to  $J_{k+2,m}^!$ ). These operators will be used later.

A Jacobi form  $\varphi \in \tilde{J}_{k,m}$  also has a Taylor expansion in  $z$  which for  $k$  even takes the form

$$\varphi(\tau, z) = \xi_0(\tau) + \left( \frac{\xi_1(\tau)}{2} + \frac{m\xi'_0(\tau)}{k} \right) (2\pi iz)^2 + \left( \frac{\xi_2(\tau)}{24} + \frac{m\xi'_1(\tau)}{2(k+2)} + \frac{m^2\xi''_0(\tau)}{2k(k+1)} \right) (2\pi iz)^4 + \dots \quad (4.13)$$

with  $\xi_\nu \in M_{k+2\nu}(SL(2, \mathbb{Z}))$  and the prime denotes  $\frac{1}{2\pi i} \frac{d}{d\tau}$  as before. In terms of the Fourier coefficients of  $\varphi$ , the modular form  $\xi_\nu$  is given by

$$\frac{(k+2\nu-2)!}{(k+\nu-2)!} \xi_\nu(\tau) = \sum_{n=0}^{\infty} \left( \sum_r P_{\nu,k}(nm, r^2) c(n, r) \right) q^n, \quad (4.14)$$

where  $P_{\nu,k}$  is a homogeneous polynomial of degree  $\nu$  in  $r^2$  and  $n$  with coefficients depending on  $k$  and  $m$ , the first few being

$$\begin{aligned} P_{0,k} &= 1, \\ P_{1,k} &= kr^2 - 2nm, \\ P_{2,k} &= (k+1)(k+2)r^4 - 12(k+1)r^2mn + 12m^2n^2. \end{aligned} \quad (4.15)$$

The Jacobi form  $\varphi$  is determined by the first  $m+1$  coefficients  $\xi_\nu$ , and the map  $\varphi \mapsto (\xi_0, \dots, \xi_m)$  is an isomorphism from  $\tilde{J}_{k,m}$  to  $M_k \oplus M_{k+2} \oplus \dots \oplus M_{k+2m}$ . For  $k$  odd, the story is similar except that (4.13) must be replaced by

$$\varphi(\tau, z) = \xi_0(\tau) (2\pi iz) + \left( \frac{\xi_1(\tau)}{6} + \frac{m\xi_0'(\tau)}{k+2} \right) (2\pi iz)^3 + \dots \quad (4.16)$$

with  $\xi_\nu \in M_{k+2\nu+1}(SL(2, \mathbb{Z}))$ , and the map  $\varphi \mapsto (\xi_0, \dots, \xi_{m-2})$  gives an isomorphism from  $\tilde{J}_{k,m}$  to  $M_{k+1} \oplus M_{k+3} \oplus \dots \oplus M_{k+2m-3}$ .

We also observe that even if  $\varphi$  is weak, so that the individual coefficients  $c(n, r)$  grow like  $C\sqrt{4nm-r^2}$ , the coefficients  $\sum_r P_{\nu,k}(nm, r^2) c(n, r)$  of  $\xi_\nu$  still have only polynomial growth. We thus have the following descriptions (analogous to those given for classical modular forms in §3.1) of holomorphic and weak Jacobi forms in terms of their asymptotic properties:

$$\begin{aligned} \text{Holomorphic Jacobi} &\iff c(n, r) = 0 \text{ for } 4mn - r^2 < 0 \\ &\iff \text{the function } q^{m\alpha^2} \varphi(\tau, \alpha\tau + \beta) \text{ (which is a modular form of weight } k \\ &\quad \text{and some level) is bounded as } \tau_2 \rightarrow \infty \text{ for every } \alpha, \beta \in \mathbb{Q} \\ &\iff \text{all } h_j(\tau) \text{ in (4.10) are bounded as } \tau_2 \rightarrow \infty \\ &\iff c(n, r) \text{ have polynomial growth.} \end{aligned}$$

$$\begin{aligned} \text{Weak Jacobi} &\iff c(n, r) = 0 \text{ for } n < 0 \\ &\iff \varphi(\tau, \alpha\tau + \beta) \text{ is bounded as } \tau_2 \rightarrow \infty \text{ for any fixed } z \in \mathbb{C} \\ &\iff \text{all } h_j(\tau) = O(q^{-j^2/4m}) \text{ as } \tau_2 \rightarrow \infty \\ &\iff \sum_r P_{\nu,k}(nm, r^2) c(n, r) \text{ have polynomial growth.} \end{aligned}$$

Finally, the relation between the Taylor expansion (4.13) of a Jacobi form and its theta expansion (4.10) is given by

$$\xi_\nu(\tau) = (4m)^\nu \binom{k+2\nu-2}{\nu}^{-1} \sum_{\ell \pmod{2m}} [h_\ell(\tau), \vartheta_{m,\ell}^0(\tau)]_\nu, \quad (4.17)$$

where  $[\ , ]_\nu = [\ , ]_\nu^{(k-\frac{1}{2}, \frac{1}{2})}$  denotes the Rankin-Cohen bracket (which, as we mentioned above, also works in half-integral weight), and  $\vartheta_{m,\ell}^0(\tau) = \vartheta_{m,\ell}(\tau, 0)$  (Thetanullwerte). There is a similar

formula in the odd case, but with  $\vartheta_{m,\ell}^0(\tau)$  replaced by

$$\vartheta_{m,\ell}^1(\tau) = \frac{1}{2\pi i} \frac{\partial}{\partial z} \vartheta_{m,\ell}(\tau, z) \Big|_{z=0} = \sum_{r \equiv \ell \pmod{2m}} r q^{r^2/4m}. \quad (4.18)$$

### 4.3 Example: Jacobi forms of index 1

If  $m = 1$ , (4.4) reduces to  $c(n, r) = C(4n - r^2)$  where  $C(\Delta)$  is a function of a single argument, because  $4n - r^2$  determines the value of  $r \pmod{2}$ . So any  $\varphi \in J_{k,1}^{\text{weak}}$  has an expansion of the form

$$\varphi(\tau, z) = \sum_{n,r \in \mathbb{Z}} C(4n - r^2) q^n y^r. \quad (4.19)$$

It also follows that  $k$  must be even, since in general,  $C(\Delta, -r) = (-1)^k C(\Delta, r)$ .

One has the isomorphisms  $J_{k,1} \cong M_k \oplus S_{k+2}$  and  $J_{k,1}^{\text{weak}} \cong M_k \oplus M_{k+2}$ . If  $\varphi \in J_{k,1}^{\text{weak}}$  with an expansion as in (4.19), then

$$\varphi(\tau, 0) = \sum_{n=0}^{\infty} a(n) q^n, \quad \frac{1}{2(2\pi i)^2} \varphi''(\tau, 0) = \sum_{n=0}^{\infty} b(n) q^n, \quad (4.20)$$

where

$$a(n) = \sum_{r \in \mathbb{Z}} C(4n - r^2), \quad b(n) = \sum_{r > 0} r^2 C(4n - r^2), \quad (4.21)$$

and the isomorphisms are given (if  $k > 0$ ) by the map  $\varphi \mapsto (\mathcal{A}, \mathcal{B})$  with

$$\mathcal{A}(\tau) = \sum a(n) q^n \in M_k, \quad \mathcal{B}(\tau) = \sum (kb(n) - na(n)) q^n \in M_{k+2}. \quad (4.22)$$

For  $J_{k,1}$  one also has the isomorphism  $J_{k,1} \cong M_{k-\frac{1}{2}}^+(\Gamma_0(4))$  given by

$$\varphi(\tau, z) \leftrightarrow g(\tau) = \sum_{\substack{\Delta \geq 0 \\ \Delta \equiv 0, 3 \pmod{4}}} C(\Delta) q^\Delta. \quad (4.23)$$

We have four particularly interesting examples  $\varphi_{k,1}$

$$\varphi_{k,1}(\tau, z) = \sum_{n,r \in \mathbb{Z}} C_k(4n - r^2) q^n y^r, \quad k = -2, 0, 10, 12, \quad (4.24)$$

which have the properties (defining them uniquely up to multiplication by scalars)

- $\varphi_{10,1}$  and  $\varphi_{12,1}$  are the two index 1 Jacobi cusp forms of smallest weight;
- $\varphi_{-2,1}$  and  $\varphi_{0,1}$  are the unique weak Jacobi forms of index 1 and weight  $\leq 0$ ;

- $\varphi_{-2,1}$  and  $\varphi_{0,1}$  generate the ring of weak Jacobi forms of even weight freely over the ring of modular forms of level 1, so that

$$J_{k,m}^{\text{weak}} = \bigoplus_{j=0}^m M_{k+2j}(SL(2, \mathbb{Z})) \cdot \varphi_{-2,1}^j \varphi_{0,1}^{m-j} \quad (k \text{ even}); \quad (4.25)$$

- $\varphi_{-2,1} = \varphi_{10,1}/\Delta$ ,  $\varphi_{0,1} = \varphi_{12,1}/\Delta$ , and the quotient  $\varphi_{0,1}/\varphi_{-2,1} = \varphi_{12,1}/\varphi_{10,1}$  is a multiple of the Weierstrass  $\wp$  function.

The Fourier coefficients of these functions can be computed from the above recursions, since the pairs  $(\mathcal{A}, \mathcal{B})$  for  $\varphi = \varphi_{-2,1}$ ,  $\varphi_{0,1}$ ,  $\varphi_{10,1}$  and  $\varphi_{12,1}$  are proportional to  $(0, 1)$ ,  $(1, 0)$ ,  $(0, \Delta)$  and  $(\Delta, 0)$ , respectively<sup>7</sup>. The results for the first few Fourier coefficients are given in Table 4.3 below. In particular, the Fourier expansions of  $\varphi_{-2,1}$  and  $\varphi_{0,1}$  begin

$$\varphi_{-2,1} = \frac{(y-1)^2}{y} - 2 \frac{(y-1)^4}{y^2} q + \frac{(y-1)^4(y^2-8y+1)}{y^3} q^2 + \dots, \quad (4.26)$$

$$\varphi_{0,1} = \frac{y^2+10y+1}{y} + 2 \frac{(y-1)^2(5y^2-22y+5)}{y^2} q + \dots. \quad (4.27)$$

$k$	$C_k(-1)$	$C_k(0)$	$C_k(3)$	$C_k(4)$	$C_k(7)$	$C_k(8)$	$C_k(11)$	$C_k(12)$	$C_k(15)$
-2	1	-2	8	-12	39	-56	152	-208	513
0	1	10	-64	108	-513	808	-2752	4016	-11775
10	0	0	1	-2	-16	36	99	-272	-240
12	0	0	1	10	-88	-132	1275	736	-8040

**Table 1:** Some Fourier coefficients

The functions  $\varphi_{k,1}$  ( $k = 10, 0, -2$ ) can be expressed in terms of the Dedekind eta function (3.12) and the Jacobi theta functions  $\vartheta_1, \vartheta_2, \vartheta_3, \vartheta_4$  by the formulas

$$\varphi_{10,1}(\tau, z) = \eta^{18}(\tau) \vartheta_1^2(\tau, z), \quad (4.28)$$

$$\varphi_{-2,1}(\tau, z) = \frac{\vartheta_1^2(\tau, z)}{\eta^6(\tau)} = \frac{\varphi_{10,1}(\tau, z)}{\Delta(\tau)}. \quad (4.29)$$

$$\varphi_{0,1}(\tau, z) = 4 \left( \frac{\vartheta_2(\tau, z)^2}{\vartheta_2(\tau)^2} + \frac{\vartheta_3(\tau, z)^2}{\vartheta_3(\tau)^2} + \frac{\vartheta_4(\tau, z)^2}{\vartheta_4(\tau)^2} \right), \quad (4.30)$$

Finally, we say a few words about Jacobi forms of odd weight. Such a form cannot have index 1, as we saw. In index 2, the isomorphisms  $J_{k,2} \cong S_{k+1}$  and  $J_{k,2}^{\text{weak}} \cong M_{k+1}$  show that the

<sup>7</sup>For  $k = 0$ , the second formula in (4.22) must be modified, and the function  $\sum b(n)q^n$  for  $\varphi_{0,1}$  is in fact  $E_2(\tau)$ .



first examples of holomorphic and weak Jacobi forms occur in weights 11 and  $-1$ , respectively, and are related by  $\varphi_{-1,2} = \varphi_{11,2}/\Delta$ . The function  $\varphi_{-1,2}$  is given explicitly by

$$\varphi_{-1,2}(\tau, z) = \frac{\vartheta_1(\tau, 2z)}{\eta^3(\tau)}, \quad (4.31)$$

with Fourier expansion beginning

$$\varphi_{-1,2} = \frac{y^2 - 1}{y} - \frac{(y^2 - 1)^3}{y^3} q - 3 \frac{(y^2 - 1)^3}{y^3} q^2 + \dots, \quad (4.32)$$

and its square is related to the index 1 Jacobi forms defined above by

$$432 \varphi_{-1,2}^2 = \varphi_{-2,1} (\varphi_{0,1}^3 - 3 E_4 \varphi_{-2,1} \varphi_{0,1} + 2 E_6 \varphi_{-2,1}^3). \quad (4.33)$$

(In fact,  $\varphi_{-1,2}/\varphi_{-2,1}^2$  is a multiple of  $\wp'(\tau, z)$  and this equation, divided by  $\varphi_{-2,1}^4$ , is just the usual equation expressing  $\wp'^2$  as a cubic polynomial in  $\wp$ .) It is convenient to introduce abbreviations

$$A = \varphi_{-2,1}, \quad B = \varphi_{0,1}, \quad C = \varphi_{-1,2}. \quad (4.34)$$

With these notations, the structure of the full bigraded ring of weak Jacobi forms is given by

$$J_{*,*}^{\text{weak}} = \mathbb{C}[E_4, E_6, A, B, C] / (432C^2 - AB^3 + 3E_4A^3B - 2E_6A^4). \quad (4.35)$$

#### 4.4 Hecke-like operators

In [52] Hecke operators  $T_\ell$  acting on  $J_{k,m}$  were introduced, but also various ‘‘Hecke-like’’ operators, again defined by the action of certain combinations of elements of  $GL(2, \mathbb{Q}) \times \mathbb{Q}^2$ , which send Jacobi forms to Jacobi forms, but now possibly changing the index. We describe these operators here. This subsection is more technical than the preceding three, but will be essential later.

The first is the very simple operator  $U_s$  ( $s \geq 1$ ) which sends  $\varphi(\tau, z)$  to  $\varphi(\tau, sz)$ , i.e.,

$$U_s : \sum_{n,r} c(n, r) q^n y^r \mapsto \sum_{n,r} c(n, r) q^n y^{sr}, \quad (4.36)$$

This operator maps  $J_{k,m}$  to  $J_{k,s^2m}$ .

The second operator,  $V_{k,t}$  ( $t \geq 1$ ), sends  $J_{k,m}$  to  $J_{k,tm}$ . It is given in terms of its action on Fourier coefficients by

$$c(\varphi|V_{k,t}; n, r) = \sum_{d|(n,r,t)} d^{k-1} c\left(\varphi; \frac{nt}{d^2}, \frac{r}{d}\right), \quad C(\varphi|V_{k,t}; \Delta, r) = \sum_{d|(\frac{t^2+\Delta}{4mt}, r, t)} d^{k-1} C\left(\varphi; \frac{\Delta}{d^2}, \frac{r}{d}\right). \quad (4.37)$$

In §9, we will also need the modified operator  $\mathcal{V}_{k,t}^{(m)} : J_{k,m} \mapsto J_{k,tm}$  defined by

$$\mathcal{V}_{k,t}^{(m)} = \sum_{\substack{s^2|t \\ (s,m)=1}} \mu(s) s^{k-1} V_{k,t/s^2} U_s, \quad (4.38)$$

so that, for instance,  $\mathcal{V}_{k,t}^{(m)} = V_{k,t}$  if  $t$  is square-free, but  $\mathcal{V}_{k,4}^{(m)} = V_{k,4} - 2^{k-1}U_2$  if  $m$  is odd.

Next, for each positive integer  $m_1$  with  $m_1 \parallel m$  (this means that  $m = m_1 m_2$  with  $m_1$  and  $m_2$  coprime) we have an involution  $W_{m_1}$  on  $J_{k,m}$ , which we will call an *Atkin-Lehner involution*,<sup>8</sup> that is defined in terms of the theta expansion of Jacobi forms by

$$W_{m_1} : \sum_{\ell \pmod{2m}} h_\ell(\tau) \vartheta_{m,\ell}(\tau, z) \mapsto \sum_{\ell \pmod{2m}} h_{\ell^*}(\tau) \vartheta_{m,\ell}(\tau, z) \quad (4.39)$$

(or equivalently by  $C(\Delta, r) \mapsto C(\Delta, r^*)$ ), where the involution  $\ell \mapsto \ell^*$  on  $\mathbb{Z}/2m\mathbb{Z}$  is defined by

$$\ell^* \equiv -\ell \pmod{2m_1}, \quad \ell^* \equiv +\ell \pmod{2m_2}. \quad (4.40)$$

These operators commute and satisfy  $W_{m/m_1} = (-1)^k W_{m_1}$ , so that we get an eigenspace decomposition

$$J_{k,m} = \bigoplus_{\substack{\boldsymbol{\varepsilon}=(\varepsilon_1, \dots, \varepsilon_t) \in \{\pm 1\}^t \\ \varepsilon_1 \cdots \varepsilon_t = (-1)^k}} J_{k,m}^{\boldsymbol{\varepsilon}}, \quad (4.41)$$

where  $m = p_1^{r_1} \cdots p_t^{r_t}$  is the prime power decomposition of  $m$  and  $\varepsilon_i$  is the eigenvalue of  $W_{p_i^{r_i}}$ . One can also introduce the “ $p$ -adic Atkin-Lehner involution”  $W_{p^\infty}$ , which acts on any  $J_{k,m}$  as  $W_{p^\nu}$  where  $p^\nu \parallel m$ . (In particular, it acts as the identity if  $p \nmid m$ .) This operator simply changes the second index  $r$  in  $C(\Delta, r)$  to  $r^*$ , where  $r$  is equal to the negative of  $r^*$   $p$ -adically and to  $r^*$   $p'$ -adically for any  $p' \neq p$ . In particular,  $W_{p^\infty}$  commutes with all  $U_s$ ,  $V_{k,t}$ , and  $\mathcal{V}_{k,t}^m$ .

We now describe a further Hecke-like operator, which was not given in [52] but is mentioned briefly in [104], p. 139, and which we will also need in the sequel. If  $t$  is a positive integer whose square divides  $m$ , then we define  $u_t : J_{k,m} \rightarrow J_{k,m/t^2}$  by

$$u_t : \sum_{n,r} c(n,r) q^n y^r \mapsto \sum_{n,r} \left( \sum_{a \pmod{t}} c\left(n + ar + \frac{m}{t^2} a^2, tr + \frac{2ma}{t}\right) \right) q^n y^r \quad (4.42)$$

or equivalently by

$$C_{\varphi|u_t}(\Delta, \ell \pmod{2m/t^2}) = \sum_{\substack{r \pmod{2m/t} \\ r \equiv \ell \pmod{2m/t^2}}} C_\varphi(\Delta t^2, rt \pmod{2m}). \quad (4.43)$$

---

<sup>8</sup>This terminology was not used in [52], where these involutions were introduced, but is justified by the results of [104] where it is shown that they correspond in a precise sense to the Atkin-Lehner involutions on the space of modular forms of weight  $2k - 2$  and level  $m$ .

It is easily checked that this map does indeed send Jacobi forms to Jacobi forms of the index stated, and that for any integers  $m, t > 0$ , the composite map  $J_{k,m/t^2} \xrightarrow{U_t} J_{k,m} \xrightarrow{u_t} J_{k,m/t^2}$  is simply multiplication by  $t$  (because, if  $\varphi = \varphi_1 | U_t$  with  $\varphi_1$  of index  $m/t^2$ , then each term on the right of (4.43) equals  $C_{\varphi_1}(\Delta, \ell)$ ). It follows that we have

$$J_{k,m} = \bigoplus_{t^2|m} J_{k, \frac{m}{t^2}}^{\text{prim}} | U_t, \quad (4.44)$$

where

$$J_{k,m}^{\text{prim}} = \bigcap_{\substack{t^2|m \\ t>1}} \ker (J_{k,m} \xrightarrow{u_t} J_{k,m/t^2}) = \bigcap_{\substack{p^2|m \\ p \text{ prime}}} \ker (J_{k,m} \xrightarrow{u_p} J_{k,m/p^2}) \quad (4.45)$$

(the equivalence of the two definitions follows because  $u_t u_{t'} = u_{tt'}$  for any  $t, t'$ ), with the projection map  $\pi^{\text{prim}} : J_{k,m} \mapsto J_{k,m}^{\text{prim}}$  onto the first factor in (4.44) given by

$$\pi^{\text{prim}}(\varphi) = \sum_{t^2|m} \frac{\mu(t)}{t} \varphi | u_t | U_t. \quad (4.46)$$

We will sometimes find it useful to work, not with the space of Jacobi forms, but with the larger space of holomorphic functions on  $\mathbb{H} \times \mathbb{C}$  that obey the elliptic transformation law (4.2) of Jacobi forms, but are not required to obey the modular transformation property (4.1). We shall call such functions *elliptic forms*. (The word *Jacobi-like forms* has been used in the literature to denote functions which, conversely, satisfy (4.1) but not (4.2).) Any elliptic form has a Fourier expansion as in (4.3) with the periodicity condition (4.4). We will denote by  $\mathcal{E}_m, \mathcal{E}_m^0, \tilde{\mathcal{E}}_m$  the spaces of all elliptic forms of index  $m$  satisfying the growth conditions (4.5), (4.6), and (4.7), respectively, and we will call them *holomorphic*, *cuspidal*, and *weak* respectively. We also denote by  $\mathcal{E}_{\pm,m}, \mathcal{E}_{\pm,m}^0, \tilde{\mathcal{E}}_{\pm,m}$  the  $(\pm 1)$ -eigenspaces of theta functions under the involution  $\varphi(\tau, z) \mapsto \varphi(\tau, -z)$ , i.e., those functions whose Fourier coefficients  $c(n, r)$  satisfy the parity condition  $c(n, -r) = \pm c(n, r)$ . Because any Jacobi form of weight  $k$  satisfies this parity condition with  $\pm 1 = (-1)^k$  (by (4.1) applied to  $-\mathbf{1}_2$ ), we have

$$J_{k,m} \subset \mathcal{E}_{\pm,m}, \quad J_{k,m}^0 \subset \mathcal{E}_{\pm,m}^0, \quad \tilde{J}_{k,m} \subset \tilde{\mathcal{E}}_{\pm,m} \quad ((-1)^k = \pm 1). \quad (4.47)$$

It is now clear from the definitions above that all of the Hecke-like operators  $V_t, U_t, W_{m_1}$  for  $(m_1, m/m_1) = 1$  and  $u_t$  for  $t^2|m$  defined in this section extend from the space of Jacobi forms to the space of elliptic forms, and that the decompositions (4.41) and (4.44) remain true if we replace all the  $J_k$ s by  $\mathcal{E}_+$ s.

Finally, we should warn the reader explicitly that, although the operators  $U_t$  and  $V_t$  also act on the spaces  $\tilde{J}_{k,*}$  and  $\tilde{\mathcal{E}}_{\pm,*}$  of weak Jacobi or weak elliptic forms, the operators  $W_{m_1}$  and  $u_t$  do not: the condition (4.7) is equivalent to saying that  $C(\Delta, \ell)$  for  $|\ell| \leq m$  vanishes whenever  $\Delta <$

$-\ell^2$ , and this condition is preserved neither under the permutation  $\ell \pmod{2m} \mapsto \ell^* \pmod{2m}$  defining  $W_{m_1}$ , nor by the summation conditions in (4.43). As a concrete example, the 7-dimensional space  $\tilde{\mathcal{J}}_{2,6}$  contains a 6-dimensional subspace of forms invariant under the involution  $W_2$ , but no anti-invariant form, and its image under the projection operator  $\pi_2^- = \frac{1}{2}(1 - W_2)$  to anti-invariant forms is the 1-dimensional space spanned by

$$\varphi_{2,6}^-(\tau, z) = j'(\tau) \frac{\vartheta_1(\tau, 4z)}{\vartheta_1(\tau, 2z)} = \sum_{n,r \in \mathbb{Z}} \chi_{12}(r) C_{2,6}(24n - r^2) q^n y^r, \quad (4.48)$$

with

$\Delta$	$\leq -49$	$-25$	$-1$	$23$	$47$	$71$	$\dots$
$C_{2,6}(\Delta)$	$0$	$-1$	$-1$	$196882$	$43184401$	$2636281193$	$\dots$

which is a weakly holomorphic but not a weak Jacobi form. (Specifically, for any  $\varphi \in \tilde{\mathcal{J}}_{2,6}$  we have  $\pi_2^-(\varphi) = 6 c_\varphi(0, 5) \varphi_{2,6}^-$ .)

#### 4.5 Quantum black holes and Jacobi forms

Jacobi forms usually arise in string theory as elliptic genera of two-dimensional superconformal field theories (SCFT) with (2, 2) or more worldsheet supersymmetry<sup>9</sup>. We denote the superconformal field theory by  $\sigma(\mathcal{M})$  when it corresponds to a sigma model with a target manifold  $\mathcal{M}$ . Let  $H$  be the Hamiltonian in the Ramond sector, and  $J$  be the left-moving  $U(1)$   $R$ -charge. The elliptic genus  $\chi(\tau, z; \mathcal{M})$  is then defined as [112, 3, 87] a trace over the Hilbert space  $\mathcal{H}_R$  in the Ramond sector

$$\chi(\tau, z; \mathcal{M}) = \text{Tr}_{\mathcal{H}_R}((-1)^F q^H y^J), \quad (4.49)$$

where  $F$  is the fermion number.

An elliptic genus so defined satisfies the modular transformation property (4.1) as a consequence of modular invariance of the path integral. Similarly, it satisfies the elliptic transformation property (4.2) as a consequence of spectral flow. Furthermore, in a unitary SCFT, the positivity of the Hamiltonian implies that the elliptic genus is a weak Jacobi form. The decomposition (4.10) follows from bosonizing the  $U(1)$  current in the standard way so that the contribution to the trace from the momentum modes of the boson can be separated into the theta function (4.11). See, for example, [69, 85] for a discussion. This notion of the elliptic genus can be generalized to a (0, 2) theory using a left-moving  $U(1)$  charge  $J$  which may not be an  $R$ -charge. In this case spectral flow is imposed as an additional constraint and follows from gauge invariance under large gauge transformations [39, 56, 72, 43].

A particularly useful example in the present context is  $\sigma(K3)$ , which is a (4, 4) SCFT whose target space is a  $K3$  surface. The elliptic genus is a topological invariant and is independent of

<sup>9</sup>An SCFT with  $(r, s)$  supersymmetries has  $r$  left-moving and  $s$  right-moving supersymmetries.

the moduli of the  $K3$ . Hence, it can be computed at some convenient point in the  $K3$  moduli space, for example, at the orbifold point where the  $K3$  is the Kummer surface. At this point, the  $\sigma(K3)$  SCFT can be regarded as a  $\mathbb{Z}_2$  orbifold of the  $\sigma(T^4)$  SCFT, which is an SCFT with a torus  $T^4$  as the target space. A simple computation using standard techniques of orbifold conformal field theory [38, 58] yields

$$\chi(\tau, z; K3) = 2\varphi_{0,1}(\tau, z) = 2 \sum C_0(4n - l^2) q^n y^l. \quad (4.50)$$

Note that for  $z = 0$ , the trace (4.49) reduces to the Witten index of the SCFT and correspondingly the elliptic genus reduces to the Euler character of the target space manifold. In our case, one can readily verify from (4.50) and (4.30) that  $\chi(\tau, 0; K3)$  equals 24, which is the Euler character of  $K3$ .

A well-known physical application of Jacobi forms is in the context of the five-dimensional Strominger-Vafa black hole[107], which is a bound state of  $Q_1$  D1-branes,  $Q_5$  D5-branes,  $n$  units of momentum and  $l$  units of five-dimensional angular momentum [12]. The degeneracies  $d_m(n, l)$  of such black holes depend only on  $m = Q_1 Q_5$ . They are given by the Fourier coefficients  $c(n, l)$  of the elliptic genus  $\chi(\tau, z; \text{Sym}^{m+1}(K3))$  of symmetric product of  $(m+1)$  copies of  $K3$ -surface.

Let us denote the generating function for the elliptic genera of symmetric products of  $K3$  by

$$\widehat{Z}(\sigma, \tau, z) := \sum_{m=-1}^{\infty} \chi(\tau, z; \text{Sym}^{m+1}(K3)) p^m \quad (4.51)$$

where  $\chi_m(\tau, z)$  is the elliptic genus of  $\text{Sym}^m(K3)$ . A standard orbifold computation [45] gives

$$\widehat{Z}(\sigma, \tau, z) = \frac{1}{p} \prod_{r>0, s\geq 0, t} \frac{1}{(1 - q^s y^t p^r)^{2C_0(rs, t)}} \quad (4.52)$$

in terms of the Fourier coefficients  $2C_o$  of the elliptic genus of a single copy of  $K3$ .

For  $z = 0$ , it can be checked that, as expected, the generating function (4.52) for elliptic genera reduces to the generating function (3.26) for Euler characters

$$\widehat{Z}(\sigma, \tau, 0) = \widehat{Z}(\sigma) = \frac{1}{\Delta(\sigma)}. \quad (4.53)$$

## 5. Siegel modular forms

### 5.1 Definitions and examples of Siegel modular forms

Let  $Sp(2, \mathbb{Z})$  be the group of  $(4 \times 4)$  matrices  $g$  with integer entries satisfying  $gJg^t = J$  where

$$J \equiv \begin{pmatrix} 0 & -I_2 \\ I_2 & 0 \end{pmatrix} \quad (5.1)$$

is the symplectic form. We can write the element  $g$  in block form as

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}, \quad (5.2)$$

where  $A, B, C, D$  are all  $(2 \times 2)$  matrices with integer entries. Then the condition  $gJg^t = J$  implies

$$AB^t = BA^t, \quad CD^t = DC^t, \quad AD^t - BC^t = \mathbf{1}, \quad (5.3)$$

Let  $\mathbb{H}_2$  be the (genus two) Siegel upper half plane, defined as the set of  $(2 \times 2)$  symmetric matrix  $\Omega$  with complex entries

$$\Omega = \begin{pmatrix} \tau & z \\ z & \sigma \end{pmatrix} \quad (5.4)$$

satisfying

$$\operatorname{Im}(\tau) > 0, \quad \operatorname{Im}(\sigma) > 0, \quad \det(\operatorname{Im}(\Omega)) > 0. \quad (5.5)$$

An element  $g \in Sp(2, \mathbb{Z})$  of the form (5.2) has a natural action on  $\mathbb{H}_2$  under which it is stable:

$$\Omega \rightarrow (A\Omega + B)(C\Omega + D)^{-1}. \quad (5.6)$$

The matrix  $\Omega$  can be thought of as the period matrix of a genus two Riemann surface<sup>10</sup> on which there is a natural symplectic action of  $Sp(2, \mathbb{Z})$ .

A Siegel form  $F(\Omega)$  of weight  $k$  is a holomorphic function  $\mathbb{H}_2 \rightarrow \mathbb{C}$  satisfying

$$F((A\Omega + B)(C\Omega + D)^{-1}) = \det(C\Omega + D)^k F(\Omega). \quad (5.7)$$

A Siegel modular form can be written in terms of its Fourier series

$$F(\Omega) = \sum_{\substack{n, r, m \in \mathbb{Z} \\ r^2 \leq 4mn}} a(n, r, m) q^n y^r p^m. \quad (5.8)$$

If one writes this as

$$F(\Omega) = \sum_{m=0}^{\infty} \varphi_m^F(\tau, z) p^m \quad (5.9)$$

with

$$\varphi_m^F(\tau, z) = \sum_{n, r} a(n, r, m) q^n y^r, \quad (5.10)$$

then each  $\varphi_m^F(m \geq 0)$  is a Jacobi form of weight  $k$  and index  $m$ .

---

<sup>10</sup>See [54, 36, 9] for a discussion of the connection with genus-two Riemann surfaces.

An important special class of Siegel forms were studied by Maass which he called the *Spezielschar*. They have the property that  $a(n, r, m)$  depends only on the discriminant  $4mn - r^2$  if  $(n, r, m)$  are coprime, and more generally

$$a(n, r, m) = \sum_{d|(n,r,m), d>0} d^{k-1} C\left(\frac{4mn - r^2}{d^2}\right) \quad (5.11)$$

for some coefficients  $C(N)$ . Specializing to  $m = 1$ , we can see that these numbers are simply the coefficients associated via (4.19) to the Jacobi form  $\varphi = \varphi_1^F \in J_{k,1}$ , and that (5.11) says precisely that the other Fourier-Jacobi coefficients of  $F$  are given by  $\varphi_m^F = \varphi_1^F|V_m$  with  $V_m$  as in (4.37). Conversely, if  $\varphi$  is any Jacobi form of weight  $k$  and index 1 with Fourier expansion (4.19), then the function  $F(\Omega)$  defined by (5.8) and (5.11) or by  $F(\Omega) = \sum_{m=0}^{\infty} (\varphi|V_m)(\tau, z) p^m$  is a Siegel modular form of weight  $k$  with  $\varphi_1^F = \varphi$ . The resulting map from  $J_{k,1}$  to the *Spezielschar* is called the Saito-Kurokawa lift or *additive lift* since it naturally gives the sum representation of a Siegel form using the Fourier coefficients of a Jacobi form as the input. (More information about the additive lift can be found in [52].)

The example of interest to us is the Igusa cusp form  $\Phi_{10}$  (the unique cusp form of weight 10) which is the Saito-Kurokawa lift of the Jacobi form  $\varphi_{10,1}$  introduced earlier, so that

$$\Phi_{10}(\Omega) = \sum_{m=1}^{\infty} (\varphi_{10,1}|V_m)(\tau, z) p^m = \sum_{n,r,m} a_{10}(n, r, m) q^n y^r p^m, \quad (5.12)$$

where  $a_{10}$  is defined by (5.11) with  $k = 10$  in terms of the coefficients  $C_{10}(d)$  given in Table 4.3.

A Siegel modular form sometimes also admits a product representation, and can be obtained as Borcherds lift or *multiplicative lift* of a weak Jacobi form of weight zero and index one. This procedure is in a sense an exponentiation of the additive lift and naturally results in the product representation of the Siegel form using the Fourier coefficients of a Jacobi form as the input. Several examples of Siegel forms that admit product representation are known but at present there is no general theory to determine under what conditions a given Siegel form admits a product representation.

For the Igusa cusp form  $\Phi_{10}$ , a product representation does exist. It was obtained by Gritsenko and Nikulin [61, 62] as a multiplicative lift of the elliptic genus  $\chi(\tau, z; K3) = 2\varphi_{0,1}(\tau, z)$  and is given by

$$\Phi_{10}(\Omega) = qyp \prod_{(r,s,t)>0} (1 - q^s y^t p^r)^{2C_0(4rs-t^2)}, \quad (5.13)$$

in terms of  $C_0$  given by (4.30, 4.24). Here the notation  $(r, s, t) > 0$  means that  $r, s, t \in \mathbb{Z}$  with either  $r > 0$  or  $r = 0, s > 0$ , or  $r = s = 0, t < 0$ . In terms of the  $V_m$ , this can be rewritten in the form  $\Phi_{10} = p \varphi_{10,1} \exp(-2 \sum_{m=1}^{\infty} (\varphi_{0,1}|V_m) p^m)$ .



## 5.2 Quantum black holes and Siegel modular forms

Siegel forms occur naturally in the context of counting of quarter-BPS dyons. The partition function for these dyons depends on three (complexified) chemical potentials  $(\sigma, \tau, z)$ , conjugate to the three  $T$ -duality invariant integers  $(m, n, \ell)$  respectively and is given by

$$Z(\Omega) = \frac{1}{\Phi_{10}(\Omega)}. \quad (5.14)$$

The product representation of the Igusa form is particularly useful for the physics application because it is closely related to the generating function for the elliptic genera of symmetric products of  $K3$  introduced earlier. This is a consequence of the fact that the multiplicative lift of the Igusa form is obtained starting with the elliptic genus of  $K3$  as the input. Comparing the product representation for the Igusa form (5.13) with (4.52), we get the relation:

$$Z(\tau, z, \sigma) = \frac{1}{\Phi_{10}(\tau, z, \sigma)} = \frac{\widehat{Z}(\tau, z, \sigma)}{\varphi_{10,1}(\tau, z)}. \quad (5.15)$$

This relation to the elliptic genera of symmetric products of  $K3$  has a deeper physical significance based on what is known as the 4d-5d lift [55]. The main idea is to use the fact that the geometry of the Kaluza-Klein monopole in the charge configuration (2.15) reduces to five-dimensional flat Minkowski spacetime in the limit when the radius of the circle  $\tilde{S}^1$  goes to infinity. In this limit, the charge  $l$  corresponding to the momentum around this circle gets identified with the angular momentum  $l$  in five dimensions. Our charge configuration (2.15) then reduces essentially to the Strominger-Vafa black hole [107] with angular momentum [12] discussed in the previous subsection. Assuming that the dyon partition function does not depend on the moduli, we thus essentially relate  $Z(\Omega)$  to  $\widehat{Z}(\Omega)$ . The additional factor in (5.15) involving  $\Phi_{10}(\sigma, \tau, z)$  comes from bound states of momentum  $n$  with the Kaluza-Klein monopole and from the center of mass motion of the Strominger-Vafa black hole in the Kaluza-Klein geometry [54, 37].

The Igusa cusp form has double zeros at  $z = 0$  and its  $Sp(2, \mathbb{Z})$  images. The partition function is therefore a *meromorphic* Siegel form (5.7) of weight  $-10$  with double poles at these divisors. This fact is responsible for much of the interesting physics of wall-crossings in this context as we explain in the next section.

Using (5.12) or (5.13), one can compute the coefficients  $\psi_m$  in the Fourier-Jacobi expan-

sion (1.1). The first few, multiplied by  $\Delta$  and a suitable integer, are given by

$$\begin{aligned}
\Delta \psi_{-1} &= A^{-1}, \\
\Delta \psi_0 &= 2 A^{-1} B, \\
4 \Delta \psi_1 &= 9 A^{-1} B^2 + 3 E_4 A, \\
27 \Delta \psi_2 &= 50 A^{-1} B^3 + 48 E_4 A B + 10 E_6 A^2, \\
384 \Delta \psi_3 &= 475 A^{-1} B^4 + 886 E_4 A B^2 + 360 E_6 A^2 B + 199 E_4^2 A^3, \\
72 \Delta \psi_4 &= 51 A^{-1} B^5 + 155 E_4 A B^3 + 93 E_6 A^2 B^2 + 102 E_4^2 A^3 B + 31 E_4 E_6 A^4.
\end{aligned} \tag{5.16}$$

(Here  $E_4, E_6, \Delta$  are as in §3 and  $A = \varphi_{-2,1}, B = \varphi_{0,1}$  as in §4.) The double zero of  $\Phi_{10}$  at  $z = 0$  is reflected by the denominator  $A$  in the  $A^{-1} B^{m+1}$  terms in these formulas. Note that, by (5.15), the right-hand side of (5.16) multiplied by  $A$  and divided by a suitable denominator are equal to the elliptic genus of symmetric products of a  $K3$  surface. For example,

$$\chi(\text{Sym}^k K3; \tau, z) = \frac{1}{384} (475 B^4 + 886 E_4 A^2 B^2 + 360 E_6 A^3 B + 199 E_4^2 A^4). \tag{5.17}$$

## 6. Walls and contours

Given the partition function (5.14), one can extract the black hole degeneracies from the Fourier coefficients. The three quadratic  $T$ -duality invariants of a given dyonic state can be organized as a  $2 \times 2$  symmetric matrix

$$\Lambda = \begin{pmatrix} N \cdot N & N \cdot M \\ M \cdot N & M \cdot M \end{pmatrix} = \begin{pmatrix} 2n & \ell \\ \ell & 2m \end{pmatrix}, \tag{6.1}$$

where the dot products are defined using the  $O(22, 6; \mathbb{Z})$  invariant metric  $L$ . The matrix  $\Omega$  in (5.14) and (5.4) can be viewed as the matrix of complex chemical potentials conjugate to the charge matrix  $\Lambda$ . The charge matrix  $\Lambda$  is manifestly  $T$ -duality invariant. Under an  $S$ -duality transformation (2.4), it transforms as

$$\Lambda \rightarrow \gamma \Lambda \gamma^t \tag{6.2}$$

There is a natural embedding of this physical  $S$ -duality group  $SL(2, \mathbb{Z})$  into  $Sp(2, \mathbb{Z})$ :

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} (\gamma^t)^{-1} & \mathbf{0} \\ \mathbf{0} & \gamma \end{pmatrix} = \begin{pmatrix} d & -c & 0 & 0 \\ -b & a & 0 & 0 \\ 0 & 0 & a & b \\ 0 & 0 & c & d \end{pmatrix} \in Sp(2, \mathbb{Z}). \tag{6.3}$$

The embedding is chosen so that  $\Omega \rightarrow (\gamma^T)^{-1}\Omega\gamma^{-1}$  and  $\text{Tr}(\Omega \cdot \Lambda)$  in the Fourier integral is invariant. This choice of the embedding ensures that the physical degeneracies extracted from the Fourier integral are  $S$ -duality invariant if we appropriately transform the moduli at the same time as we explain below.

To specify the contours, it is useful to define the following moduli-dependent quantities. One can define the matrix of right-moving  $T$ -duality invariants

$$\Lambda_R = \begin{pmatrix} Q_R \cdot Q_R & Q_R \cdot P_R \\ P_R \cdot Q_R & P_R \cdot P_R \end{pmatrix}, \quad (6.4)$$

which depends both on the integral charge vectors  $N, M$  as well as the  $T$ -moduli  $\mu$ . One can then define two matrices naturally associated to the  $S$ -moduli  $S = S_1 + iS_2$  and the  $T$ -moduli  $\mu$  respectively by

$$\mathcal{S} = \frac{1}{S_2} \begin{pmatrix} |S|^2 & S_1 \\ S_1 & 1 \end{pmatrix}, \quad \mathcal{T} = \frac{\Lambda_R}{|\det(\Lambda_R)|^{\frac{1}{2}}}. \quad (6.5)$$

Both matrices are normalized to have unit determinant. In terms of them, we can construct the moduli-dependent ‘central charge matrix’

$$\mathcal{Z} = |\det(\Lambda_R)|^{\frac{1}{4}} (\mathcal{S} + \mathcal{T}), \quad (6.6)$$

whose determinant equals the BPS mass

$$M_{Q,P} = |\det \mathcal{Z}|. \quad (6.7)$$

We define

$$\tilde{\Omega} \equiv \begin{pmatrix} \sigma & -z \\ -z & \tau \end{pmatrix}. \quad (6.8)$$

This is related to  $\Omega$  by an  $SL(2, \mathbb{Z})$  transformation

$$\tilde{\Omega} = \hat{S}\Omega\hat{S}^{-1} \quad \text{where} \quad \hat{S} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (6.9)$$

so that, under a general  $S$ -duality transformation  $\gamma$ , we have the transformation  $\tilde{\Omega} \rightarrow \gamma\tilde{\Omega}\gamma^T$  as  $\Omega \rightarrow (\gamma^T)^{-1}\Omega\gamma^{-1}$ .

With these definitions,  $\Lambda, \Lambda_R, \mathcal{Z}$  and  $\tilde{\Omega}$  all transform as  $X \rightarrow \gamma X \gamma^T$  under an  $S$ -duality transformation (2.4) and are invariant under  $T$ -duality transformations. The moduli-dependent Fourier contour can then be specified in a duality-invariant fashion by [22]

$$\mathcal{C} = \{\text{Im}(\tilde{\Omega}) = \varepsilon^{-1}\mathcal{Z}; \quad 0 \leq \text{Re}(\tau), \text{Re}(\sigma), \text{Re}(z) < 1\}, \quad (6.10)$$

where  $\varepsilon \rightarrow 0^+$ . For a given set of charges, the contour depends on the moduli  $S, \mu$  through the definition of the central charge vector (6.6). The degeneracies  $d(n, \ell, m)|_{S, \mu}$  of states with the  $T$ -duality invariants  $(n, \ell, m)$  at a given point  $(S, \mu)$  in the moduli space are then given by<sup>11</sup>

$$d(n, \ell, m)|_{S, \mu} = \int_{\mathcal{C}} e^{-i\pi \text{Tr}(\Omega \cdot \Lambda)} \mathcal{Z}(\Omega) d^3\Omega. \quad (6.11)$$

This contour prescription thus specifies how to extract the degeneracies from the partition function for a given set of charges and in any given region of the moduli space. In particular, it also completely summarizes all wall-crossings as one moves around in the moduli space for a fixed set of charges. Even though the indexed partition function has the same functional form throughout the moduli space, the spectrum is moduli dependent because of the moduli dependence of the contours of Fourier integration and the pole structure of the partition function. Since the degeneracies depend on the moduli *only* through the dependence of the contour  $\mathcal{C}$ , moving around in the moduli space corresponds to deforming the Fourier contour. This does not change the degeneracy except when one encounters a pole of the partition function. Crossing a pole corresponds to crossing a wall in the moduli space. The moduli space is thus divided up into domains separated by “walls of marginal stability.” In each domain the degeneracy is constant but it jumps upon crossing a wall as one goes from one domain to the other. The jump in the degeneracy has a nice mathematical characterization. It is simply given by the residue at the pole that is crossed while deforming the Fourier contour in going from one domain to the other.

We now turn to the degeneracies of single-centered black holes. Given the  $T$ -duality invariants  $\Lambda$ , a single centered black hole solution is known to exist in all regions of the moduli space as long as  $\det(\Lambda)$  is large and positive. The moduli fields can take any values  $(\lambda_\infty, \mu_\infty)$  at asymptotic infinity far away from the black hole but they vary in the black hole geometry. Because of the attractor phenomenon [53, 105], the moduli adjust themselves so that near the horizon of the black hole of charge  $\Lambda$  they get attracted to the values  $(\lambda_*(\Lambda), \mu_*(\Lambda))$  which are determined by the requirement that the central charge  $\mathcal{Z}_*(\Lambda)$  evaluated using these moduli becomes proportional to  $\Lambda$ . These attractor values are independent of the asymptotic values and depend only on the charge of black hole. We call these moduli the attractor moduli. This enables us to define the *attractor contour* for a given charge  $\Lambda$  by fixing the asymptotic moduli to the attractor values corresponding to this charge. In this case

$$\mathcal{Z}(\lambda_\infty, \mu_\infty) = \mathcal{Z}(\lambda_*(\Lambda), \mu_*(\Lambda)) \sim \Lambda \quad (6.12)$$

and we have the attractor contour

$$\mathcal{C}_* = \{\text{Im}(\tilde{\Omega}) = \varepsilon^{-1}\Lambda; \quad 0 \leq \text{Re}(\tau), \text{Re}(\sigma), \text{Re}(z) < 1\} \quad (6.13)$$

---

<sup>11</sup>The physical degeneracies have an additional multiplicative factor of  $(-1)^{\ell+1}$  which we omit here for simplicity of notation in later chapters.

which depends only on the integral charges and not on the moduli. The significance of the attractor moduli in our context stems from the fact if the asymptotic moduli are tuned to these values for given  $(n, \ell, m)$ , then only single-centered black hole solution exists. The degeneracies  $d^*(n, \ell, m)$  obtained using the attractor contour

$$d^*(n, \ell, m) = \int_{\mathcal{C}_*} e^{-i\pi \text{Tr}(\Omega \cdot \Lambda)} Z(\Omega) d^3\Omega \quad (6.14)$$

are therefore expected to be the degeneracies of the immortal single-centered black holes.

## 7. Mock modular forms

Mock modular forms are a relatively new class of modular objects, although individual examples had been known for some time. They were first isolated explicitly by S. Zwegers in his thesis [119] as the explanation of the “mock theta functions” introduced by Ramanujan in his famous last letter to Hardy. An expository account of this work can be found in [118].

Ramanujan’s mock theta functions are examples of what we will call “pure” mock modular forms. By this we mean a holomorphic function in the upper half plane which transforms under modular transformations almost, but not quite, as a modular form. The non-modularity is of a very special nature and is governed by another holomorphic function called the *shadow* which is itself an ordinary modular form. We will describe this more precisely in §7.1 and give a number of examples. In §7.2, we introduce a notion of mock Jacobi forms (essentially, holomorphic functions of  $\tau$  and  $z$  with theta expansions like that of usual Jacobi forms, but in which the coefficients  $h_\ell(\tau)$  are mock modular forms) and show how all the examples given in the first subsection occur naturally as pieces of mock Jacobi forms. Finally in §7.3 we define a more general notion of mock modular forms in which the shadow is replaced by a sum of products of holomorphic and anti-holomorphic modular forms. This has the advantage that there are now many examples of strongly holomorphic mock modular forms, whereas pure mock modular forms almost always have a negative power of  $q$  at at least one cusp, and also that almost all of the examples occurring naturally in number theory, algebraic geometry, or physics are of this more general “mixed” type, with the pure mock modular forms arising as quotients of mixed mock modular forms by eta products.

### 7.1 Pure mock modular forms

We define a (weakly holomorphic) *pure mock modular form* of weight  $k \in \frac{1}{2}\mathbb{Z}$  as the first member of a pair  $(h, g)$ , where

1.  $h$  is a holomorphic function in  $\mathbb{H}$  with at most exponential growth at all cusps,

2. the function  $g(\tau)$ , called the *shadow* of  $h$ , is a holomorphic<sup>12</sup> modular form of weight  $2 - k$ , and
3. the sum  $\widehat{h} := h + g^*$ , called the *completion* of  $h$ , transforms like a holomorphic modular form of weight  $k$ , i.e.  $\widehat{h}(\tau)/\theta(\tau)^{2k}$  is invariant under  $\tau \rightarrow \gamma\tau$  for all  $\tau \in \mathbb{H}$  and for all  $\gamma$  in some congruence subgroup of  $SL(2, \mathbb{Z})$ .

Here  $g^*(\tau)$ , called the *non-holomorphic Eichler integral*, is a solution of the differential equation

$$(4\pi\tau_2)^k \frac{\partial g^*(\tau)}{\partial \bar{\tau}} = -2\pi i \overline{g(\tau)}. \quad (7.1)$$

If  $g$  has the Fourier expansion  $g(\tau) = \sum_{n \geq 0} b_n q^n$ , we fix the choice of  $g^*$  by setting

$$g^*(\tau) = \bar{b}_0 \frac{(4\pi\tau_2)^{-k+1}}{k-1} + \sum_{n>0} n^{k-1} \bar{b}_n \Gamma(1-k, 4\pi n\tau_2) q^{-n}, \quad (7.2)$$

where  $\tau_2 = \text{Im}(\tau)$  and  $\Gamma(1-k, x) = \int_x^\infty t^{-k} e^{-t} dt$  denotes the incomplete gamma function, and where the first term must be replaced by  $-\bar{b}_0 \log(4\pi\tau_2)$  if  $k = 1$ . Note that the series in (7.2) converges despite the exponentially large factor  $q^{-n}$  because  $\Gamma(1-k, x) = O(x^{-k}e^{-x})$ . If we assume either that  $k > 1$  or that  $b_0 = 0$ , then we can define  $g^*$  alternatively by the integral

$$g^*(\tau) = \left(\frac{i}{2\pi}\right)^{k-1} \int_{-\bar{\tau}}^\infty (z + \tau)^{-k} \overline{g(-\bar{z})} dz. \quad (7.3)$$

(The integral is independent of the path chosen because the integrand is holomorphic in  $z$ .) Since  $h$  is holomorphic, (7.1) implies that the completion of  $h$  is related to its shadow by

$$(4\pi\tau_2)^k \frac{\partial \widehat{h}(\tau)}{\partial \bar{\tau}} = -2\pi i \overline{g(\tau)}. \quad (7.4)$$

We denote by  $\mathbb{M}_{k|0}^!$  the space of weakly holomorphic pure mock modular forms of weight  $k$  and arbitrary level. (The “0” in the notation corresponds to the word “pure” and will be replaced by an arbitrary integer or half-integer in §7.3.) This space clearly contains the space  $M_k^!$  of ordinary weakly holomorphic modular forms (the special case  $g = 0$ ,  $h = \widehat{h}$ ) and we have an exact sequence

$$0 \longrightarrow M_k^! \longrightarrow \mathbb{M}_{k|0}^! \xrightarrow{\mathcal{S}} \overline{M_{2-k}}, \quad (7.5)$$

where the *shadow map*  $\mathcal{S}$  sends  $h$  to  $\bar{g}$ .<sup>13</sup>

<sup>12</sup>One can also consider the case where the shadow is allowed to be a weakly holomorphic modular form, but we do not do this since none of our examples will be of this type.

<sup>13</sup>We will use the word “shadow” to denote either  $g(\tau)$  or  $\overline{g(\tau)}$ , but the *shadow map*, which should be linear over  $\mathbb{C}$ , always sends  $h$  to  $\bar{g}$ , the complex conjugate of its holomorphic shadow. We will also often be sloppy and say that the shadow of a certain mock modular form “is” some modular form  $g$  when in fact it is merely proportional to  $g$ , since the constants occurring are arbitrary and are often messy.

In the special case when the shadow  $g$  is a unary theta series as in (3.10) or (3.11) (which can only happen if  $k$  equals  $3/2$  or  $1/2$ , respectively), the mock modular form  $h$  is called a *mock theta function*. All of Ramanujan's examples, and all of ours in this paper, are of this type. In these cases the incomplete gamma functions in (7.2) can be expressed in terms of the complementary error function  $\operatorname{erfc}(x) = 2\pi^{-1/2} \int_x^\infty e^{-t^2} dt$ :

$$\Gamma\left(-\frac{1}{2}, x\right) = \frac{2}{\sqrt{x}} e^{-x} - 2\sqrt{\pi} \operatorname{erfc}(\sqrt{x}), \quad \Gamma\left(\frac{1}{2}, x\right) = \sqrt{\pi} \operatorname{erfc}(\sqrt{x}). \quad (7.6)$$

A very simple, though somewhat artificial, example of a mock modular form is the weight 2 Eisenstein series  $E_2(\tau)$  mentioned in §3.1, which is also a quasimodular form. Here the shadow  $g(\tau)$  is a constant and its non-holomorphic Eichler integral  $g^*(\tau)$  is simply a multiple of  $1/\tau_2$ . This example, however, is atypical, since most quasimodular forms (like  $E_2(\tau)^2$ ) are not mock modular forms. We end this subsection by giving a number of less trivial examples. Many more will occur later.

**Example 1.** In Ramanujan's famous last letter to Hardy in 1920, he gives 17 examples of mock theta functions, though without giving any complete definition of this term. All of them have weight  $1/2$  and are given as  $q$ -hypergeometric series. A typical example (Ramanujan's second mock theta function of "order 7" — a notion that he also does not define) is

$$\mathcal{F}_{7,2}(\tau) = -q^{-25/168} \sum_{n=1}^{\infty} \frac{q^{n^2}}{(1-q^n) \cdots (1-q^{2n-1})} = -q^{143/168} (1+q+q^2+2q^3+\cdots). \quad (7.7)$$

This is a mock theta function of weight  $1/2$  on  $\Gamma_0(4) \cap \Gamma(7)$  with shadow the unary theta series

$$\sum_{n \equiv 2 \pmod{7}} \chi_{12}(n) n q^{n^2/168}, \quad (7.8)$$

with  $\chi_{12}(n)$  as in (3.13). The product  $\eta(\tau)\mathcal{F}_{7,2}(\tau)$  is a strongly holomorphic mixed mock modular form of weight  $(1, 1/2)$ , and by an identity of Hickerson [64] is equal to an indefinite theta series:

$$\eta(\tau) \mathcal{F}_{7,2}(\tau) = \sum_{r,s \in \mathbb{Z} + \frac{5}{14}} \frac{1}{2} (\operatorname{sgn}(r) + \operatorname{sgn}(s)) (-1)^{r-s} q^{(3r^2+8rs+3s^2)/2}. \quad (7.9)$$

**Example 2.** The second example is the generating function of the Hurwitz-Kronecker class numbers  $H(N)$ . These numbers are defined for  $N > 0$  as the number of  $PSL(2, \mathbb{Z})$ -equivalence classes of integral binary quadratic forms of discriminant  $-N$ , weighted by the reciprocal of the number of their automorphisms (if  $-N$  is the discriminant of an imaginary quadratic field

$K$  other than  $\mathbb{Q}(i)$  or  $\mathbb{Q}(\sqrt{-3})$ , this is just the class number of  $K$ , and for other values of  $N$  by  $H(0) = -1/12$  and  $H(N) = 0$  for  $N < 0$ . It was shown in [115] that the function

$$\mathbf{H}(\tau) := \sum_{N=0}^{\infty} H(N) q^N = -\frac{1}{12} + \frac{1}{3}q^3 + \frac{1}{2}q^4 + q^7 + q^8 + q^{11} + \cdots \quad (7.10)$$

is a mock modular form of weight  $3/2$  on  $\Gamma_0(4)$ , with shadow the classical theta function  $\theta(\tau) = \sum q^{n^2}$ . Here  $\mathbf{H}(\tau)$  is strongly holomorphic, but this is very exceptional. In fact, up to minor variations it is essentially the *only* known non-trivial example of a strongly holomorphic pure mock modular form, which is why we will introduce mixed mock modular forms below. As mentioned earlier, this mock modular form has appeared in the physics literature in the context of topologically twisted supersymmetric Yang-Mills theory on  $\mathbb{CP}^2$  [109].

**Example 3.** This example is taken from [118]. Set

$$F_2^{(6)}(\tau) = - \sum_{r>s>0} \chi_{12}(r^2 - s^2) s q^{rs/6} = q + 2q^2 + q^3 + 2q^4 - q^5 + \cdots \quad (7.11)$$

with  $\chi_{12}$  as in (3.13), and let  $E_2(\tau)$  be the quasimodular Eisenstein series of weight 2 on  $SL(2, \mathbb{Z})$  as defined in §3.1. Then the function

$$h^{(6)}(\tau) = \frac{12F_2^{(6)}(\tau) - E_2(\tau)}{\eta(\tau)} = q^{-1/24} (-1 + 35q + 130q^2 + 273q^3 + 595q^4 + \cdots) \quad (7.12)$$

is a weakly holomorphic mock modular form of weight  $3/2$  on  $SL(2, \mathbb{Z})$  with shadow proportional to  $\eta(\tau)$ . More generally, if we define

$$F_k^{(6)}(\tau) = - \sum_{r>s>0} \chi_{12}(r^2 - s^2) s^{k-1} q^{rs/6} \quad (k > 2, k \text{ even}), \quad (7.13)$$

then for all  $\nu \geq 0$  we have

$$\frac{24^\nu}{\binom{2\nu}{\nu}} [h^{(6)}, \eta]_\nu = 12F_k^{(6)} - E_k + \text{cusp form of weight } k, \quad k = 2\nu + 2, \quad (7.14)$$

where  $[h^{(6)}, \eta]_\nu$  denotes the  $\nu$ th Rankin-Cohen bracket of the mock modular form  $h^{(6)}$  and the modular form  $\eta$  in weights  $(3/2, 1/2)$ . This statement, and the similar statements for other mock modular forms which come later, are proved by the method of holomorphic projection, which we do not explain here, and are intimately connected with the mock Jacobi forms introduced in the next subsection. That connection will also explain the superscript “6” in (7.11)–(7.14).

**Example 4.** Our last example is very similar to the preceding one. Set

$$F_2^{(2)}(\tau) = \sum_{\substack{r>s>0 \\ r-s \text{ odd}}} (-1)^r s q^{rs/2} = q + q^2 - q^3 + q^4 - q^5 + \cdots \quad (7.15)$$



Then the function

$$h^{(2)}(\tau) = \frac{24F_2^{(2)}(\tau) - E_2(\tau)}{\eta(\tau)^3} = q^{-1/8} (-1 + 45q + 231q^2 + 770q^3 + 2277q^4 + \dots) \quad (7.16)$$

is a weakly holomorphic mock modular form of weight  $1/2$  on  $SL(2, \mathbb{Z})$  with shadow proportional to  $\eta(\tau)^3$  and, as in Example 3, if we set

$$F_k^{(2)}(\tau) = - \sum_{\substack{r>s>0 \\ r-s \text{ odd}}} (-1)^r s^{k-1} q^{rs/2} \quad (k > 2, k \text{ even}), \quad (7.17)$$

then for all  $\nu \geq 0$  we have

$$\frac{8^\nu}{\binom{2\nu}{\nu}} [h^{(2)}, \eta^3]_\nu = 24F_k^{(2)} - E_k + \text{cusp form of weight } k, \quad k = 2\nu + 2, \quad (7.18)$$

where  $[h^{(2)}, \eta^3]_\nu$  denotes the Rankin-Cohen bracket in weights  $(1/2, 3/2)$ .

In fact, the mock modular form  $h^{(2)}$ , with the completely different definition

$$h^{(2)}(\tau) = \frac{\vartheta_2(\tau)^4 - \vartheta_4(\tau)^4}{\eta(\tau)^3} - \frac{24}{\vartheta_3(\tau)} \sum_{n \in \mathbb{Z}} \frac{q^{n^2/2-1/8}}{1 + q^{n-1/2}}, \quad (7.19)$$

arose from the works of Eguchi-Taormina [51] and Eguchi–Ooguri–Taormina–Yang [48] in the late 1980’s in connection with characters of the  $\mathcal{N} = 4$  superconformal algebra in two dimensions and the elliptic genus of  $K3$  surfaces, and appears explicitly in the work of Ooguri [88] (it is  $q^{-1/8}(-1 + F(\tau)/2)$  in Ooguri’s notation) and Wendland [110]. It has recently aroused considerable interest because of the “Mathieu moonshine” discovery by Eguchi, Ooguri and Tachikawa (see [47, 20]) that the coefficients 45, 231, 770,  $\dots$  in (7.16) are the dimensions of certain representations of the Mathieu group  $M_{24}$ . The equality of the right-hand sides of (7.16) and (7.19), which is not a priori obvious, follows because both expressions define mock modular forms whose shadow is the same multiple of  $\eta(\tau)^3$ , and the first few Fourier coefficients agree.

## 7.2 Mock Jacobi forms

By a (pure) *mock Jacobi form*<sup>14</sup> (resp. *weak mock Jacobi form*) of weight  $k$  and index  $m$  we will mean a holomorphic function  $\varphi$  on  $\mathbb{H} \times \mathbb{C}$  that satisfies the elliptic transformation property (4.2), and hence has a Fourier expansion as in (4.3) with the periodicity property (4.4) and a theta expansion as in (4.10), and that also satisfies the same cusp conditions (4.5) (resp. (4.7)) as in

<sup>14</sup>There are more general definitions of mock Jacobi forms in the literature (see [14], [23]) in which the completion is non-holomorphic in  $z$  as well as in  $\tau$ , but they will not be used in this paper.

the classical case, but in which the modularity property with respect to the action of  $SL(2, \mathbb{Z})$  on  $\mathbb{H} \times \mathbb{Z}$  is weakened: the coefficients  $h_\ell(\tau)$  in (4.10) are now mock modular forms rather than modular forms of weight  $k - \frac{1}{2}$ , and the modularity property of  $\varphi$  is that the *completed* function

$$\widehat{\varphi}(\tau, z) = \sum_{\ell \in \mathbb{Z}/2m\mathbb{Z}} \widehat{h}_\ell(\tau) \vartheta_{m,\ell}(\tau, z), \quad (7.20)$$

rather than  $\varphi$  itself, transforms according to (4.1). If  $g_\ell$  denotes the shadow of  $h_\ell$ , then we have

$$\widehat{\varphi}(\tau, z) = \varphi(\tau, z) + \sum_{\ell \in \mathbb{Z}/2m\mathbb{Z}} g_\ell^*(\tau) \vartheta_{m,\ell}(\tau, z)$$

with  $g_\ell^*$  as in (7.2) and hence, by (7.1),

$$\psi(\tau, z) := \tau_2^{k-1/2} \frac{\partial}{\partial \bar{\tau}} \widehat{\varphi}(\tau, z) \doteq \sum_{\ell \in \mathbb{Z}/2m\mathbb{Z}} \overline{g_\ell(\tau)} \vartheta_{m,\ell}(\tau, z). \quad (7.21)$$

(Here  $\doteq$  indicates an omitted constant.) The function  $\psi(\tau, z)$  is holomorphic in  $z$ , satisfies the same elliptic transformation property (4.2) as  $\varphi$  does (because each  $\vartheta_{m,\ell}$  satisfies this), satisfies the heat equation  $(8\pi im \frac{\partial}{\partial \tau} - \frac{\partial^2}{\partial z^2})\psi = 0$  (again, because each  $\vartheta_{m,\ell}$  does), and, by virtue of the modular invariance property of  $\widehat{\varphi}(\tau, z)$ , also satisfies the transformation property

$$\psi\left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}\right) = |c\tau + d| (c\bar{\tau} + d)^{2-k} e^{\frac{2\pi imcz^2}{c\tau + d}} \psi(\tau, z) \quad \forall \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2; \mathbb{Z}) \quad (7.22)$$

with respect to the action of the modular group. These properties say precisely that  $\psi$  is a *skew-holomorphic Jacobi form* of weight  $3 - k$  and index  $m$  in the sense of Skoruppa [101, 102], and the above discussion can be summarized by saying that we have an exact sequence

$$0 \longrightarrow J_{k,m}^{\text{weak}} \longrightarrow \mathbb{J}_{k|0,m}^{\text{weak}} \xrightarrow{\mathcal{S}} J_{3-k,m}^{\text{skew}} \quad (7.23)$$

(and similarly with the word “weak” omitted), where  $\mathbb{J}_{k|0,m}^{\text{weak}}$  (resp.  $\mathbb{J}_{k|0,m}$ ) denotes the space of weak (resp. strong) pure mock Jacobi forms and the “shadow map”  $\mathcal{S}$  sends  $\varphi$  to  $\psi$ .

It turns out that most of the classical examples of mock theta functions occur as the components of a vector-valued mock modular form which gives the coefficients in the theta series expansion of a mock Jacobi form. We illustrate this for the four examples introduced in the previous subsection.

**Example 1.** The function  $\mathcal{F}_{7,2}(\tau)$  in the first example of §7.1 is actually one of three “order 7 mock theta functions”  $\{\mathcal{F}_{7,j}\}_{j=1,2,3}$  defined by Ramanujan, each given by a  $q$ -hypergeometric formula like (7.7), each having a shadow  $\Theta_{7,j}$  like in (7.8) (but now with the summation over  $n \equiv j$  rather than  $n \equiv 2$  modulo 7), and each satisfying an indefinite theta series identity like

(7.9). We extend  $\{\mathcal{F}_{7,j}\}$  to all  $j$  by defining it to be an odd periodic function of  $j$  of period 7, so that the shadow of  $\mathcal{F}_{7,j}$  equals  $\Theta_{7,j}$  for all  $j \in \mathbb{Z}$ . Then the function

$$\mathcal{F}_{42}(\tau, z) = \sum_{\ell \pmod{84}} \chi_{12}(\ell) \mathcal{F}_{7,\ell}(\tau) \vartheta_{42,\ell}(\tau, z) \quad (7.24)$$

belongs to  $\mathbb{J}_{1,42}^{\text{weak}}$ . The Taylor coefficients  $\xi_\nu$  as defined in equation (4.16) are proportional to  $\sum_{j=1}^3 [\mathcal{F}_{7,j}, \Theta_{7,j}]_\nu$  and have the property that their completions  $\widehat{\xi}_\nu = \sum_{j=1}^3 [\widehat{\mathcal{F}}_{7,j}, \Theta_{7,j}]_\nu$  transform like modular forms of weight  $2\nu + 2$  on the full modular group  $SL(2, \mathbb{Z})$ .

**Example 2.** Set  $\mathcal{H}_0(\tau) = \sum_{n=0}^{\infty} H(4n)q^n$  and  $\mathcal{H}_1(\tau) = \sum_{n=1}^{\infty} H(4n-1)q^{n-\frac{1}{4}}$  so that  $\mathcal{H}_0(\tau) + \mathcal{H}_1(\tau) = \mathbf{H}(\tau/4)$ . Then the function

$$\mathcal{H}(\tau, z) = \mathcal{H}_0(\tau)\vartheta_{1,0}(\tau, z) + \mathcal{H}_1(\tau)\vartheta_{1,1}(\tau, z) = \sum_{\substack{n, r \in \mathbb{Z} \\ 4n-r^2 \geq 0}} H(4n-r^2) q^n y^r, \quad (7.25)$$

is a mock Jacobi form of weight 2 and index 1 with shadow  $\overline{\vartheta_{1,0}(\tau, 0)} \vartheta_{1,0}(\tau, z) + \overline{\vartheta_{1,1}(\tau, 0)} \vartheta_{1,1}(\tau, z)$ . The  $\nu$ th Taylor coefficient  $\xi_\nu$  of  $\mathcal{H}$  is given by

$$\frac{4^\nu}{\binom{2\nu}{\nu}} \sum_{j=0}^1 [\vartheta_{1,j}, \mathcal{H}_j]_\nu = \delta_{k,2} E_k - F_k^{(1)} + (\text{cusp form of weight } k \text{ on } SL(2, \mathbb{Z})), \quad (7.26)$$

where  $k = 2\nu + 2$  and

$$F_k^{(1)}(\tau) := \sum_{n>0} \left( \sum_{d|n} \min(d, \frac{n}{d})^{k-1} \right) q^n \quad (k \text{ even, } k \geq 2). \quad (7.27)$$

In fact the cusp form appearing in (7.26) is a very important one, namely (up to a factor  $-2$ ) the sum of the normalized Hecke eigenforms in  $S_k(SL(2, \mathbb{Z}))$ , and equation (7.26) is equivalent to the famous formula of Eichler and Selberg expressing the traces of Hecke operators on  $S_k(SL(2, \mathbb{Z}))$  in terms of class numbers of imaginary quadratic fields.

**Example 3.** Write the function  $h^{(6)}$  defined in (7.12) as

$$h^{(6)}(\tau) = \sum_{\substack{\Delta \geq -1 \\ \Delta \equiv -1 \pmod{24}}} C^{(6)}(\Delta) q^{\Delta/24} \quad (7.28)$$

with the first few coefficients  $C^{(6)}(\Delta)$  given by

$\Delta$	-1	23	47	71	95	119	143	167	191	215	239
$C^{(6)}(\Delta)$	-1	35	130	273	595	1001	1885	2925	4886	7410	11466

Then the function

$$\mathcal{F}_6(\tau, z) = \sum_{\substack{n, r \in \mathbb{Z} \\ 24n - r^2 \geq -1}} \chi_{12}(r) C^{(6)}(24n - r^2) q^n y^r \quad (7.29)$$

is a mock Jacobi form of index 6 (explaining the notation  $h^{(6)}$ ). Note that, surprisingly, this is even *simpler* than the expansion of the index 1 mock Jacobi form just discussed, because its twelve Fourier coefficients  $h_\ell$  are all multiples of just one of them, while the two Fourier coefficients  $h_\ell$  of  $\mathcal{H}(\tau, z)$  were not proportional. (This is related to the fact that the shadow  $\eta(\tau)$  of  $h^{(6)}(\tau)$  is a modular form on the full modular group, while the shadow  $\theta(\tau)$  of  $\mathcal{H}(\tau)$  is a modular form on a congruence subgroup.) Specifically, we have  $h_\ell(\tau) = \chi_{12}(\ell) h^{(6)}(\tau)$  for all  $\ell \in \mathbb{Z}/12\mathbb{Z}$ , where  $\chi_{12}$  is the character defined in (3.13), so that  $\mathcal{F}_6$  has the factorization

$$\mathcal{F}_6(\tau, z) = h^{(6)}(\tau) \sum_{\ell \pmod{12}} \chi_{12}(\ell) \vartheta_{6,\ell}(\tau, z) = \eta(\tau) h^{(6)}(\tau) \frac{\vartheta_1(\tau, 4z)}{\vartheta_1(\tau, 2z)}. \quad (7.30)$$

Combining this with (4.17) and noting that  $\vartheta_{6,1}^0 - \vartheta_{6,5}^0 = \vartheta_{6,11}^0 - \vartheta_{6,7}^0 = \eta$ , we see that the functions described in (7.14) are proportional to the Taylor coefficients  $\xi_\nu(\tau)$  of  $\mathcal{F}_6(\tau, z)$ .

**Example 4.** The fourth example is very similar. Write the mock modular form (7.16) as

$$h^{(2)}(\tau) = \sum_{\substack{\Delta \geq -1 \\ \Delta \equiv -1 \pmod{8}}} C^{(2)}(\Delta) q^{\Delta/8} \quad (7.31)$$

with initial coefficients given by

$\Delta$	-1	7	15	23	31	39	47	55	63	71
$C^{(2)}(\Delta)$	-1	45	231	770	2277	5796	13915	30843	65550	132825

Then the function

$$\begin{aligned} \mathcal{F}_2(\tau, z) &= \sum_{\substack{n, r \in \mathbb{Z} \\ 8n - r^2 \geq -1}} \chi_4(r) C^{(2)}(8n - r^2) q^n y^r = h^{(2)}(\tau) (\vartheta_{2,1}(\tau, z) - \vartheta_{2,3}(\tau, z)) \\ &= h^{(2)}(\tau) \vartheta_1(\tau, 2z) = \eta(\tau)^3 h^{(2)}(\tau) C(\tau, z) \end{aligned} \quad (7.32)$$

(where  $\chi_4(r) = \pm 1$  for  $r \equiv \pm 1 \pmod{4}$  and  $\chi_4(r) = 0$  for  $r$  even) is a mock Jacobi form of index 2. (Here  $C$  is the Jacobi form of weight  $-1$  and index 2 introduced in §4.3.) As in Example 3, the functions given in (7.18) are proportional to the Taylor coefficients  $\xi_\nu$  of  $\mathcal{F}_2$ , because  $\vartheta_{2,1}^1 = -\vartheta_{2,3}^1 = \eta^3$ , where  $\vartheta_{2,\ell}^1$  is defined by (4.18).

### 7.3 Mock modular forms: the general case

If we replace the condition “exponential growth” in 1. of 7.1 by “polynomial growth,” we get the class of *strongly holomorphic mock modular forms*, which we can denote  $\mathbb{M}_{k|0}$ , and an exact sequence  $0 \rightarrow M_k \rightarrow \mathbb{M}_{k|0} \rightarrow \overline{M_{2-k}}$ . This is not very useful, however, because there are almost no examples of “pure” mock modular forms that are strongly holomorphic, essentially the only ones being the function  $\mathcal{H}$  of Example 2 of §7.1 and its variants. It becomes useful if we generalize to *mixed mock modular forms* of mixed weight  $k|\ell$ . (Here  $k$  and  $\ell$  can be integers or half-integers.) These are holomorphic functions  $h(\tau)$ , with polynomial growth at the cusps, that have completions  $\widehat{h}$  of the form  $\widehat{h} = h + \sum_j f_j g_j^*$  with  $f_j \in M_\ell$ ,  $g_j \in M_{2-k-\ell}$  that transform like modular forms of weight  $k$ . The space  $\mathbb{M}_{k|\ell}$  of such forms thus fits into an exact sequence

$$0 \longrightarrow M_k \longrightarrow \mathbb{M}_{k|\ell} \xrightarrow{\mathcal{S}} M_\ell \otimes \overline{M_{2-k+\ell}}, \quad (7.33)$$

where the shadow map  $\mathcal{S}$  now sends  $h$  to  $\sum_j f_j \overline{g_j}$ . If  $\ell = 0$  this reduces to the previous definition, since  $M_0 = \mathbb{C}$ , but with the more general notion of mock modular forms there are now plenty of strongly holomorphic examples, and, just as for ordinary modular forms, they have much nicer properties (notably, polynomial growth of their Fourier coefficients) than the weakly holomorphic ones. If the shadow of a mixed mock modular form  $h \in \mathbb{M}_{k|\ell}$  happens to contain only one term  $f(\tau)\overline{g(\tau)}$ , and if  $f(\tau)$  has no zeros in the upper half-plane, then  $f^{-1}h$  is a weakly holomorphic pure mock modular form of weight  $k-\ell$  (and in fact, all weakly holomorphic pure mock modular forms arise in this way). For functions in the larger space  $\mathbb{M}_{k|\ell}^!$  of weakly holomorphic mixed mock modular forms, defined in the obvious way, this always happens, so  $\mathbb{M}_{k|\ell}^!$  simply coincides with  $M_\ell \otimes \mathbb{M}_{k-\ell}^!$  and does not give anything new, but the smaller space  $\mathbb{M}_{k|\ell}$  does not have such a decomposition and is more interesting. Moreover, there are many examples. In fact, apart from Ramanujan’s original examples of mock theta functions, which were defined as  $q$ -hypergeometric series and were weakly holomorphic pure mock modular forms of weight  $1/2$ , most of the examples of mock modular forms occurring “in nature,” such as Appell-Lerch sums or indefinite theta series, are strongly holomorphic mixed modular forms.

We can also define “even more mixed” mock modular forms by replacing  $\mathbb{M}_{k|\ell}$  by the space  $\mathbb{M}_k = \sum_\ell \mathbb{M}_{k|\ell}$ , i.e., by allowing functions whose shadow is a finite sum of products  $f_j(\tau)\overline{g_j(\tau)}$  with the  $f_j$  of varying weights  $\ell_j$  and  $g_j$  of weight  $2-k+\ell_j$ . This space fits into an exact sequence

$$0 \longrightarrow M_k \longrightarrow \mathbb{M}_k \xrightarrow{\mathcal{S}} \bigoplus_\ell M_\ell \otimes \overline{M_{2-k+\ell}}. \quad (7.34)$$

In fact, the functions in this space are the most natural objects, and from now on we will use “mock modular forms” to refer to this more general class, including the adjectives “mixed” and “pure” only when needed for emphasis. To justify this claim, we consider the space  $\mathfrak{M}_k$  of all *real-analytic modular forms of weight  $k$* , i.e., real-analytic functions that transform like modular

forms of weight  $k$  and have polynomial growth at all cusps. More generally, one can consider the space  $\mathfrak{M}_{k,\ell}$  of real-analytic functions of polynomial growth that transform according to  $F(\gamma\tau) = (c\tau + d)^k (c\bar{\tau} + d)^\ell F(\tau)$  for all matrices  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  belonging to some subgroup of finite index of  $SL(2, \mathbb{Z})$ . Since the function  $\tau_2 = \text{Im}(\tau)$  transforms by  $(\gamma\tau)_2 = \tau_2/|c\tau + d|^2$ , the spaces  $\mathfrak{M}_{k,\ell}$  and  $\tau_2^r \mathfrak{M}_{k+r,\ell+r}$  coincide for all  $r$ . (Here,  $k$ ,  $\ell$  and  $r$  can each be integral or half-integral.) Obviously the space  $\mathfrak{M}_{k,\ell}$  contains the space  $M_k \otimes \overline{M_\ell}$ , so by the comment just made, it also contains the direct sum  $\bigoplus_{r \in \mathbb{Z}} \tau_2^r M_{k+r} \otimes \overline{M_{\ell+r}}$ . Elements of this subspace will be called *decomposable*. Now, if a function  $F \in \mathfrak{M}_k = \mathfrak{M}_{k,0}$  has the property that its  $\bar{\tau}$ -derivative (which automatically belongs to  $\mathfrak{M}_{k,2}$ ) is decomposable, say as  $\sum_j \tau_2^{r_j} f_j \bar{g}_j$  (where the weights of  $f_j$  and  $g_j$  are necessarily equal to  $k + r_j$  and  $2 + r_j$ , respectively), then  $F$  is the sum of a holomorphic function  $h$  and a multiple of  $\sum_j f_j g_j^*$  and the function  $h$  is a (mixed) mock modular form in the sense just explained. Conversely, a holomorphic function  $h$  is a (mixed) mock modular form of weight  $k$  if and only if there exist numbers  $r_j$  and modular forms  $f_j \in M_{k+r_j}$  and  $g_j \in M_{2+r_j}$  such that the sum  $\hat{h} := h + \sum_j f_j g_j^*$  belongs to  $\mathfrak{M}_k$ . We can thus summarize the entire discussion by the following very simple definition:

*A mock modular form is the holomorphic part of a real-analytic modular form whose  $\bar{\tau}$ -derivative is decomposable.*

As examples of this more general class we mention the functions  $F_2^{(6)}$  and  $F_2^{(2)}$  defined in equations (7.11) and (7.15), which are both (strong) mixed mock modular forms of total weight 2 and trivial character on the full modular group  $SL(2, \mathbb{Z})$ , but with different mixed weights:  $12F_2^{(6)}$  is the sum of the functions  $E_2(\tau)$  and  $\eta(\tau) h^{(6)}(\tau)$  of weights  $2|0$  and  $\frac{3}{2}|\frac{1}{2}$ , while  $24F_2^{(2)}$  is the sum of the functions  $E_2(\tau)$  and  $\eta(\tau)^3 h^{(2)}(\tau)$  of weights  $2|0$  and  $\frac{1}{2}|\frac{3}{2}$ .

We remark that in the pure case we have  $\tau_2^k \cdot \partial_{\bar{\tau}} F \in \overline{M_{2-k}} = \text{Ker}(\partial_{\bar{\tau}}: \mathfrak{M}_{0,2-k} \rightarrow \mathfrak{M}_{2,2-k})$ , so that the subspace  $\mathbb{M}_{k,0}$  of  $\mathbb{M}_k$  can be characterized by a *single* second-order differential equation  $\Delta_k \hat{h} = 0$ , where  $\Delta_k = \tau_2^{2-k} \partial_{\tau} \tau_2^k \partial_{\bar{\tau}}$  is the Laplacian in weight  $k$ . For this reason, (weak) pure mock modular forms can be, and often are, viewed as the holomorphic parts of (weak) harmonic Maass forms, i.e., of functions in  $\mathfrak{M}_k^!$  annihilated by  $\Delta_k$ . But in the general case, there is no connection between mock modular forms or their completions and any second-order operators. Only the Cauchy-Riemann operator  $\partial/\partial\bar{\tau}$  is needed, and the new definition of mock modular forms as given above is both more flexible and more readily generalizable than the more familiar definition of pure mock modular forms in terms of harmonic Maass forms.

Finally, we can define mixed mock Jacobi forms in exactly the same way as we defined pure ones (i.e., they have theta expansions in which the coefficients are mixed mock modular forms, with completions that transform like ordinary Jacobi forms). We will see in the next section that such objects arise very naturally as the “finite parts” of meromorphic Jacobi forms.

## 7.4 Superconformal characters and mock modular forms

As already mentioned, mock modular forms first made their appearance in physics some ten years before the work of Zwegers, in connection with the characters of superconformal algebras [51, 48]. The representations of the  $\mathcal{N} = 4$  superconformal algebra with central charge  $c = 6k$  are labeled by two quantum numbers, the energy  $h$  and spin  $l$  of the lowest weight state  $|\Omega\rangle$ , which satisfy  $h > k/4$ ,  $l \in \{1/2, 1, \dots, k/2\}$  for the non-BPS (massive) case and  $h = k/4$ ,  $l \in \{0, 1/2, \dots, k/2\}$  for the BPS (massless) case. The corresponding character, defined by

$$\text{ch}_{k,h,l}^{\tilde{R}}(\tau, z) = \text{Tr}(q^{L_0-c/24} e^{4\pi iz J_0}) \quad (L_0|\Omega\rangle = h|\Omega\rangle, \quad J_0|\Omega\rangle = l|\Omega\rangle),$$

turns out to be, up to a power of  $q$ , a Jacobi form in the non-BPS case but a “mock” object in the BPS case. In particular, for  $k = 1$  the non-BPS characters are given by

$$\text{ch}_{1,h,\frac{1}{2}}^{\tilde{R}}(\tau, z) = -q^{h-\frac{3}{8}} \frac{\vartheta_1(\tau, z)^2}{\eta(\tau)^3} = -q^{h-\frac{3}{8}} \eta(\tau)^3 \varphi_{-2,1}(\tau, z) \quad (h > \frac{1}{4}) \quad (7.35)$$

and the BPS characters by

$$\text{ch}_{1,\frac{1}{4},0}^{\tilde{R}}(\tau, z) = -\frac{\vartheta_1(\tau, z)^2}{\eta(\tau)^3} \mu(\tau, z), \quad \text{ch}_{1,\frac{1}{4},\frac{1}{2}}^{\tilde{R}}(z, \tau) + 2\text{ch}_{1,\frac{1}{4},0}^{\tilde{R}}(\tau, z) = -q^{-\frac{1}{8}} \frac{\vartheta_1(\tau, z)^2}{\eta(\tau)^3}, \quad (7.36)$$

where  $\mu(\tau, z)$  is the function defined by

$$\mu(\tau, z) = \frac{e^{\pi iz}}{\vartheta_1(z, \tau)} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{\frac{n^2+n}{2}} e^{2\pi in z}}{1 - q^n e^{2\pi iz}} \quad (q = e^{2\pi i\tau}, \quad y = e^{2\pi iz}), \quad (7.37)$$

an “Appell-Lerch sum” of a type that also played a central role in Zwegers’s work<sup>15</sup> and that we will discuss in the next section. The “mock” aspect is the fact that  $\mu(\tau, z)$  must be completed by an appropriate non-holomorphic correction term in order to transform like a Jacobi form.

The mock modular form  $h^{(2)}$  arises in connection with the decomposition of the elliptic genus of a  $K3$  surface as a linear combination of these characters. This elliptic genus is, as discussed in 4.5, equal to the weak Jacobi form  $2B = 2\varphi_{0,1}$ . It can be represented as a specialization of the partition function (= generating function of states) of superstrings compactified on a  $K3$  surface [48] and as such can be decomposed into pieces corresponding to the representations of the  $N = 4$  superconformal algebra with central charge  $6k = 6$ . The decomposition has the form [88, 110, 50]

$$2\varphi_{0,1}(\tau, z) = 20 \text{ch}_{1,\frac{1}{4},0}^{\tilde{R}}(z, \tau) + A_0 \text{ch}_{1,\frac{1}{4},\frac{1}{2}}^{\tilde{R}}(z, \tau) + \sum_{n=1}^{\infty} A_n \text{ch}_{1,n+\frac{1}{4},\frac{1}{2}}^{\tilde{R}}(z, \tau), \quad (7.38)$$

<sup>15</sup>More precisely, the function  $\mu(\tau, z)$  is the specialization to  $u = v = z$  of the 2-variable function  $\mu(u, v; \tau)$  considered by Zwegers.



where  $A_0 = -2$ ,  $A_1 = 90$ ,  $A_2 = 462$ ,  $\dots$  are the Fourier coefficients  $2C^{(2)}(8n - 1)$  of the mock modular form  $2h^{(2)}(\tau)$ . Inserting (7.35) and (7.36), we can rewrite this as

$$\eta(\tau)^{-3} \frac{\varphi_{0,1}(\tau, z)}{\varphi_{-2,1}(\tau, z)} = -12\mu(\tau, z) - h^{(2)}(\tau). \quad (7.39)$$

In the next section we will see how to interpret this equation as an example of a certain canonical decomposition of meromorphic Jacobi forms (8.52).

## 8. From meromorphic Jacobi forms to mock modular forms

In this section we consider Jacobi forms  $\varphi(\tau, z)$  that are meromorphic with respect to the variable  $z$ . It was discovered by Zwegers [119] that such forms, assuming that their poles occur only at points  $z \in \mathbb{Q}\tau + \mathbb{Q}$  (i.e., at torsion points on the elliptic curve  $\mathbb{C}/\mathbb{Z}\tau + \mathbb{Z}$ ), have a modified theta expansion related to mock modular forms. Our treatment is based on his, but the presentation is quite different and the results go further in one key respect. We show that  $\varphi$  decomposes canonically into two pieces, one constructed directly from its poles and consisting of a finite linear combination of Appell-Lerch sums with modular forms as coefficients, and the other being a mock Jacobi form in the sense introduced in the preceding section. Each piece separately transforms like a Jacobi form with respect to elliptic transformations. Neither piece separately transforms like a Jacobi form with respect to modular transformations, but each can be completed by the addition of an explicit and elementary non-holomorphic correction term so that it does transform correctly with respect to the modular group.

In §8.1 we explain how to modify the Fourier coefficients  $h_\ell$  defined in (4.8) when  $\varphi$  has poles, and use these to define a “finite part” of  $\varphi$  by the theta decomposition (4.10). In §8.2 we define (in the case when  $\varphi$  has simple poles only) a “polar part” of  $\varphi$  as a finite linear combination of standard Appell-Lerch sums times modular forms arising as the residues of  $\varphi$  at its poles, and show that  $\varphi$  decomposes as the sum of its finite part and its polar part. Subsection 8.3 gives the proof that the finite part of  $\varphi$  is a mock Jacobi form and a description of the non-holomorphic correction term needed to make it transform like a Jacobi form. This subsection also contains a summary in tabular form of the various functions that have been introduced and the relations between them. In §8.4 we describe the modifications needed in the case of double poles (the case actually needed in this paper) and in §8.5 we present a few examples to illustrate the theory. Among the “mock” parts of these are two of the most interesting mock Jacobi forms from §7 (the one related to class numbers of imaginary quadratic fields and the one conjecturally related to representations of the Mathieu group  $M_{24}$ ). Many other examples will be given in §9.

Throughout the following sections, we use the convenient notation  $\mathbf{e}(x) := e^{2\pi ix}$ .



## 8.1 The Fourier coefficients of a meromorphic Jacobi form

As indicated above, the main problem we face is to find an analogue of the theta decomposition (4.10) of holomorphic Jacobi forms in the meromorphic case. We will approach this problem from two sides: computing the Fourier coefficients of  $\varphi(\tau, z)$  with respect to  $z$ , and computing the contribution from the poles. In this subsection we treat the first of these questions.

Consider a meromorphic Jacobi form  $\varphi(\tau, z)$  of weight  $k$  and index  $m$ . We assume that  $\varphi(\tau, z)$  for each  $\tau \in \mathbb{H}$  is a meromorphic function of  $z$  which has poles only at points  $z = \alpha\tau + \beta$  with  $\alpha$  and  $\beta$  rational. In the case when  $\varphi$  was holomorphic, we could write its Fourier expansion in the form (4.8). By Cauchy's theorem, the coefficient  $h_\ell(\tau)$  in that expansion could also be given by the integral formula

$$h_\ell^{(z_0)}(\tau) = q^{-\ell^2/4m} \int_{z_0}^{z_0+1} \varphi(\tau, z) \mathbf{e}(-\ell z) dz, \quad (8.1)$$

where  $z_0$  is an arbitrary point of  $\mathbb{C}$ . From the holomorphy and transformation properties of  $\varphi$  it follows that the value of this integral is independent of the choice of  $z_0$  and of the path of integration and depends only on  $\ell$  modulo  $2m$  (implying that we have the theta expansion (4.10)) and that each  $h_\ell$  is a modular form of weight  $k - \frac{1}{2}$ . Here each of these properties fails: the integral (8.1) is not independent of the path of integration (it jumps when the path crosses a pole); it is not independent of the choice of the initial point  $z_0$ ; it is not periodic in  $\ell$  (changing  $\ell$  by  $2m$  corresponds to changing  $z_0$  by  $\tau$ ); it is not modular; and of course the expansion (4.10) cannot possibly hold since the right-hand-side has no poles in  $z$ .

To take care of the first of these problems, we specify the path of integration in (8.1) as the horizontal line from  $z_0$  to  $z_0 + 1$ . If there are poles of  $\varphi(\tau, z)$  along this line, this does not make sense; in that case, we define the value of the integral as the average of the integral over a path deformed to pass just above the poles and the integral over a path just below them. (We do not allow the initial point  $z_0$  to be a pole of  $\varphi$ , so this makes sense.) To take care of the second and third difficulties, the dependence on  $z_0$  and the non-periodicity in  $\ell$ , we play one of these problems off against the other. From the elliptic transformation property (4.2) we find that

$$\begin{aligned} h_\ell^{(z_0+\tau)}(\tau) &= q^{-\ell^2/4m} \int_{z_0}^{z_0+1} \varphi(\tau, z + \tau) \mathbf{e}(-\ell(z + \tau)) dz \\ &= q^{-(\ell+2m)^2/4m} \int_{z_0}^{z_0+1} \varphi(\tau, z) \mathbf{e}(-(\ell + 2m)z) dz = h_{\ell+2m}^{(z_0)}(\tau), \end{aligned}$$

i.e., changing  $z_0$  by  $\tau$  corresponds to changing  $\ell$  by  $2m$ , as already mentioned. It follows that if we choose  $z_0$  to be  $-\ell\tau/2m$  (or  $-\ell\tau/2m + B$  for any  $B \in \mathbb{R}$ , since it is clear that the value of the integral (8.1) depends only on the height of the path of integration and not on the initial point on this line), then the quantity

$$h_\ell(\tau) := h_\ell^{(-\ell\tau/2m)}(\tau), \quad (8.2)$$

which we will call the  $\ell$ th **canonical Fourier coefficient** of  $\varphi$ , depends only on the value of  $\ell \pmod{2m}$ . This in turn implies that the sum

$$\varphi^F(\tau, z) := \sum_{\ell \in \mathbb{Z}/2m\mathbb{Z}} h_\ell(\tau) \vartheta_{m,\ell}(\tau, z), \quad (8.3)$$

which we will call the **finite (or Fourier) part** of  $\varphi$ , is well defined. If  $\varphi$  is holomorphic, then of course  $\varphi^F = \varphi$ , by virtue of (4.10).

Note that the definition of  $h_\ell(\tau)$  can also be written

$$h_\ell(\tau) = q^{\ell^2/4m} \int_{\mathbb{R}/\mathbb{Z}} \varphi(\tau, z - \ell\tau/2m) \mathbf{e}(-\ell z) dz, \quad (8.4)$$

with the same convention as above if  $\varphi(\tau, z - \ell\tau/2m)$  has poles on the real line.

Finally, we remark that none of these definitions have anything to do with the modular transformation property (4.1), and would also work if our initial function  $\varphi(\tau, z)$  satisfied only the elliptic transformation property (4.2). (In that case, however, the functions  $h_\ell$  in general need not be modular, and in fact will be mock modular when  $\varphi(\tau, z)$  is a meromorphic Jacobi form.) What we have said so far can be summarized by a commutative diagram

$$\begin{array}{ccc} \mathcal{E}_{\pm, m}^{\text{Mer}} & \xrightarrow{F} & \mathcal{E}_{\pm, m} \\ \cup & & \cup \\ J_{k, m}^{\text{Mer}} & \xrightarrow{F} & \mathbb{J}_{k, m} \end{array}, \quad (8.5)$$

where  $J_{k, m}^{\text{Mer}}$  (resp.  $\mathcal{E}_{\pm, m}^{\text{Mer}}$ ) denote the subspace of meromorphic Jacobi (resp. elliptic) forms having poles only at torsion points  $z \in \mathbb{Q}\tau + \mathbb{Q}$ , and the map  $F$  sends  $\varphi \mapsto \varphi^F$ .

## 8.2 The polar part of $\varphi$ (case of simple poles)

We now consider the contribution from the poles. To present the results we first need to fix notations for the positions and residues of the poles of our meromorphic function  $\varphi$ . We assume for now that the poles are all simple.

By assumption,  $\varphi(\tau, z)$  has poles only at points of the form  $z = z_s = \alpha\tau + \beta$  for  $s = (\alpha, \beta)$  belonging to some subset  $S$  of  $\mathbb{Q}^2$ . The double periodicity property (4.2) implies that  $S$  is invariant under translation by  $\mathbb{Z}^2$ , and of course  $S/\mathbb{Z}^2$  must be finite. The modular transformation property (4.1) of  $\varphi$  implies that  $S$  is  $SL(2, \mathbb{Z})$ -invariant. For each  $s = (\alpha, \beta) \in S$ , we set

$$D_s(\tau) = 2\pi i \mathbf{e}(m\alpha z_s) \text{Res}_{z=z_s}(\varphi(\tau, z)) \quad (s = (\alpha, \beta) \in S, z_s = \alpha\tau + \beta). \quad (8.6)$$

The functions  $D_s(\tau)$  are holomorphic modular forms of weight  $k - 1$  and some level, and only finitely many of them are distinct. More precisely, they satisfy

$$\bullet \quad D_{(\alpha+\lambda, \beta+\mu)}(\tau) = \mathbf{e}(m(\mu\alpha - \lambda\beta)) D_{(\alpha, \beta)}(\tau) \quad \text{for } (\lambda, \mu) \in \mathbb{Z}^2, \quad (8.7)$$

$$\bullet \quad D_s\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^{k-1} D_{s\gamma}(\tau) \quad \text{for } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}), \quad (8.8)$$

as one sees from the transformation properties of  $\varphi$ . (The calculation for equation (8.8) is given in §8.4.) It is precisely to obtain the simple transformation equation (8.8) that we included the factor  $\mathbf{e}(m\alpha z_s)$  in (8.6). Since we are assuming for the time being that there are no higher-order poles, all of the information about the non-holomorphy of  $\varphi$  is contained in these functions.

The strategy is to define a “polar part” of  $\varphi$  by taking the poles  $z_s$  in some fundamental parallelogram for the action of the lattice  $\mathbb{Z}\tau + \mathbb{Z}$  on  $\mathbb{C}$  (i.e., for  $s = (\alpha, \beta)$  in the intersection of  $S$  with some square box  $[A, A + 1) \times [B, B + 1)$ ) and then averaging the residues at these poles over all translations by the lattice. But we must be careful to do this in just the right way to get the desired invariance properties. For each  $m \in \mathbb{N}$  we introduce the *averaging operator*

$$\text{Av}^{(m)}[F(y)] := \sum_{\lambda} q^{m\lambda^2} y^{2m\lambda} F(q^\lambda y) \quad (8.9)$$

which sends any function of  $y$  ( $= \mathbb{Z}$ -invariant function of  $z$ ) of polynomial growth in  $y$  to a function of  $z$  transforming like an index  $m$  Jacobi form under translations by the full lattice  $\mathbb{Z}\tau + \mathbb{Z}$ . For example, we have

$$q^{\ell^2/4m} \text{Av}^{(m)}[y^\ell] = \sum_{\lambda \in \mathbb{Z}} q^{(\ell+2m\lambda)^2/4m} y^{\ell+2m\lambda} = \vartheta_{m, \ell}(\tau, z) \quad (8.10)$$

for any  $\ell \in \mathbb{Z}$ . If  $F(y)$  itself is given as the average

$$F(y) = \text{Av}_{\mathbb{Z}}[f(z)] := \sum_{\mu \in \mathbb{Z}} f(z + \mu) \quad (z \in \mathbb{C}, y = \mathbf{e}(z)) \quad (8.11)$$

of a function  $f(z)$  in  $\mathbb{C}$  (of sufficiently rapid decay at infinity), then we have

$$\text{Av}^{(m)}[F(y)] = \text{Av}_{\mathbb{Z}\tau + \mathbb{Z}}^{(m)}[f(z)] := \sum_{\lambda, \mu \in \mathbb{Z}} e^{2\pi i m(\lambda^2 \tau + 2\lambda z)} f(z + \lambda\tau + \mu). \quad (8.12)$$

We want to apply the averaging operator (8.9) to the product of the function  $D_s(\tau)$  with a standard rational function of  $y$  having a simple pole of residue 1 at  $y = y_s = \mathbf{e}(z_s)$ , but the choice of this rational function is not obvious. The right choice turns out to be  $\mathcal{R}_{-2m\alpha}(y)$ , where  $\mathcal{R}_c(y)$  for  $c \in \mathbb{R}$  is defined by

$$\mathcal{R}_c(y) = \begin{cases} \frac{1}{2} y^c \frac{y+1}{y-1} & \text{if } c \in \mathbb{Z}, \\ y^{\lceil c \rceil} \frac{1}{y-1} & \text{if } c \in \mathbb{R} \setminus \mathbb{Z}. \end{cases} \quad (8.13)$$

(Here  $\lceil c \rceil$  denotes the ‘‘ceiling’’ of  $c$ , i.e., the smallest integer  $\geq c$ . The right-hand side can also be written more uniformly as  $\frac{1}{2} \frac{y^{\lceil c \rceil + 1} + y^{\lfloor c \rfloor}}{y - 1}$ , where  $\lfloor c \rfloor = -\lceil -c \rceil$  denotes the ‘‘floor’’ of  $c$ , i.e., the largest integer  $\leq c$ .) This function looks artificial, but is in fact quite natural. First of all, by expanding the right-hand side of (8.13) in a geometric series we find

$$\mathcal{R}_c(y) = \begin{cases} -\sum_{\ell \geq c}^* y^\ell & \text{if } |y| < 1, \\ \sum_{\ell \leq c}^* y^\ell & \text{if } |y| > 1, \end{cases} \quad (8.14)$$

where the asterisk on the summation sign means that the term  $\ell = c$  is to be counted with multiplicity 1/2 when it occurs (which happens only for  $c \in \mathbb{Z}$ , explaining the case distinction in (8.13)). This formula, which can be seen as the prototypical example of wall-crossing, can also be written in terms of  $z$  as a Fourier expansion (convergent for all  $z \in \mathbb{C} \setminus \mathbb{R}$ )

$$\mathcal{R}_c(\mathbf{e}(z)) = -\sum_{\ell \in \mathbb{Z}} \frac{\operatorname{sgn}(\ell - c) + \operatorname{sgn}(z_2)}{2} \mathbf{e}(\ell z) \quad (y = \mathbf{e}(z), \quad z_2 = \operatorname{Im}(z) \neq 0), \quad (8.15)$$

without any case distinction. Secondly,  $R_c(y)$  can be expressed in a natural way as an average:

**Proposition 8.1.** *For  $c \in \mathbb{R}$  and  $z \in \mathbb{C} \setminus \mathbb{Z}$  we have*

$$\mathcal{R}_c(\mathbf{e}(z)) = \operatorname{Av}_{\mathbb{Z}} \left[ \frac{\mathbf{e}(cz)}{2\pi iz} \right]. \quad (8.16)$$

*Proof:* If  $c \in \mathbb{Z}$ , then

$$\operatorname{Av}_{\mathbb{Z}} \left[ \frac{\mathbf{e}(cz)}{2\pi iz} \right] = \frac{y^c}{2\pi i} \sum_{n \in \mathbb{Z}} \frac{1}{z - n} = \frac{y^c}{2\pi i} \frac{\pi}{\tan \pi z} = \frac{y^c}{2} \frac{y + 1}{y - 1}$$

by a standard formula of Euler. If  $c \notin \mathbb{Z}$  then the Poisson summation formula and (8.15) give

$$\begin{aligned} \operatorname{Av}_{\mathbb{Z}} \left[ \frac{\mathbf{e}(cz)}{2\pi iz} \right] &= \sum_{n \in \mathbb{Z}} \frac{\mathbf{e}(c(z + n))}{2\pi i(z + n)} = \sum_{\ell \in \mathbb{Z}} \left( \int_{iz_2 - \infty}^{iz_2 + \infty} \frac{\mathbf{e}(c(z + u))}{2\pi i(z + u)} \mathbf{e}(-\ell u) du \right) \mathbf{e}(\ell z) \\ &= -\sum_{\ell \in \mathbb{Z}} \frac{\operatorname{sgn}(\ell - c) + \operatorname{sgn}(z_2)}{2} \mathbf{e}(\ell z) = \mathcal{R}_c(\mathbf{e}(z)) \end{aligned}$$

if  $z_2 \neq 0$ , and the formula remains true for  $z_2 = 0$  by continuity. An alternative proof can be obtained by noting that  $\mathbf{e}(-cz)\mathcal{R}_c(\mathbf{e}(z))$  is periodic of period 1 with respect to  $c$  and expanding it as a Fourier series in  $c$ , again by the Poisson summation formula.

For  $m \in \mathbb{N}$  and  $s = (\alpha, \beta) \in \mathbb{Q}^2$  we now define a *universal Appell-Lerch sum*  $\mathcal{A}_m^s(\tau, z)$  by

$$\mathcal{A}_m^s(\tau, z) = \mathbf{e}(-m\alpha z_s) \operatorname{Av}^{(m)} \left[ \mathcal{R}_{-2m\alpha}(y/y_s) \right] \quad (y_s = \mathbf{e}(z_s) = \mathbf{e}(\beta)q^\alpha). \quad (8.17)$$

It is easy to check that this function satisfies

$$\mathcal{A}_m^{(\alpha+\lambda, \beta+\mu)} = \mathbf{e}(-m(\mu\alpha - \lambda\beta)) \mathcal{A}_m^{(\alpha, \beta)} \quad (\lambda, \mu \in \mathbb{Z}) \quad (8.18)$$

and hence, in view of the corresponding property (8.7) of  $D_s$ , that the product  $D_s(\tau)\mathcal{A}_m^s(\tau, z)$  depends only on the class of  $s$  in  $S/\mathbb{Z}^2$ . We can therefore define

$$\varphi^P(\tau, z) := \sum_{s \in S/\mathbb{Z}^2} D_s(\tau) \mathcal{A}_m^s(\tau, z), \quad (8.19)$$

and it is obvious from the above discussion that this function, which we will call the **polar part** of  $\varphi$ , satisfies the index  $m$  elliptic transformation property (4.2) and has the same poles and residues as  $\varphi$ , so that the difference  $\varphi - \varphi^P$  is holomorphic and has a theta expansion. In fact, we have:

**Theorem 8.1.** *Let  $\varphi(\tau, z)$  be a meromorphic Jacobi form with simple poles at  $z = z_s = \alpha\tau + \beta$  for  $s = (\alpha, \beta) \in S \subset \mathbb{Q}^2$ , with Fourier coefficients  $h_\ell(\tau)$  defined by (8.1) and (8.2) or by (8.4) and residues  $D_s(\tau)$  defined by (8.6). Then  $\varphi$  has the decomposition*

$$\varphi(\tau, z) = \varphi^F(\tau, z) + \varphi^P(\tau, z), \quad (8.20)$$

where  $\varphi^F$  and  $\varphi^P$  are defined by equations (8.3) and (8.19), respectively.

*Proof:* Fix a point  $P = A\tau + B \in \mathbb{C}$  with  $(A, B) \in \mathbb{R}^2 \setminus S$ . Since  $\varphi$ ,  $\varphi^F$  and  $\varphi^P$  are meromorphic, it suffices to prove the decomposition (8.20) on the horizontal line  $\text{Im}(z) = \text{Im}(P) = A\tau_2$ . On this line we have the Fourier expansion

$$\varphi(\tau, z) = \sum_{\ell \in \mathbb{Z}} q^{\ell^2/4m} h_\ell^{(P)}(\tau) y^\ell,$$

where the coefficients  $h_\ell^{(P)}$  are defined by (8.1) (modified as explained in the text there if  $A = \alpha$  for any  $(\alpha, \beta) \in S$ , but for simplicity we will simply assume that this is not the case, since we are free to choose  $A$  any way we want). Comparing this with (8.3) gives

$$\varphi(\tau, z) - \varphi^F(\tau, z) = \sum_{\ell \in \mathbb{Z}} (h_\ell^{(P)}(\tau) - h_\ell(\tau)) q^{\ell^2/4m} y^\ell \quad (\text{Im}(z) = \text{Im}(P)). \quad (8.21)$$

But  $q^{\ell^2/4m}(h_\ell^{(P)}(\tau) - h_\ell(\tau))$  is just  $2\pi i$  times the sum of the residues of  $\varphi(\tau, z)\mathbf{e}(-\ell z)$  in the parallelogram with width 1 and horizontal sides at heights  $A\tau_2$  and  $-\ell\tau_2/2m$ , with the residues of any poles on the latter line being counted with multiplicity 1/2 because of the way we defined  $h_\ell$  in that case, so

$$\begin{aligned} q^{\ell^2/4m}(h_\ell^{(P)}(\tau) - h_\ell(\tau)) &= 2\pi i \sum_{s=(\alpha, \beta) \in S/\mathbb{Z}} \frac{\text{sgn}(\alpha - A) - \text{sgn}(\alpha + \ell/2m)}{2} \text{Res}_{z=z_s}(\varphi(\tau, z)\mathbf{e}(-\ell z)) \\ &= \sum_{s=(\alpha, \beta) \in S/\mathbb{Z}} \frac{\text{sgn}(\alpha - A) - \text{sgn}(\ell + 2m\alpha)}{2} D_s(\tau) \mathbf{e}(-(\ell + m\alpha)z_s). \end{aligned}$$

(Here “ $(\alpha, \beta) \in S/\mathbb{Z}$ ” means that we consider all  $\alpha$ , but  $\beta$  only modulo 1, which is the same by periodicity as considering only the  $(\alpha, \beta)$  with  $B \leq \beta < B + 1$ .) Inserting this formula into (8.21) and using (8.15), we find

$$\begin{aligned}
\varphi(\tau, z) - \varphi^F(\tau, z) &= - \sum_{s=(\alpha, \beta) \in S/\mathbb{Z}} \mathbf{e}(-m\alpha z_s) D_s(\tau) \sum_{\ell \in \mathbb{Z}} \frac{\operatorname{sgn}(\operatorname{Im}(z - z_s)) + \operatorname{sgn}(\ell + 2m\alpha)}{2} \left(\frac{y}{y_s}\right)^\ell \\
&= \sum_{s=(\alpha, \beta) \in S/\mathbb{Z}} \mathbf{e}(-m\alpha z_s) D_s(\tau) \mathcal{R}_{-2m\alpha}(y/y_s) \\
&= \sum_{s=(\alpha, \beta) \in S/\mathbb{Z}^2} \sum_{\lambda \in \mathbb{Z}} \mathbf{e}(-m(\alpha - \lambda)(z_s - \lambda\tau)) D_{(\alpha - \lambda, \beta)}(\tau) \mathcal{R}_{-2m(\alpha - \lambda)}(q^\lambda y/y_s) \\
&= \sum_{s=(\alpha, \beta) \in S/\mathbb{Z}^2} D_s(\tau) \mathbf{e}(-m\alpha z_s) \sum_{\lambda \in \mathbb{Z}} q^{m\lambda^2} y^{2m\lambda} \mathcal{R}_{-2m\alpha}(q^\lambda y/y_s),
\end{aligned}$$

where in the last line we have used the periodicity property (8.7) of  $D_s(\tau)$  together with the obvious periodicity property  $\mathcal{R}_{c+n}(y) = y^n \mathcal{R}_c(y)$  of  $\mathcal{R}_c(y)$ . But the inner sum in the last expression is just  $\operatorname{Av}^{(m)}[\mathcal{R}_{-2m\alpha}(y/y_s)]$ , so from the definition (8.17) we see that this agrees with  $\varphi^P(\tau, z)$ , as claimed.

### 8.3 Mock modularity of the Fourier coefficients

In subsections §8.1 and §8.2 we introduced a canonical splitting of a meromorphic Jacobi form  $\varphi$  into a finite part  $\varphi^F$  and a polar part  $\varphi^P$ , but there is no reason yet (apart from the simplicity of equation (8.2)) to believe that the choice we have made is the “right” one: we could have defined periodic Fourier coefficients  $h_\ell(\tau)$  in many other ways (for instance, by taking  $P = P_0 - \ell/2m\tau$  with any fixed  $P_0 \in \mathbb{C}$  or more generally  $P = P_\ell - \ell\tau/2m$  where  $P_\ell$  depends only on  $\ell$  modulo  $2m$ ) and obtained other functions  $\varphi^F$  and  $\varphi^P$ . What makes the chosen decomposition special is that, as we will now show, the Fourier coefficients defined in (8.2) are (mixed) mock modular forms and the function  $\varphi^F$  therefore a (mixed) mock Jacobi form in the sense of §7.3. The corresponding shadows will involve theta series that we now introduce.

For  $m \in \mathbb{N}$ ,  $\ell \in \mathbb{Z}/2m\mathbb{Z}$  and  $s = (\alpha, \beta) \in \mathbb{Q}^2$  we define the unary theta series

$$\Theta_{m, \ell}^s(\tau) = \mathbf{e}(-m\alpha\beta) \sum_{\lambda \in \mathbb{Z} + \alpha + \ell/2m} \lambda \mathbf{e}(2m\beta\lambda) q^{m\lambda^2} \tag{8.22}$$

of weight  $3/2$  and its Eichler integral<sup>16</sup>

$$\Theta_{m, \ell}^{s*}(\tau) = \frac{\mathbf{e}(m\alpha\beta)}{2} \sum_{\lambda \in \mathbb{Z} + \alpha + \ell/2m} \operatorname{sgn}(\lambda) \mathbf{e}(-2m\beta\lambda) \operatorname{erfc}(2|\lambda|\sqrt{\pi m\tau_2}) q^{-m\lambda^2} \tag{8.23}$$

---

<sup>16</sup>Strictly speaking, the Eichler integral as defined by equation (7.2) with  $k = 1/2$  would be this multiplied by  $2\sqrt{\pi/m}$ , but this normalization will lead to simpler formulas and, as already mentioned, there is no good universal normalization for the shadows of mock modular forms.

(cf. (7.2) and (7.6)). One checks easily that these functions transform by

$$\Theta_{m,\ell}^{(\alpha+\lambda,\beta+\mu)}(\tau) = \mathbf{e}(m(\mu\alpha - \lambda\beta)) \Theta_{m,\ell}^{(\alpha,\beta)}(\tau) \quad (\lambda, \mu \in \mathbb{Z}), \quad (8.24)$$

$$\Theta_{m,\ell}^{(\alpha+\lambda,\beta+\mu)*}(\tau) = \mathbf{e}(-m(\mu\alpha - \lambda\beta)) \Theta_{m,\ell}^{(\alpha,\beta)*}(\tau) \quad (\lambda, \mu \in \mathbb{Z}). \quad (8.25)$$

with respect to translations of  $s$  by elements of  $\mathbb{Z}^2$ . From this and (8.7) it follows that the products  $D_s \overline{\Theta_{m,\ell}^s}$  and  $D_s \Theta_{m,\ell}^{s*}$  depend only on the class of  $s$  in  $S/\mathbb{Z}^2$ , so that the sums over  $s$  occurring in the following theorem make sense.

**Theorem 8.2.** *Let  $\varphi$ ,  $h_\ell$  and  $\varphi^F$  be as in Theorem 8.1. Then each  $h_\ell$  is a mixed mock modular form of weight  $k - \frac{1}{2} \mid k - 1$ , with shadow  $\sum_{s \in S/\mathbb{Z}^2} D_s(\tau) \overline{\Theta_{m,\ell}^s(\tau)}$ , and the function  $\varphi^F$  is a mixed mock Jacobi form. More precisely, for each  $\ell \in \mathbb{Z}/2m\mathbb{Z}$  the completion of  $h_\ell$  defined by*

$$\widehat{h}_\ell(\tau) := h_\ell(\tau) - \sum_{s \in S/\mathbb{Z}^2} D_s(\tau) \Theta_{m,\ell}^{s*}(\tau), \quad (8.26)$$

with  $\Theta_{m,\ell}^{s*}$  as in (8.23), transforms like a modular form of weight  $k - 1/2$  with respect to some congruence subgroup of  $SL(2, \mathbb{Z})$ , and the completion of  $\varphi^F$  defined by

$$\widehat{\varphi}^F(\tau, z) := \sum_{\ell \pmod{2m}} \widehat{h}_\ell(\tau) \vartheta_{m,\ell}(\tau, z) \quad (8.27)$$

transforms like a Jacobi form of weight  $k$  and index  $m$  with respect to the full modular group.

The key property needed to prove this theorem is the following proposition, essentially due to Zwegers, which says that the functions  $\mathcal{A}_m^s(\tau, z)$  defined in §8.2 are (meromorphic) mock Jacobi forms of weight 1 and index  $m$ , with shadow  $\sum_{\ell \pmod{2m}} \overline{\Theta_{m,\ell}^s(\tau)} \vartheta_{m,\ell}(\tau, z)$  (more precisely, that each  $\mathcal{A}_m^s$  is a meromorphic mock Jacobi form of this weight, index and shadow with respect to some congruence subgroup of  $SL(2, \mathbb{Z})$  depending on  $s$  and that the collection of all of them is a vector-valued mock Jacobi form on the full modular group):

**Proposition 8.2.** *For  $m \in \mathbb{N}$  and  $s \in \mathbb{Q}^2$  the completion  $\widehat{\mathcal{A}}_m^s$  of  $\mathcal{A}_m^s$  defined by*

$$\widehat{\mathcal{A}}_m^s(\tau, z) := \mathcal{A}_m^s(\tau, z) + \sum_{\ell \pmod{2m}} \Theta_{m,\ell}^{s*}(\tau) \vartheta_{m,\ell}(\tau, z). \quad (8.28)$$

satisfies

$$\widehat{\mathcal{A}}_m^{(\alpha+\lambda,\beta+\mu)}(\tau, z) = \mathbf{e}(-m(\mu\alpha - \lambda\beta)) \widehat{\mathcal{A}}_m^{(\alpha,\beta)}(\tau) \quad (\lambda, \mu \in \mathbb{Z}), \quad (8.29)$$

$$\widehat{\mathcal{A}}_m^s(\tau, z + \lambda\tau + \mu) = \mathbf{e}(-m(\lambda^2\tau + 2\lambda z)) \widehat{\mathcal{A}}_m^s(\tau) \quad (\lambda, \mu \in \mathbb{Z}), \quad (8.30)$$

$$\widehat{\mathcal{A}}_m^s\left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}\right) = (c\tau + d) \mathbf{e}\left(\frac{mcz^2}{c\tau + d}\right) \widehat{\mathcal{A}}_m^{s\gamma}(\tau, z) \quad \left(\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})\right). \quad (8.31)$$

*Proof:* The first two properties are easy to check because they hold for each term in (8.28) separately. The modular transformation property, considerably less obvious, is essentially the content of Proposition 3.5 of [119], but the functions he studies are different from ours and we need a small calculation to relate them. Zwegers defines two functions  $f_u^{(m)}(z; \tau)$  and  $\tilde{f}_u^{(m)}(z; \tau)$  ( $m \in \mathbb{N}$ ,  $\tau \in \mathbb{H}$ ,  $z, u \in \mathbb{C}$ ) by

$$f_u^{(m)}(z; \tau) = \text{Av}^{(m)} \left[ \frac{1}{1 - y/\mathbf{e}(u)} \right], \quad \tilde{f}_u^{(m)}(z; \tau) = f_u^{(m)}(z; \tau) - \frac{1}{2} \sum_{\ell \pmod{2m}} R_{m,\ell}(u; \tau) \vartheta_{m,\ell}(\tau, z)$$

(here we have rewritten Definition 3.2 of [119] in our notation), where

$$R_{m,\ell}(u; \tau) = \sum_{r \in \ell + 2m\mathbb{Z}} \left\{ \text{sgn}\left(r + \frac{1}{2}\right) - \text{erf}\left(\sqrt{\pi} \frac{r\tau_2 + 2mu_2}{\sqrt{m\tau_2}}\right) \right\} q^{-r^2/4m} \mathbf{e}(-ru), \quad (8.32)$$

and shows (Proposition 3.5) that  $\tilde{f}_u^{(m)}$  satisfies the modular transformation property

$$\tilde{f}_{u/(c\tau+d)}^{(m)}\left(\frac{z}{c\tau+d}; \frac{a\tau+b}{c\tau+d}\right) = (c\tau+d) \mathbf{e}\left(\frac{mc(z^2-u^2)}{c\tau+d}\right) \tilde{f}_u^{(m)}(z; \tau) \quad (8.33)$$

for all  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ . Noting that  $\text{erf}(x) = \text{sgn}(x)(1 - \text{erfc}(|x|))$ , we find that

$$\frac{1}{2} R_{m,\ell}(z_s; \tau) = \sum_{r \equiv \ell \pmod{2m}} \frac{\text{sgn}(r + \frac{1}{2}) - \text{sgn}(r + 2m\alpha)}{2} q^{-r^2/4m} y_s^{-r} + \mathbf{e}(m\alpha z_s) \Theta_{m,\ell}^{s*}(\tau)$$

in our notation. On the other hand, from (8.13) we have

$$\mathcal{R}_{-2m\alpha}(y) = \frac{1}{y-1} + \sum_{r \in \mathbb{Z}} \frac{\text{sgn}(r + \frac{1}{2}) - \text{sgn}(r + 2m\alpha)}{2} y^r \quad (8.34)$$

(note that the sum here is finite). Replacing  $y$  by  $y/y_s$  and applying the operator  $\text{Av}^{(m)}$ , we find (using (8.10))

$$\mathbf{e}(m\alpha z_s) \mathcal{A}_m^s(\tau, z) = -f_{z_s}^{(m)}(z; \tau) + \sum_{r \in \mathbb{Z}} \frac{\text{sgn}(r + \frac{1}{2}) - \text{sgn}(r + 2m\alpha)}{2} q^{-r^2/4m} y_s^{-r} \vartheta_{m,r}(\tau, z).$$

Combining these two equations and rearranging, we obtain

$$\widehat{\mathcal{A}}_m^s(\tau, z) = -\mathbf{e}(-m\alpha z_s) \tilde{f}_{z_s}^{(m)}(z; \tau),$$

and the modularity property (8.31) then follows from (8.33) after a short calculation.



The proof of Theorem 8.2 follows easily from Proposition 8.2. We define the completion of the function  $\varphi^P$  studied in §8.2 by

$$\widehat{\varphi}^P(\tau, z) := \sum_{s \in S/\mathbb{Z}^2} D_s(\tau) \widehat{\mathcal{A}}_m^s(\tau, z). \quad (8.35)$$

The sum makes sense by (8.29), and from the transformation equations (8.30)–(8.31) together with the corresponding properties (8.7)–(8.8) of the residue functions  $D_s(\tau)$  it follows that  $\widehat{\varphi}^P(\tau, z)$  transforms like a Jacobi form of weight  $k$  and index  $m$  with respect to the full modular group. Comparing equations (8.35) and (8.28) with equations (8.27) and (8.26), we find that

$$\widehat{\varphi}^P(\tau, z) - \varphi^P(\tau, z) = \sum_{\ell \in \mathbb{Z}/2m\mathbb{Z}} \sum_{s \in S/\mathbb{Z}^2} D_s(\tau) \Theta_{m,\ell}^{s*}(\tau) \vartheta_{m,\ell}(\tau, z) = \varphi^F(\tau, z) - \widehat{\varphi}^F(\tau, z)$$

and hence, using Theorem 8.1, that

$$\widehat{\varphi}^F(\tau, z) + \widehat{\varphi}^P(\tau, z) = \varphi^F(\tau, z) + \varphi^P(\tau, z) = \varphi(\tau, z).$$

Since both  $\varphi(\tau, z)$  and  $\widehat{\varphi}^P(\tau, z)$  transform like Jacobi forms of weight  $k$  and index  $m$ , it follows that  $\widehat{\varphi}^F(\tau, z)$  also does, and then the fact that each  $\widehat{h}_\ell$  transforms like a modular form of weight  $k-1/2$  (and hence that each  $h_\ell$  is a mixed mock modular form with the weight and shadow given in the theorem) follows by the same argument that proves the modularity of the coefficients  $h_\ell$  in the theta expansion (4.10) in the classical case.

**Summary.** For the reader's convenience, we give a brief summary of the results given up to now. We have the following six functions of two variables  $(\tau, z) \in \mathbb{H} \times \mathbb{C}$ :

- $\varphi(\tau, z)$ , a meromorphic Jacobi form of weight  $k$  and index  $m$ , assumed to have only simple poles at  $z = z_s = \alpha\tau + \beta$  for  $s = (\alpha, \beta)$  in some discrete subset  $S \subset \mathbb{Q}^2$ ;
- $\varphi^F(\tau, z)$ , the **finite** part of  $\varphi$ , defined by the theta expansion  $\sum_{\ell \pmod{2m}} h_\ell(\tau) \vartheta_{m,\ell}(\tau, z)$  where  $h_\ell(\tau)$  is  $q^{\ell^2/4m}$  times the  $\ell$ th Fourier coefficient of  $\varphi(\tau, z - \ell\tau/2m)$  on the real line;
- $\varphi^P(\tau, z)$ , the **polar** part of  $\varphi$ , defined as  $\sum_{s \in S/\mathbb{Z}^2} D_s(\tau) \mathcal{A}_m^s(\tau, z)$ , where  $\mathcal{A}_m^s$  is an explicit Appell-Lerch sum having simple poles at  $z \in z_s + \mathbb{Z}\tau + \mathbb{Z}$ ;
- $\varphi^C(\tau, z)$ , a non-holomorphic **correction** term, defined as  $\sum_s \sum_\ell D_s(\tau) \Theta_{m,\ell}^{s*}(\tau) \vartheta_{m,\ell}(\tau, z)$  where the  $\Theta_{m,\ell}^{s*}$  are the Eichler integrals of explicit unary theta series of weight  $3/2$ ;
- $\widehat{\varphi}^F(\tau, z)$ , the completed finite part, defined as  $\varphi^F(\tau, z) - \varphi^C(\tau, z)$ ;
- $\widehat{\varphi}^P(\tau, z)$ , the completed polar part, defined as  $\varphi^P(\tau, z) + \varphi^C(\tau, z)$ .

These six functions are related by

$$\varphi^F + \varphi^P = \varphi = \widehat{\varphi}^F + \widehat{\varphi}^P, \quad \varphi^F - \widehat{\varphi}^F = \varphi^C = \widehat{\varphi}^P - \varphi^P. \quad (8.36)$$

Each of them is real-analytic in  $\tau$ , meromorphic in  $z$ , satisfies the elliptic transformation property (4.2) with respect to  $z$ , and has precisely two of four further desirable properties of such a function (note that  $6 = \binom{4}{2}$ ), as shown in the following table

	$\varphi$	$\varphi^F$	$\varphi^P$	$\varphi^C$	$\widehat{\varphi}^F$	$\widehat{\varphi}^P$
holomorphic in $\tau$ ?	✓	✓	✓	–	–	–
transforms like a Jacobi form?	✓	–	–	–	✓	✓
holomorphic in $z$ ?	–	✓	–	✓	✓	–
determined by the poles of $\varphi$ ?	–	–	✓	✓	–	✓

in which the three checked entries in each row correspond to one of the equations (10.1). Each Fourier coefficient  $h_\ell$  of  $\varphi$  is a mixed mock modular form of weight  $(k-1, 1/2)$ , and the finite part  $\varphi^F$  is a mixed mock Jacobi form. In the holomorphic case, the functions  $\varphi$ ,  $\varphi^F$  and  $\widehat{\varphi}^F$  coincide and the functions  $\varphi^P$ ,  $\varphi^C$  and  $\widehat{\varphi}^P$  vanish.

We end by mentioning one further property of the canonical decomposition  $\varphi = \varphi^F + \varphi^P$  that seems of interest. The finite part  $\varphi^P$  of  $\varphi$  is a linear combination of terms  $h_\ell(\tau) \vartheta_{m,\ell}(\tau, z)$  where  $h_\ell(\tau)$  is a Laurent power series in  $q$  and  $\vartheta_{m,\ell}(\tau, z)$  a linear combination of terms  $q^{r^2/4m} y^r$ , so it satisfies the same growth condition as a weak Jacobi form, viz., that it contains only monomials  $q^n y^r$  with discriminant  $4mn - r^2$  bounded from below. But the Appell-Lerch sums  $\mathcal{A}_m^s(\tau, z)$  occurring in  $\varphi^P(\tau, z)$  have the opposite property: substituting the Fourier expansion (8.15) of  $\mathcal{R}_c$  into the definition (8.17), we find that they contain only terms  $q^n y^r$  with  $4mn - r^2 = -\delta^2$  where the numbers  $\delta$  occurring are rational (more precisely,  $\delta \in r + 2m(\mathbb{Z} + \alpha)$ ) and unbounded (more precisely,  $\delta$  lies between 0 and  $r + O(1)$ ).

#### 8.4 The case of double poles

In this subsection we extend our considerations to the case when  $\varphi$  is allowed to have double poles, again assumed to be at points  $z = z_s = \alpha\tau + \beta$  for  $s = (\alpha, \beta)$  belonging to some discrete subset  $S$  of  $\mathbb{Q}^2$ . The first thing we need to do is to generalize the definition (8.6) to this case. For  $s \in S$  we define functions  $E_s$  and  $D_s$  on  $\mathbb{H}$  by the Laurent expansion

$$e(m\alpha z_s) \varphi(\tau, z_s + \varepsilon) = \frac{E_s(\tau)}{(2\pi i \varepsilon)^2} + \frac{D_s(\tau) - 2m\alpha E_s(\tau)}{2\pi i \varepsilon} + O(1) \quad \text{as } \varepsilon \rightarrow 0. \quad (8.37)$$

(Notice that  $D_s(\tau)$  is the same function as in (8.6) when the pole is simple.) It is easily checked that the behavior of these functions under translations of  $s$  by elements of  $\mathbb{Z}^2$  is given by equation (8.7) and its analogue for  $E_s$ . For the modular behavior, we have:

**Proposition 8.3.** *The functions  $E_s(\tau)$  and  $D_s(\tau)$  defined by (8.37) are modular forms of weight  $k-2$  and  $k-1$ , respectively. More respectively, for all  $s \in S$  and  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$  we have*

$$E_s\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^{k-2} E_{s\gamma}(\tau), \quad D_s\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^{k-1} D_{s\gamma}(\tau). \quad (8.38)$$

*Proof:* We rewrite (8.37) as

$$\mathbf{e}(m\alpha(z_s + 2\varepsilon)) \varphi(\tau, z_s + \varepsilon) = \frac{E_s(\tau)}{(2\pi i\varepsilon)^2} + \frac{D_s(\tau)}{2\pi i\varepsilon} + \mathbf{O}(1),$$

and also write  $\alpha_s$  and  $z_s(\tau)$  instead of just  $\alpha$  and  $z_s$ . Then using the easily checked identities

$$z_s(\gamma\tau) = \frac{z_{s\gamma}(\tau)}{c\tau + d}, \quad \alpha_{s\gamma} z_{s\gamma}(\tau) - \alpha_s z_s(\gamma\tau) = \frac{c z_{s\gamma}(\tau)^2}{c\tau + d} \quad \left( \gamma\tau := \frac{a\tau + b}{c\tau + d} \right),$$

and the modular transformation equation (4.1), we find

$$\begin{aligned} \frac{(c\tau + d)^2 E_s(\gamma\tau)}{(2\pi i\varepsilon)^2} + \frac{(c\tau + d) D_s(\gamma\tau)}{2\pi i\varepsilon} &\equiv \mathbf{e}\left(m\alpha\left(z_s(\gamma\tau) + \frac{2\varepsilon}{c\tau + d}\right)\right) \varphi\left(\gamma\tau, z_s(\gamma\tau) + \frac{\varepsilon}{c\tau + d}\right) \\ &\equiv \mathbf{e}\left(m\alpha \frac{z_{s\gamma}(\tau) + 2\varepsilon}{c\tau + d}\right) \varphi\left(\gamma\tau, \frac{z_{s\gamma}(\tau) + \varepsilon}{c\tau + d}\right) \\ &\equiv (c\tau + d)^k \mathbf{e}\left(m\alpha \frac{z_{s\gamma} + 2\varepsilon}{c\tau + d} + \frac{mc(z_{s\gamma} + \varepsilon)^2}{c\tau + d}\right) \varphi\left(\tau, z_{s\gamma} + \varepsilon\right) \\ &\equiv (c\tau + d)^k \mathbf{e}(m\alpha_{s\gamma}(z_{s\gamma} + 2\varepsilon)) \varphi(\tau, z_{s\gamma} + \varepsilon) \\ &\equiv (c\tau + d)^k \left[ \frac{E_{s\gamma}(\tau)}{(2\pi i\varepsilon)^2} + \frac{D_{s\gamma}(\tau)}{2\pi i\varepsilon} \right], \end{aligned}$$

where “ $\equiv$ ” means equality modulo a quantity that is bounded as  $\varepsilon \rightarrow 0$ . The claim follows.

Next, we must define a standard Appell-Lerch sum with a double pole at  $z = z_s$ . We begin by defining a rational function  $\mathcal{R}_c^{(2)}(y)$  with a double pole at  $y = 1$  for each  $c \in \mathbb{R}$ . Motivated by Proposition 8.1, we require

$$\mathcal{R}_c^{(2)}(\mathbf{e}(z)) = \text{Av}_{\mathbb{Z}} \left[ \frac{\mathbf{e}(cz)}{(2\pi iz)^2} \right] = \frac{1}{(2\pi i)^2} \sum_{n \in \mathbb{Z}} \frac{\mathbf{e}(c(z-n))}{(z-n)^2}. \quad (8.39)$$

To calculate this explicitly as a rational function, we could imitate the proof of Proposition 8.1, but it is easier to note that  $\mathcal{R}_c^{(2)} = \left(-\frac{1}{2\pi i} \frac{d}{dz} + c\right) \mathcal{R}_c = \left(-y \frac{d}{dy} + c\right) \mathcal{R}_c$  and hence from equations (8.15), (8.14) and (8.13) we get three alternative definitions

$$\mathcal{R}_c^{(2)}(y) = \sum_{\ell \in \mathbb{Z}} \frac{|\ell - c| + \text{sgn}(z_2)(\ell - c)}{2} y^\ell \quad (8.40)$$

$$= \begin{cases} \sum_{\ell \geq c} (\ell - c) y^\ell & \text{if } |y| < 1 \\ \sum_{\ell \leq c} (c - \ell) y^\ell & \text{if } |y| > 1 \end{cases} \quad (8.41)$$

$$= y^{\lfloor c \rfloor + 1} \left( \frac{1}{(y-1)^2} + \frac{c - \lfloor c \rfloor}{y-1} \right). \quad (8.42)$$

of  $\mathcal{R}_c^{(2)}(y)$ . (Notice that in these equations neither the asterisk on the summation sign nor the case distinction for  $c \in \mathbb{Z}$  and  $c \notin \mathbb{Z}$  are needed as before, and that the function  $\mathcal{R}_c^{(2)}(y)$ , unlike  $\mathcal{R}_c(y)$ , is continuous in  $c$ .) For  $s = (\alpha, \beta) \in \mathbb{Q}^2$  and  $m \in \mathbb{N}$  we set

$$\mathcal{B}_m^s(\tau, z) = \mathbf{e}(-m\alpha z_s) \text{Av}^{(m)}[\mathcal{R}_{-2m\alpha}^{(2)}(y/y_s)], \quad (8.43)$$

in analogy with (8.17). If we then define the polar part  $\varphi^P$  of  $\varphi$  by

$$\varphi^P(\tau, z) = \sum_{s \in S/\mathbb{Z}^2} \left( D_s(\tau) \mathcal{A}_m^s(\tau, z) + E_s(\tau) \mathcal{B}_m^s(\tau, z) \right); \quad (8.44)$$

then the definitions of the functions  $D_s$ ,  $E_s$ ,  $\mathcal{A}_m^s$  and  $\mathcal{B}_m^s$  immediately imply that  $\varphi^P$  has the same singularities as  $\varphi$ , so that the difference

$$\varphi^F(\tau, z) = \varphi(\tau, z) - \varphi^P(\tau, z) \quad (8.45)$$

is a holomorphic function of  $z$ .

As before, the key property of the Appell-Lerch sums is that they are again mock Jacobi forms, of a somewhat more complicated type than before. We introduce the unary theta series

$$\vartheta_{m,\ell}^s(\tau) = \mathbf{e}(-m\alpha\beta) \sum_{\lambda \in \mathbb{Z} + \alpha + \ell/2m} \mathbf{e}(2m\beta\lambda) q^{m\lambda^2} \quad (8.46)$$

of weight  $1/2$  and its (again slightly renormalized) Eichler integral

$$\vartheta_{m,\ell}^{s*}(\tau) = \frac{\overline{\vartheta_{m,\ell}^s(\tau)}}{2\pi\sqrt{m\tau_2}} - \mathbf{e}(m\alpha\beta) \sum_{\lambda \in \mathbb{Z} + \alpha + \ell/2m} |\lambda| \mathbf{e}(-2m\beta\lambda) \text{erfc}(2|\lambda|\sqrt{\pi m\tau_2}) q^{-m\lambda^2} \quad (8.47)$$

(cf. (7.2) and (7.6)). Then we can define the completion  $\widehat{\mathcal{B}}_m^s$  of  $\mathcal{B}_m^s$  by

$$\widehat{\mathcal{B}}_m^s(\tau, z) := \mathcal{B}_m^s(\tau, z) + m \sum_{\ell \pmod{2m}} \vartheta_{m,\ell}^{s*}(\tau) \vartheta_{m,\ell}(\tau, z). \quad (8.48)$$

**Proposition 8.4.** *For  $m \in \mathbb{N}$  and  $s \in \mathbb{Q}^2$  the completion  $\widehat{\mathcal{B}}_m^s$  of  $\mathcal{B}_m^s$  defined by (8.48) satisfies*

$$\widehat{\mathcal{B}}_m^{(\alpha+\lambda, \beta+\mu)}(\tau, z) = \mathbf{e}(-m(\mu\alpha - \lambda\beta + \lambda\mu)) \widehat{\mathcal{B}}_m^{(\alpha, \beta)}(\tau) \quad (\lambda, \mu \in \mathbb{Z}), \quad (8.49)$$

$$\widehat{\mathcal{B}}_m^s(\tau, z + \lambda\tau + \mu) = \mathbf{e}(-m(\lambda^2\tau + 2\lambda z)) \widehat{\mathcal{B}}_m^s(\tau) \quad (\lambda, \mu \in \mathbb{Z}), \quad (8.50)$$

$$\widehat{\mathcal{B}}_m^s\left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}\right) = (c\tau + d)^2 \mathbf{e}\left(\frac{mcz^2}{c\tau + d}\right) \widehat{\mathcal{B}}_m^{s\gamma}(\tau, z) \quad \left(\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})\right). \quad (8.51)$$

The proof is exactly similar to that of 8.2. We define functions  $g_u^{(m)}(z; \tau)$  and  $\tilde{g}_u^{(m)}(z; \tau)$  by applying the operator  $\frac{1}{2\pi i} \frac{\partial}{\partial u} - 2m \frac{u_2}{\tau_2}$  to  $f_u^{(m)}(z; \tau)$  and  $\tilde{f}_u^{(m)}(z; \tau)$ ; then the transformation equation (8.33) of  $\tilde{f}_u^{(m)}$  implies the same transformation equation for  $\tilde{g}_u^{(m)}$  but with the initial factor  $(c\tau + d)$  replaced by  $(c\tau + d)^2$ , and a calculation exactly similar to the one given before shows that  $\mathbf{e}(m\alpha z_s) \mathcal{B}_m^s(\tau, z)$  differs from  $g_{z_s}^{(m)}(z; \tau)$  by a finite linear combination of functions  $\vartheta_{m,r}(\tau, z)$  and that  $\mathbf{e}(m\alpha z_s) \widehat{\mathcal{B}}_m^s(\tau, z) = \tilde{g}_{z_s}^{(m)}(z; \tau)$ . We omit the details.

**Theorem 8.3.** *Let  $\varphi$  be as above, with singularities at  $z = z_s$  ( $s \in S \subset \mathbb{Q}^2$ ) given by (8.37). Then the finite part  $\varphi^F$  as defined by (8.45) coincides with the finite part defined by the theta expansion (8.3), the coefficients  $h_\ell(\tau)$  in this expansion are mixed mock modular forms, with completion given by*

$$\widehat{h}_\ell(\tau) = h_\ell(\tau) + \sum_{s \in S/\mathbb{Z}^2} \left( D_s(\tau) \Theta_{m,\ell}^{s*}(\tau) + E_s(\tau) \vartheta_{m,\ell}^{s*}(\tau) \right),$$

and the completion  $\widehat{\varphi}^F$  defined by (8.27) transforms like a Jacobi form of weight  $k$  and index  $m$ .

The proof follows the same lines as before: the equivalence of (8.3) and (8.45) is proved by expanding  $\varphi(\tau, z)$  as a Fourier series along the horizontal line  $\text{Im}(z) = \text{Im}(P)$  for some generic point  $P \in \mathbb{C}$  and calculating the difference  $\varphi - \varphi^F$  as a sum of residues, and the mock modularity is proved by decomposing  $\varphi$  as  $\widehat{\varphi}^F + \widehat{\varphi}^P$  with  $\widehat{\varphi}^F$  as in (8.27) and  $\widehat{\varphi}^P = \sum_{s \in S/\mathbb{Z}^2} (D_s \widehat{\mathcal{A}}_m^s + E_s \widehat{\mathcal{B}}_m^s)$ , which transforms like a Jacobi form by virtue of Proposition 8.4. Again the details are left to the reader. Note that here the mock modular forms  $h_\ell(\tau)$  are of the “even more mixed” variety mentioned at the end of §7.2, since they now have a shadow that is a linear combination of two terms  $\sum_s D_s \overline{\Theta_{m,\ell}^s}$  and  $\sum_s E_s \overline{\vartheta_{m,\ell}^s}$  belonging to two different tensor products  $M_{k-1} \otimes \overline{M_{3/2}}$  and  $M_{k-2} \otimes \overline{M_{1/2}}$  and hence two different mixed weights  $k - \frac{1}{2} | k - 1$  and  $k - \frac{1}{2} | k - 2$  rather than a single mixed weight  $k - \frac{1}{2} | k - 1$  as in the case of simple poles.

## 8.5 Examples

We end this section by giving five examples of meromorphic Jacobi forms and their canonical decompositions into a mock Jacobi form and a finite linear combination of Appell-Lerch sums. We systematically use the notations  $A = \varphi_{-2,1}$ ,  $B = \varphi_{0,1}$ ,  $C = \varphi_{-1,2}$  for the three basic generators of the ring of weak Jacobi forms as described in equations (4.26)–(4.35).

**Example 1: Weight 1, index 1, simple pole at  $z = 0$ .** As our first example we take the Jacobi form  $\varphi = C/A \in J_{1,1}^{\text{mer}}$ , which has a simple pole of residue  $1/\pi i$  at  $z = 0$  and a Fourier expansion beginning

$$\frac{y+1}{y-1} - (y^2 - y^{-2})q - 2(y^3 - y^{-3})q^2 - 2(y^4 - y^{-4})q^3 - (2y^5 + y^4 - y^{-4} - 2y^{-5})q^4 - \dots$$

Calculating the Fourier expansion of the polar part  $\varphi^P = \text{Av}^{(1)}\left[\frac{y+1}{y-1}\right]$ , we find that it begins the same way, and indeed, we must have  $\varphi = \varphi^P$  because the Fourier coefficients  $h_\ell$  all vanish (we have  $h_{-\ell} = -h_\ell$  because the weight is odd and  $h_{\ell+2} = h_\ell$  because the index is 1). So here there is no mock Jacobi form at all, but only the polar correction term given by the Appell-Lerch sum, a kind of a ‘‘Cheshire cat’’ example which is all smile and no cat.

**Example 2: Weight 1, index 2, simple pole at  $z = 0$ .** As a second example take  $\varphi = BC/A \in J_{1,2}^{\text{mer}}$ . Here we find

$$\begin{aligned}\varphi^P &= 12 \text{Av}^{(2)}\left[\frac{y+1}{y-1}\right] = 12 \frac{y+1}{y-1} - 12(y^4 - y^{-4})q^2 + 24(y^5 - y^{-5})q^3 + \dots \\ \varphi^F &= (y - y^{-1}) - (y^3 + 45y - 45y^{-1} - y^{-3})q + (45y^3 - 231y + 231y^{-1} - 45y^{-3})q^2 + \dots,\end{aligned}$$

and by comparing the shadows and the first few Fourier coefficients (cf. eq. (7.32)), we see that  $\varphi^F$  is the negative of the mock Jacobi form  $\mathcal{F}_2(\tau, z) = h^{(2)}(\tau) \vartheta_1(\tau, 2z)$  discussed in Example 4 of §7.2 that is conjecturally related to representations of  $M_{24}$ . Using (4.31) and dividing by  $\vartheta_1(\tau, 2z)$ , we can rewrite the identity  $\varphi = \varphi^P + \varphi^F$  as

$$\eta(\tau)^{-3} \frac{\varphi_{0,1}(\tau, z)}{\varphi_{-2,1}(\tau, z)} = -\frac{12}{\vartheta_1(\tau, 2z)} \text{Av}^{(2)}\left[\frac{1+y}{1-y}\right] - h^{(2)}(\tau). \quad (8.52)$$

One can check that  $\vartheta_1(\tau, 2z)\mu(\tau, z) = \text{Av}^{(2)}\left[\frac{1+y}{1-y}\right]$ , so this agrees with the splitting (7.39) coming from the decomposition of the elliptic genus of  $K3$  into superconformal characters.

**Example 3: Weight 1, index 1/2, simple pole at  $z = 0$ .** We can also multiply (7.39) by  $\vartheta_1(\tau, z)$  and write it in the form

$$\frac{B}{\sqrt{A}} = -12 \text{Av}^{(\frac{1}{2}, -)}\left[\frac{\sqrt{y}}{1-y}\right] - h^{(2)}(\tau) \vartheta_1(\tau, z), \quad (8.53)$$

where  $\text{Av}^{(\frac{1}{2}, -)}[F(y)] := \sum_{n \in \mathbb{Z}} (-1)^n q^{n^2/2} y^n F(q^n y)$ . This decomposition also fits into the scheme explained in this section, but slightly modified because the index here is half-integral.

**Example 4: Weight 2, index 0, double pole at  $z = 0$ .** Now take  $\varphi = B/A$ , which, as we saw in §4.3, is just a multiple of the Weierstrass  $\wp$ -function. This example does not quite fit in with the discussion given so far, since we did not allow  $m = 0$  in the definition (8.9) of the averaging operator ( $m < 0$  doesn't work at all, because the sum defining the average diverges, and  $m = 0$  is less interesting since a form of index 0 is clearly determined up to a function of  $\tau$  alone by its singularities, so that in our discussion we excluded that case too), but nevertheless it works. The decomposition  $\varphi = \varphi^P + \varphi^F$  takes the form

$$\frac{\varphi_{0,1}(\tau, z)}{\varphi_{-2,1}(\tau, z)} = 12 \text{Av}^{(0)}\left[\frac{y}{(1-y)^2}\right] + E_2(\tau), \quad (8.54)$$

with the finite part being simply the quasimodular Eisenstein series  $E_2(\tau)$ , which is also a mock modular form. (It is automatically independent of  $z$  since the index is 0.)

**Example 5: Weight 2, index 1, double pole at  $z = 0$ .** The next example is a special case of an infinite family that will be treated in detail in §9. Take  $\varphi = B^2/A \in J_{2,1}^{\text{mer}}$ . Then we find

$$\begin{aligned}\varphi &= \left( \frac{144y}{(1-y)^2} + y + 22 + y^{-1} \right) + \left( 22y^2 + 152y - 636 + 152y^{-1} + 22y^{-2} \right) q \\ &\quad + \left( 145y^3 - 636y^2 + 3831y - 7544 + 3831y^{-1} - 636y^{-2} + 145y^{-3} \right) q^2 + \dots, \\ \varphi^P &= \text{Av}^{(1)} \left[ \frac{144y}{(1-y)^2} \right] = \frac{144y}{(1-y)^2} + 0q + 144(y^3 + y^{-3})q^2 + 288(y^4 + y^{-4})q^3 + \dots\end{aligned}$$

and hence that  $\varphi^F = \varphi - \varphi^P = \sum_{4n-r^2 \geq -1} C(4n-r^2) q^n y^r$  with the first few  $C(\Delta)$  given by

$\Delta$	-1	0	3	4	7	8	11	12	15
$C(\Delta)$	1	22	152	-636	3831	-7544	33224	-53392	191937

On the other hand, we have the weak Jacobi form  $E_4(\tau)\varphi_{-2,1}(\tau, z) = \sum C^*(4n-r^2)q^n y^r$  with

$\Delta$	-1	0	3	4	7	8	11	12	15
$C^*(\Delta)$	1	-2	248	-492	4119	-7256	33512	-53008	192513

We see that  $C(\Delta)$  and  $C^*(\Delta)$  are very close to each other (actually, asymptotically the same) and that their difference is precisely  $-288$  times the Hurwitz-Kronecker class number  $H(d)$ . We thus have  $\varphi^F = E_4A - 288\mathcal{H}$ , and we have found  $\mathcal{H}(\tau, z)$ , the simplest of all mock Jacobi forms, as the finite part of the meromorphic Jacobi form  $(E_4A^2 - B^2)/288A$ . We remark that the numbers  $-C^*(\Delta)$  have an interesting interpretation as “traces of singular moduli” [117].

**Example 6: Weight  $-5$ , index 1, simple poles at the 2-torsion points.** Finally, we give an example with more than one pole. Take  $\varphi = A^3/C \in J_{1,-5}$ . This function has three poles, all simple, at the three non-trivial 2-torsion points on the torus  $\mathbb{C}/(\mathbb{Z}\tau + \mathbb{Z})$ . The three corresponding modular forms, each of weight  $-6$ , are given by

$$D_{(0, \frac{1}{2})}(\tau) = 16 \frac{\eta(2\tau)^{12}}{\eta(\tau)^{24}}, \quad D_{(\frac{1}{2}, 0)}(\tau) = -\frac{1}{4} \frac{\eta(\frac{\tau}{2})^{12}}{\eta(\tau)^{24}}, \quad D_{(\frac{1}{2}, \frac{1}{2})}(\tau) = \frac{i}{4} \frac{\eta(\frac{\tau+1}{2})^{12}}{\eta(\tau)^{24}},$$

and one finds

$$\begin{aligned}\varphi = \varphi^P &= D_{(0, \frac{1}{2})}(\tau) \text{Av}^{(1)} \left[ \frac{1}{2} \frac{y-1}{y+1} \right] + q^{1/4} D_{(\frac{1}{2}, 0)}(\tau) \text{Av}^{(1)} \left[ \frac{1}{2y} \frac{y + \sqrt{q}}{y - \sqrt{q}} \right] \\ &\quad + q^{1/4} D_{(\frac{1}{2}, \frac{1}{2})}(\tau) \text{Av}^{(1)} \left[ \frac{1}{2y} \frac{y - \sqrt{q}}{y + \sqrt{q}} \right], \quad \varphi^F \equiv 0,\end{aligned}$$

another “Cheshire cat” example (of necessity, for the same reason as in Example 1, since again  $m = 1$  and  $k$  is odd).

## 9. Special mock Jacobi forms

We now turn to the study of certain families of meromorphic Jacobi forms of weights 1 or 2 and arbitrary positive index and of their related mock Jacobi forms, the weight 2 case being the one relevant for the application to black hole microstate counting. In this section, we introduce these families and formulate a number of properties of them that were found experimentally and that we will prove in §10. In particular, it will turn out that all of these functions can be expressed, using the Hecke-like operators introduced in §4.4, in terms of a collection of mock Jacobi forms  $\mathcal{Q}_m$  defined (though not quite uniquely: there is some choice involved starting at  $m = 91$ ) for all square-free  $m$ , of weight either 1 or 2 depending on whether  $m$  has an odd or even number of prime factors. These functions, which seem to have particularly nice arithmetic properties, include all of the examples studied in §7, and several other special mock modular forms that have appeared in the literature.

### 9.1 The simplest meromorphic Jacobi forms

The two simplest situations for the theory described in §8 are when the meromorphic Jacobi form  $\varphi = \varphi_{k,m}$  satisfies either

- $k = 1$  and  $\varphi(\tau, z)$  has a simple pole at  $z = 0$  and no other poles in  $\mathbb{C}/(\mathbb{Z}\tau + \mathbb{Z})$  or
- $k = 2$  and  $\varphi(\tau, z)$  has a double pole at  $z = 0$  and no other poles in  $\mathbb{C}/(\mathbb{Z}\tau + \mathbb{Z})$ ,

since in these two cases the modular forms  $D_s(\tau)$  (resp.  $E_s(\tau)$ ) defined in §8.2 (resp. §8.4) are simply constants and the canonical Fourier coefficients  $h_\ell$  of  $\varphi$  are pure, rather than mixed, mock modular forms. If we normalize  $\varphi$  by  $\varphi(\tau, z) \sim (2\pi iz)^{-1}$  in the first case and  $\varphi(\tau, z) \sim (2\pi iz)^{-2}$  in the second case as  $z \rightarrow 0$ , then any two choices of  $\varphi$  differ by a (weakly) holomorphic Jacobi form whose Fourier coefficients are true modular forms and are therefore well understood. It therefore suffices in principle to make some specific choice of  $\varphi_{1,m}$  and  $\varphi_{2,m}$  for each  $m$  and to study their finite parts  $\varphi_{k,m}^F$ , but, as we shall see, the problem of choosing a good representative  $\varphi$  is very interesting and has several subtleties. For any choice of  $\varphi_{k,m}$ , we will use the notation  $\Phi_{k,m} := \varphi_{k,m}^F$ , with decorations like  $\Phi_{k,m}^{\text{stand}}$  or  $\Phi_{k,m}^{\text{opt}}$  corresponding to the choice of  $\varphi_{k,m}$ . The shadow of the mock Jacobi form  $\Phi_{k,m}$  is independent of the choice of representative and is a multiple of  $\sum_{\ell \pmod{2m}} \overline{\vartheta_{m,\ell}^{2-k}(\tau)} \vartheta_{m,\ell}(\tau, z)$ , where  $\vartheta_{m,\ell}^1$  and  $\vartheta_{m,\ell}^0$  are defined as at the end of §4.2.

The Jacobi forms  $\varphi_{2,m}$  are the ones related to the Fourier coefficients  $\psi_m$  of  $\Phi_{10}(\Omega)^{-1}$  that appear in the application to black holes (eq. (1.1)). Indeed, since the principal part of the pole of  $\psi_m(\tau, z)$  at  $z = 0$  equals  $p_{24}(m+1) \Delta(\tau)^{-1} (2\pi iz)^{-2}$ , we can write  $\Delta(\tau)\psi_m(\tau, z)$  as the sum of  $p_{24}(m+1) \varphi_{2,m}$  and a weak Jacobi form. If we make the simplest choice for  $\varphi_{2,m}$ , namely

$$\varphi_{2,m}^{\text{stand}} = \frac{B^{m+1}}{12^{m+1}A}, \quad (9.1)$$



where  $A = \varphi_{-2,1}$ ,  $B = \varphi_{0,1}$  are the special Jacobi forms introduced in §4 (which has the correct principal part because  $A(\tau, z) \sim (2\pi iz)^2$  and  $B(\tau, z) \rightarrow 12$  as  $z \rightarrow 0$ ), then the first few values of  $\Delta(\tau) \psi_m(\tau, z) - p_{24}(m+1) \varphi_{2,m}^{\text{stand}}(\tau, z)$  can be read off from (5.16). For instance, we have  $p_{24}(3) = 3200$  and the fourth equation of (5.16) says that  $\Delta\psi_2 = 3200 \varphi_{2,2}^{\text{stand}} + \frac{16}{9} E_4 AB + \frac{10}{27} E_6 A^2$ .

In the weight one case we can also make a simple standard choice of  $\varphi$ . It has to be divisible by  $C$  because the weight is odd, and since  $C \sim 4\pi iz$  for  $z \rightarrow 0$  we can choose

$$\varphi_{1,m}^{\text{stand}} = \frac{B^{m-1} C}{2 \cdot 12^{m-1} A} = \frac{C}{2} \varphi_{2,m-2}^{\text{stand}}. \quad (9.2)$$

as our standard functions. However, we emphasize that (9.1) and (9.2) are only the starting points for our discussion and will be replaced by better choices as we go along.

We will apply the decomposition of §8 to the functions  $\varphi$ . The polar part is independent of the choice of  $\varphi$  and is given by  $\varphi_{k,m}^{\text{P}} = \mathcal{A}_{k,m}$ , where  $\mathcal{A}_{k,m}$  is given (in the notation of equations (8.17) and (8.43)) by

$$\mathcal{A}_{1,m} = \mathcal{A}_{1,m}^{(0,0)} = \text{Av}^{(m)} \left[ \frac{1}{2} \frac{y+1}{y-1} \right], \quad \mathcal{A}_{2,m} = \mathcal{A}_{2,m}^{(0,0)} = \text{Av}^{(m)} \left[ \frac{y}{(y-1)^2} \right] \quad (9.3)$$

or—written out explicitly—by

$$\mathcal{A}_{1,m}(\tau, z) = -\frac{1}{2} \sum_{s \in \mathbb{Z}} q^{ms^2} y^{2ms} \frac{1+q^s y}{1-q^s y}, \quad \mathcal{A}_{2,m}(\tau, z) = \sum_{s \in \mathbb{Z}} \frac{q^{ms^2+s} y^{2ms+1}}{(1-q^s y)^2}. \quad (9.4)$$

Also independent of the choice of  $\varphi$  is the correction term which must be added to the finite part and subtracted from the polar part of  $\varphi_{2,m}$  in order to make these functions transform like Jacobi forms. In particular, for the double pole case, we find from (8.48) that the completion  $\widehat{\Phi}_{2,m}$  of  $\Phi_{2,m} = \varphi_{2,m}^{\text{F}}$  for any choice of  $\varphi_{2,m}$  is given by

$$\widehat{\Phi}_{2,m}(\tau, z) = \Phi_{2,m}(\tau, z) - \sqrt{\frac{m}{4\pi}} \sum_{\ell \pmod{2m}} (\vartheta_{m,\ell}^0)^*(\tau) \vartheta_{m,\ell}(\tau, z), \quad (9.5)$$

where  $\vartheta_{m,\ell}^0(\tau) = \vartheta_{m,\ell}(\tau, 0)$  and  $(\vartheta_{m,\ell}^0)^*$  is normalized<sup>17</sup> as in equations (7.2) and (7.6).

The functions  $\mathcal{A}_{1,m}$ ,  $\mathcal{A}_{2,m}$  have wall-crossings, so their Fourier expansions depend on the value of  $z_2 = \text{Im}(z)$ . In the range  $0 < z_2 < \tau_2$ , or  $|q| < |y| < 1$ , they are given by

$$\mathcal{A}_{1,m} = \left( -\sum_{s \geq 0} \sum_{\ell \geq 0}^* + \sum_{s < 0} \sum_{\ell \leq 0}^* \right) q^{ms^2+\ell s} y^{2ms+\ell}, \quad (9.6)$$

$$\mathcal{A}_{2,m} = \left( \sum_{s \geq 0} \sum_{\ell \geq 0}^* - \sum_{s < 0} \sum_{\ell \leq 0}^* \right) \ell q^{ms^2+\ell s} y^{2ms+\ell}, \quad (9.7)$$

---

<sup>17</sup>Note that the function  $\vartheta_{m,\ell}^{(0,0)*}$  as defined in (8.47) is equal to  $1/\sqrt{4\pi m}$  times  $(\vartheta_{m,\ell}^0)^*$ .

where the asterisk on the summation sign has the usual meaning (count the term  $\ell = 0$  with multiplicity  $1/2$ ). Hence the coefficient of  $q^n y^r$  is zero unless  $\Delta = 4mn - r^2$  is the negative of the square of an integer congruent to  $r$  modulo  $2m$ , in which case it is  $0, \pm\frac{1}{2}, \pm 1$  or  $\pm 2$  if  $k = 1$  and is  $0, \pm\sqrt{-\Delta}$  or  $\pm 2\sqrt{-\Delta}$  if  $k = 2$ . (Compare the remark at the end of §8.3.) If we expand  $\mathcal{A}_{1,m}$  and  $\mathcal{A}_{2,m}$  in Fourier series for  $z_2$  in some other interval  $(N, N+1)\tau_2$ , then the formulas (9.6) and (9.7) are unchanged except for a shift in the inequalities defining the summation. In summary, the Fourier coefficients of the polar part of  $\varphi$ , although not quite like the coefficients of a usual Jacobi form—they do not only depend on  $\Delta$  and on  $r \bmod 2m$ , they are non-zero for infinitely many negative  $\Delta$ , and there is wall-crossing—are extremely simple and have very slow growth (at most like  $\sqrt{|\Delta|}$ ).

The interesting part of the Fourier expansion of  $\varphi_{k,m}$  is therefore the mock part  $\Phi_{k,m}$ . Here the choice of  $\varphi_{k,m}$  makes a big difference. For instance, Examples 1–4 of §8.5 say that the mock Jacobi forms  $\Phi_{k,m}^{\text{stand}}$  corresponding to the standard choices (9.1) and (9.2) for small  $m$  are

$$\Phi_{1,1}^{\text{stand}} = 0, \quad \Phi_{1,2}^{\text{stand}} = \frac{1}{24} \mathcal{F}_2, \quad \Phi_{2,0}^{\text{stand}} = \frac{1}{12} E_2, \quad \Phi_{2,1}^{\text{stand}} = \frac{1}{144} E_4 A - 2\mathcal{H}, \quad (9.8)$$

and we see that this choice is a good one in the first three cases (indeed, in these cases there *is* no choice!) but that, as we have already observed, a better choice of  $\varphi$  in the fourth case would have been the function  $\varphi_{2,1}^{\text{opt}} = \varphi_{2,1}^{\text{stand}} - \frac{1}{144} E_4 A$ , with mock part  $\Phi_{2,1}^{\text{opt}} = -2\mathcal{H}$ , because it has no terms  $q^n y^r$  with discriminant  $\Delta = 4n - r^2$  smaller than zero and (consequently) has Fourier coefficients that grow polynomially in  $\Delta$  (actually, like  $\Delta^{1/2+\varepsilon}$ ) rather than exponentially in  $\sqrt{\Delta}$ . Our next main objective, to which we devote the next subsection, is therefore to replace our initial choice of  $\varphi_{k,m}$  as given by (9.1) or (9.2) by an optimal choice  $\varphi_{k,m}^{\text{opt}}$ , where “optimal” means “with the negative discriminants  $d = 4mn - r^2$  that occur being as little negative as possible” or, equivalently, “with Fourier coefficients growing as slowly as possible.” This choice turns out to have Fourier coefficients of very much smaller growth than those of our initial choice  $\varphi_{k,m}^{\text{stand}}$  in all cases, and to be unique in many cases.

## 9.2 Choosing the function $\varphi$ for weight 2 and small index

We now consider in more detail the meromorphic Jacobi forms of weight 2 and index  $m$  with a double pole at the origin introduced in §9.1. Any such form will have the form

$$\varphi_{2,m} = \frac{\psi_{0,m+1}}{12^{m+1}A}, \quad \psi_{0,m+1} \in B^{m+1} + A \cdot \tilde{\mathcal{J}}_{2,m} \subset \tilde{\mathcal{J}}_{0,m+1}. \quad (9.9)$$

For each  $m$ , we want to choose a meromorphic Jacobi form  $\varphi_{2,m}^{\text{opt}}$  differing from  $\varphi_{2,m}^{\text{stand}}$  by a weak Jacobi form of weight 2 and index  $m$  whose coefficients grow as slowly as possible. We are interested both in the form of the corresponding  $\psi_{0,m+1} = \psi_{0,m+1}^{\text{opt}}$  as a polynomial in  $A$  and  $B$  with modular form coefficients, and in the nature of the finite part  $\Phi_{2,m}^{\text{opt}} = \varphi_{2,m}^{\text{opt}} - \mathcal{A}_{2,m}$  as a

mock Jacobi form. In this subsection we will look at the optimal choice for  $0 \leq m \leq 11$  to see what phenomena occur, while in the next subsection, we report on the calculations for higher values up to about 250 and formulate the general results.

For  $\mathbf{m}=\mathbf{0}$ , we have to choose

$$\varphi_{2,0}^{\text{opt}} = \varphi_{2,0}^{\text{stand}} = \frac{B}{12A}, \quad \psi_{0,1}^{\text{opt}} = B$$

because there are no weak Jacobi forms of weight 2 and index 0. The mock part  $\Phi_{2,0}^{\text{opt}}$  here is  $E_2(\tau)/12$ , as we have already seen (Example 3 of §8.5).

For  $\mathbf{m}=\mathbf{1}$  the only freedom is the addition of a multiple of  $E_4A$  and (as recalled in §9.1) the optimal choice is

$$\varphi_{2,1}^{\text{opt}} = \varphi_{2,1}^{\text{stand}} - \frac{E_4A}{144}, \quad \psi_{0,2}^{\text{opt}} = B^2 - E_4A^2, \quad (9.10)$$

because then  $\Phi_{2,1}^{\text{opt}} = -2\mathcal{H}$  has no terms  $q^n y^r$  with  $4n - r^2 < 0$  and its Fourier coefficients  $-2H(4n - r^2)$  have only polynomial growth as  $4n - r^2 \rightarrow \infty$ .

Now we look at  $\mathbf{m}=\mathbf{2}$ . Here we can modify  $12^3 \Phi_{2,2}^{\text{stand}} = B^3/A - 12^3 \mathcal{A}_{2,2}$  by adding to it any linear combination of the two weak Jacobi forms  $E_4AB$  and  $E_6A^2$ . Each of these three functions has Fourier coefficients of the form  $c(n, r) = C_\varphi(8n - r^2)$  (because the  $(n, r)$ -coefficient of a Jacobi form of prime or prime power index  $m$  and even weight always depends only on the discriminant  $4mn - r^2$ ), with the first few given by

$\Delta$	-4	-1	0	4	7	8	12	15	16
$C(12^3 \Phi_{2,2}^{\text{stand}}; \Delta)$	1	32	366	528	1056	-10144	-14088	92832	-181068
$C(E_4AB; \Delta)$	1	8	-18	144	2232	-4768	-22536	96264	-151884
$C(E_6A^2; \Delta)$	1	-4	6	-480	1956	-2944	-28488	96252	-139452

It follows from this table that the linear combination

$$\varphi_{2,2}^{\text{opt}} = \varphi_{2,2}^{\text{stand}} - \frac{1}{576} E_4AB + \frac{1}{864} E_6A^2 = \frac{B^3 - 3E_4A^2B + 2E_6A^3}{12^3 A} \quad (9.11)$$

has a mock Jacobi part  $\Phi_{2,2}^{\text{opt}} = \varphi_{2,2}^{\text{opt}} - \mathcal{A}_{2,2}$  with no negative terms and much smaller coefficients:

$d$	-4	-1	0	4	7	8	12	15	16
$C(\Phi_{2,2}^{\text{opt}}; \Delta)$	0	0	1/4	-1/2	-1	-1	-2	-2	-5/2

On inspection of these numbers and the next few values, we recognize them as simple linear combinations of Hurwitz class numbers, namely:

$$C(\Phi_{2,2}^{\text{opt}}; \Delta) = \begin{cases} -H(\Delta) & \text{if } \Delta \equiv 7 \pmod{8}, \\ -H(\Delta) - 2H(\Delta/4) & \text{if } \Delta \equiv 0 \pmod{4}, \end{cases}$$

which says that  $\Phi_{2,2}^{\text{opt}}$  is simply  $-\mathcal{H}|V_2$ , with  $\mathcal{H}$  as in (7.25) and  $V_2$  as in (4.37). (Note, by the way, the similarity between (9.11) and (4.33). Together they say that  $\varphi_{2,2}^{\text{opt}} = (C/2A)^2$ .)

The cases  $\mathbf{m=3}$ ,  $\mathbf{m=4}$ ,  $\mathbf{m=5}$ ,  $\mathbf{m=7}$ ,  $\mathbf{m=8}$ ,  $\mathbf{m=9}$  and  $\mathbf{m=11}$  are similar to the case  $m = 2$ . For each of these values of  $m$  the coefficient of  $q^n y^r$  in  $\Phi_{2,m}^{\text{stand}}$  depends only on  $\Delta = 4nm - r^2$ , the number of negative  $\Delta$  for which this coefficient is non-zero equals the dimension of the space of weak Jacobi forms of weight 2 and index  $m$ , and we can subtract a unique weak Jacobi form from  $\varphi_{2,m}^{\text{stand}}$  to obtain a meromorphic Jacobi form  $\varphi_{2,m}^{\text{opt}}$  whose finite part is a strongly holomorphic mock Jacobi form with very small Fourier coefficients. For example, for  $m = 3$  the optimal choice of  $\varphi_{2,m}^{\text{opt}}$  is  $\varphi_{2,3}^{\text{opt}} = \psi_{0,4}^{\text{opt}}/12^4 A$  with

$$\psi_{0,4}^{\text{opt}} = B^4 - 6E_4 A^2 B^2 + 8E_6 A^3 B - 3E_4^2 A^4 \quad (9.12)$$

and the first few Fourier coefficients of  $12^4 \Phi_{2,3}^{\text{stand}}$  and  $\Phi_{2,3}^{\text{opt}}$  are given by

$\Delta$	-9	-4	-1	0	3	8	11	12	15
$12^4 C(\Phi_{2,3}^{\text{stand}}; \Delta)$	1	42	687	5452	1104	8088	-1488	-139908	-19629
$C(\Phi_{2,3}^{\text{opt}}; \Delta)$	0	0	0	1/3	-1/3	-1	-1	-4/3	-2

Again we recognize the coefficients  $C(\Phi_{2,3}^{\text{opt}}; \Delta)$  as  $-H(\Delta)$  if  $9 \nmid \Delta$  and  $-H(\Delta) - 3H(\Delta/9)$  if  $9 \mid \Delta$ , so that just as for  $m = 2$  we find that the mock part of  $\varphi_{2,3}^{\text{opt}}$  equals  $-\mathcal{F}_1|V_3$ . The same thing also happens for  $m = 5, 7$  and  $11$ , suggesting that for prime values of  $m$  we can always choose

$$\Phi_{2,p}^{\text{opt}} = -\mathcal{H}|V_p \quad (p \text{ prime}). \quad (9.13)$$

This will be proved in §10. For the prime-power values  $m = 4, 8$  and  $9$  this formula no longer holds, but the coefficients of  $\Phi_{2,m}^{\text{opt}}$  are still simple linear combinations of Hurwitz-Kronecker class numbers and we find the expressions

$$\Phi_{2,4}^{\text{opt}} = -\mathcal{H}|V_4 + 2\mathcal{H}|U_2, \quad \Phi_{2,8}^{\text{opt}} = -\mathcal{H}|V_8 + 2\mathcal{H}|U_2 V_2, \quad \Phi_{2,9}^{\text{opt}} = -\mathcal{H}|V_9 + 3\mathcal{H}|U_3,$$

where  $U_t$  is the Hecke-like operator defined in §4.4 that multiplies the index of a Jacobi form (or mock Jacobi form) by  $t^2$ . This suggests that for prime power values of  $m$  we can choose

$$\Phi_{2,m}^{\text{opt}} = -2\mathcal{H}|\mathcal{V}_{2,m}^{(1)} \quad (m \text{ a prime power}) \quad (9.14)$$

where  $\mathcal{V}_{k,t}^{(1)}$  for  $t \in \mathbb{N}$  is the Hecke-like operator from Jacobi (or weakly holomorphic Jacobi, or mock Jacobi) forms of weight  $k$  and index 1 to forms of weight  $k$  and index  $t$  defined in (4.38). Formula (9.14) will also be proved in §10.

The data for the values discussed so far ( $0 \leq m \leq 11$ ,  $m \neq 6, 10$ ) is summarized in the table below, which expresses the weak Jacobi form  $\psi_{0,m+1}^{\text{opt}} = 12^{m+1} A \varphi_{2,m}^{\text{opt}}$  for each of these values of  $m$  as a polynomial in  $A$  and  $B$ . In this table we notice many regularities, e.g.,

the coefficients of  $E_4 A^2 B^{m-2}$ ,  $E_6 A^3 B^{m-3}$  and  $E_4^2 A^4 B^{m-4}$  in  $\psi_{0,m}^{\text{opt}}$  are always given by  $-\binom{m}{2}$ ,  $2\binom{m}{3}$  and  $-3\binom{m}{4}$ , respectively, and more generally, if we write  $\psi_{0,m}^{\text{opt}}$  as  $\sum_{i=0}^m f_{m,i} A^i B^{m-i}$  with  $f_{m,i} = f_{m,i}(\tau) \in M_{2i}$ ,  $f_{m,0} = 1$ , then the constant terms  $f_{m,i}(\infty)$  of the modular forms  $f_{m,i}$  (which can be read off from the table by setting  $E_4 = E_6 = 1$  and  $D = 0$ ) are given in each case by  $f_{m,i}(\infty) = (-1)^{i-1}(i-1)\binom{m}{i}$  or equivalently by the generating function

$$\sum_{i=0}^m f_{m,i}(\infty) X^{m-i} = (X-1)^m + m(X-1)^{m-1} = (X-1)^{m-1}(X+m-1), \quad (9.15)$$

e.g.  $X^8 - 28X^6 + 112X^5 - 210X^4 + 224X^3 - 140X^2 + 48X - 7 = (X-1)^7(X+7)$ . This formula has a simple interpretation: the constant term with respect to  $q$  of  $\sum f_{m,i} A^i B^{m-i}$  is  $\sum f_{m,i}(\infty)(y^{-1} - 2 + y)^i (y^{-1} + 10 + y)^{m-i}$  and this Laurent polynomial—subject to the condition  $f_{m,1}(\infty) = 0$ , which comes from the fact that there are no modular forms of weight 2 on  $SL(2, \mathbb{Z})$ —has minimal growth  $O(y + y^{-1})$  precisely for the coefficients defined by (9.15).

$m$	1	2	3	4	5	6	8	9	10	12
$B^m$	1	1	1	1	1	1	1	1	1	1
$E_4 \cdot A^2 B^{m-2}$	0	-1	-3	-6	-10	-15	-28	-36	-45	-66
$E_6 \cdot A^3 B^{m-3}$	0	0	2	8	20	40	112	168	240	440
$E_4^2 \cdot A^4 B^{m-4}$	0	0	0	-3	-15	-45	-210	-378	-630	-1485
$E_4 E_6 \cdot A^5 B^{m-5}$	0	0	0	0	4	24	224	504	1008	3168
$E_4^3 \cdot A^6 B^{m-6}$	0	0	0	0	0	-5	-140	-420	-1050	-4620
$D \cdot A^6 B^{m-6}$	0	0	0	0	0	1	10	30	48	222
$E_4^2 E_6 \cdot A^7 B^{m-7}$	0	0	0	0	0	0	48	216	720	4752
$E_4^4 \cdot A^8 B^{m-8}$	0	0	0	0	0	0	-7	-63	-315	-3465
$E_4 D \cdot A^8 B^{m-8}$	0	0	0	0	0	0	-4	-36	-18	-360
$E_4^3 E_6 \cdot A^9 B^{m-9}$	0	0	0	0	0	0	0	8	80	1760
$E_6 D \cdot A^9 B^{m-9}$	0	0	0	0	0	0	0	20	-16	80
$E_4^5 \cdot A^{10} B^{m-10}$	0	0	0	0	0	0	0	0	-9	-594
$E_4^2 D \cdot A^{10} B^{m-10}$	0	0	0	0	0	0	0	0	18	702
$E_4^4 E_6 \cdot A^{11} B^{m-11}$	0	0	0	0	0	0	0	0	0	120
$E_4 E_6 D \cdot A^{11} B^{m-11}$	0	0	0	0	0	0	0	0	0	-672
$E_4^6 \cdot A^{12} B^{m-12}$	0	0	0	0	0	0	0	0	0	-11
$E_4^3 D \cdot A^{12} B^{m-12}$	0	0	0	0	0	0	0	0	0	188
$D^2 \cdot A^{12} B^{m-12}$	0	0	0	0	0	0	0	0	0	-8

**Table 2:** Coefficients of  $\psi_{0,m}^{\text{opt}}$  in terms of the standard basis. Here  $D := 2^{11}3^3\Delta$ .

For  $\mathbf{m}=6$  several new things happen, all connected with the fact that 6 is the first value

of  $m$  having more than one prime factor. First of all, the formula (9.14), which worked for prime powers, now fails:  $-2\mathcal{H}|\mathcal{V}_{2,6}^{(1)}$  is not a possible choice for  $\Phi_{2,6}$ . The reason is that, as we will see later in general, the form  $\mathcal{H}|\mathcal{V}_{2,m}^{(1)}$  (for any  $m$ ) is invariant under all Atkin-Lehner involutions, but  $\Phi_{2,m}$  is not when  $m$  is not a prime power. The expression  $-2\mathcal{H}|\mathcal{V}_{2,6}^{(1)}$  does, however, give the optimal choice of the eigencomponent  $\Phi_{2,6}^{++}$  of  $\Phi_{2,6}$  with eigenvalues  $W_2 = W_3 = +1$ . The other eigencomponent, corresponding to  $W_2 = W_3 = -1$ , is obtained by subtracting a multiple of the non-weak Jacobi form  $\varphi_{2,6}^-$  given in (4.48) from  $\pi_2^-(\Phi_{2,6}^{\text{opt}})$  (which is weakly holomorphic but not weak), and turns out to be simply a multiple of the mock Jacobi form  $\mathcal{F}_6$  defined in §7.2 (Example 3), giving as our first choice of the “optimal” mock modular form

$$\Phi_{2,6}^{\text{I}} = -2\mathcal{H}|\mathcal{V}_{2,6}^{(1)} - \frac{1}{24}\mathcal{F}_6. \quad (9.16)$$

The second new phenomenon for  $m = 6$  is that requiring the order of the poles to be as small as possible no longer fixes a unique optimal choice. The function  $\Phi_{2,6}^{\text{I}}$  defined above has an even part that is strongly holomorphic (because  $\mathcal{H}$  is) and an odd part that has discriminant  $\Delta \geq -1$  (because  $C^{(6)}(\Delta) = 0$  for  $\Delta < -1$ ), but we can change this choice without disturbing the discriminant bound  $\Delta \geq -1$  by adding to it an arbitrary multiple of the function

$$\mathcal{K}_6 = \frac{E_4AB^5 - 5E_6A^2B^4 + 10E_4^2A^3B^3 - 10E_4E_6A^4B^2 + (5E_4^3 - \frac{1}{4}D)A^5B - E_4^2E_6A^5}{12^5} \quad (9.17)$$

which is (up to a scalar factor, fixed here by requiring  $\mathcal{K}_6 = y - 2 + y^{-1} + O(q)$ ) the unique Jacobi form of weight 2 and index 6 satisfying the same discriminant bound. Two particularly nice choices besides the function  $\varphi_{2,6}^{\text{I}}$  are the functions

$$\varphi_{2,6}^{\text{II}} = \varphi_{2,6}^{\text{I}} - \frac{1}{24}\mathcal{K}_6 \quad \varphi_{2,6}^{\text{III}} = \varphi_{2,6}^{\text{I}} + \frac{1}{4}\mathcal{K}_6 \quad (9.18)$$

The special properties of these two choices are that  $\Phi_{2,6}^{\text{II}}$  is the unique choice with  $c(\Phi; 0, 1) = 0$  and (hence) has the form  $12^{-7} \sum_{i=0}^7 f_{7,i} A^{i-1} B^{7-i}$  with  $f_{7,i}$  satisfying (9.15), while  $\Phi_{2,6}^{\text{III}}$  is the unique choice for which  $c(n, r) = 0$  whenever  $24n = r^2$ . (This last choice is from some points of view the most canonical one.) Each of these three special choices, as we shall see below, has analogues for all  $m$ , although in general none of them is canonical, since for large  $m$  we can change them by adding cusp forms without affecting their defining properties and we do not know any criterion that selects a unique form.

The following table, which gives the first few Fourier coefficients  $C(\Phi; \Delta, r \pmod{12})$  for the Jacobi form  $\mathcal{K}_6$  and each of the five mock Jacobi forms  $-2\mathcal{H}|\mathcal{V}_{2,6}^{(1)}$ ,  $\mathcal{F}_6$ ,  $\Phi_{2,6}^{\text{I}}$ ,  $12\Phi_{2,6}^{\text{II}}$  and  $\Phi_{2,6}^{\text{III}}$ , summarizes the above discussion. In this table we have set in boldface the Fourier coefficients  $C(\Delta, r)$  whose special properties determine the choice of  $\Phi$  (namely,  $C(-1, \pm 1) + C(-1, \pm 5) = 0$  for  $\Phi^{\text{I}}$ ,  $C(-1, \pm 1) = 0$  for  $\Phi^{\text{II}}$ , and  $C(0, 0) = 0$  for  $\Phi^{\text{III}}$ ). We mention in passing

$\Delta$	-1	-1	0	8	12	15	20	23	23	24	32	36	39
$r \pmod{12}$	$\pm 1$	$\pm 5$	0	$\pm 4$	6	$\pm 3$	$\pm 2$	$\pm 1$	$\pm 5$	0	$\pm 4$	6	$\pm 3$
$\mathcal{K}_6$	1	1	-2	10	28	10	-12	13	13	-44	58	104	2
$-2\mathcal{H}   \mathcal{V}_{2,6}$	0	0	$\frac{1}{2}$	$-\frac{1}{2}$	-1	-1	-1	$-\frac{3}{2}$	$-\frac{3}{2}$	-1	$-\frac{5}{2}$	-2	-2
$\mathcal{F}_6$	-1	1	0	0	0	0	0	35	-35	0	0	0	0
$\Phi_{2,6}^I$	$\frac{1}{24}$	$-\frac{1}{24}$	$\frac{1}{2}$	$-\frac{1}{2}$	-1	-1	-1	$-\frac{71}{24}$	$-\frac{1}{24}$	-1	$-\frac{5}{2}$	-2	-2
$12 \Phi_{2,6}^{II}$	<b>0</b>	-1	7	-11	-26	-17	-6	-42	-7	10	-59	-76	-25
$\Phi_{2,6}^{III}$	$\frac{7}{24}$	$\frac{5}{24}$	<b>0</b>	2	6	$\frac{3}{2}$	-4	$\frac{7}{24}$	$\frac{77}{24}$	-12	12	24	$-\frac{3}{2}$

**Table 3:** Jacobi and mock Jacobi forms of weight 2 and index 6

that the function  $\psi_{0,7}^I = 12^7 A \varphi_{2,6}^I$  corresponding to (9.16) is precisely equal to  $B$  times  $\psi_{0,6}^{\text{opt}}$  as given in Table 2. We do not know if this has any special significance. It may be connected with the fact that  $\psi_{0,6}^{\text{opt}}$ , which is equal to the quotient  $\vartheta_1(\tau, 4z)/\vartheta_1(\tau, 2z)$  (Watson quintuple product identity), is a  $(-1)$ -eigenvector of the involution  $W_2 = W_3$  on  $\tilde{\mathcal{J}}_{0,6}$ .

For  $\mathbf{m}=\mathbf{10}$  the situation is similar. Again, we can achieve  $\Delta \geq -1$  but not uniquely because there is a weak Jacobi form  $\mathcal{K}_{10} \in \tilde{\mathcal{J}}_{2,10}$  (unique up to a normalizing factor, which we fix in the same way as before) satisfying the same discriminant bound and we can change  $\varphi_{2,10}$  an arbitrary multiple of  $\mathcal{K}_{10}$ . We again have three distinguished choices

$$\varphi_{2,10}^{II} = \frac{\psi_{0,11}^{II}}{12^{11}A}, \quad \varphi_{2,10}^I = \varphi_{2,10}^{II} + \frac{1}{12} \mathcal{K}_{10} \quad \varphi_{2,10}^{III} = \varphi_{2,10}^I + \frac{3}{8} \mathcal{K}_{10}, \quad (9.19)$$

where  $\psi_{0,11}^{II} \in \tilde{\mathcal{J}}_{0,11}$  is given (with  $D = 32(E_4^3 - E_6^2) = 2^{11}3^3\Delta$  as in Table 2) by

$$\begin{aligned} \Psi_{11}^{II} = & B^{11} - 55E_4A^2B^9 + 330E_6A^3B^8 - 990E_4^2A^4B^7 + 1848E_4E_6A^5B^6 - (2310E_4^3 - 111D)A^6B^5 \\ & + 1980E_4^2E_6A^7B^4 - (1155E_4^4 + 120E_4D)A^8B^3 + (440E_4^3E_6 + 20E_6D)A^9B^2 \\ & - (99E_4^5 - 117E_4^2D)A^{10}B + (10E_4^4E_6 - 56E_4E_6D)A^{11} \end{aligned} \quad (9.20)$$

and is the unique choice satisfying (9.15),  $\Phi_{2,10}^{III}$  is the unique choice with no  $\Delta = 0$  coefficients, and  $\Phi_{2,6}^I$  is the unique choice whose invariant part under  $W_2$  (or  $W_5$ ) is strongly holomorphic. Explicitly, we have

$$\Phi_{2,10}^I = -2\mathcal{H} | \mathcal{V}_{2,10}^{(1)} - \frac{1}{12} \mathcal{F}_{10} \quad (9.21)$$

where  $\mathcal{F}_{10}$  is a mock Jacobi form of weight 2 and index 10 with Fourier coefficients given by

$$c(\mathcal{F}_{10}; n, r) = \begin{cases} C^{(10)}(40n - r^2) & r \equiv \pm 1 \pmod{20} \text{ or } r \equiv \pm 3 \pmod{20} \\ -C^{(10)}(40n - r^2) & r \equiv \pm 7 \pmod{20} \text{ or } r \equiv \pm 9 \pmod{20} \end{cases} \quad (9.22)$$

for a function  $C^{(10)}(\Delta)$  whose first few values are



$\Delta$	-1	31	39	71	79	111	119	151	159	191	199	231	239
$C^{(10)}(\Delta)$	-1	9	21	35	63	57	112	126	207	154	306	315	511

or equivalently, where  $\mathcal{F}_{10}$  has a theta expansion of the form

$$\mathcal{F}_{10}(\tau, z) = \sum_{\substack{\ell \pmod{20} \\ (\ell, 10)=1}} h_\ell(\tau) \vartheta_{10, \ell}(\tau, z) \quad (9.23)$$

with mock modular forms  $h_\ell = h_\ell(\tau)$  having expansions beginning

$$h_1 = -h_9 = -h_{11} = h_{19} = q^{-1/40} (-1 + 21q + 63q^2 + 112q^3 + \dots), \quad (9.24)$$

$$h_3 = -h_7 = -h_{13} = h_{17} = q^{31/40} (9 + 35q + 57q^2 + 126q^3 \dots). \quad (9.25)$$

### 9.3 Further data and formulation of main results for the weight 2 family

In §9.2, we found that the functions  $\Phi_{2,m}$  for  $m \leq 11$  could all be obtained, using the Hecke-like operators from §4.4, from just three functions  $\mathcal{H}$ ,  $\mathcal{F}_6$  and  $\mathcal{F}_{10}$ . Continuing, we find that for  $\mathbf{m} = \mathbf{12}$  the same statement holds: the  $(+1, +1)$  eigencomponent of this form with respect to the involutions  $W_4$  and  $W_3$  can be chosen to be  $-2\mathcal{H}|\mathcal{V}_{2,12}^{(1)}$  and the  $(-1, -1)$  eigencomponent can be chosen to be  $\frac{1}{12}\mathcal{F}_6|\mathcal{V}_{2,2}^{(6)}$ , with  $\mathcal{V}_{k,t}^{(M)}$  as in (4.38) (so  $V_{2,12}^{(1)} = \frac{1}{4}(V_{12} - 2V_2U_3)$  and  $\mathcal{V}_{2,2}^{(6)} = \frac{1}{2}V_2$ ). Thus

$$\Phi_{2,12} = -2\mathcal{H}|\mathcal{V}_{2,12}^{(1)} + \frac{1}{12}\mathcal{F}_6|\mathcal{V}_{2,2}^{(6)} \quad (9.26)$$

is a possible choice for  $\Phi_{2,12}$ . Similarly, for  $\mathbf{m} = \mathbf{18}$  and  $\mathbf{m} = \mathbf{24}$  we can take the functions

$$\Phi_{2,18} = -2\mathcal{H}|\mathcal{V}_{2,18}^{(1)} + \frac{1}{24}\mathcal{F}_6|\mathcal{V}_{2,3}^{(6)}, \quad \Phi_{2,24} = -2\mathcal{H}|\mathcal{V}_{2,24}^{(1)} + \frac{1}{24}\mathcal{F}_6|\mathcal{V}_{2,4}^{(6)} \quad (9.27)$$

as our representatives of the coset  $\Phi_{2,m}^{\text{stand}} + \tilde{J}_{2,m}$ .

On the other hand, the indices  $\mathbf{m} = \mathbf{14}$  and  $\mathbf{m} = \mathbf{15}$  are similar to the indices  $m = 6$  and  $m = 10$  looked at previously: here we find optimal mock Jacobi forms of the form

$$\Phi_{2,14} = -2\mathcal{H}|\mathcal{V}_{2,14}^{(1)} - \frac{1}{8}\mathcal{F}_{14}, \quad \Phi_{2,15} = -2\mathcal{H}|\mathcal{V}_{2,15}^{(1)} - \frac{1}{6}\mathcal{F}_{15}, \quad (9.28)$$

where  $\mathcal{F}_{14}$  and  $\mathcal{F}_{15}$  are forms anti-invariant under  $W_p$  ( $p|m$ ) with Fourier expansions of the form

$$c(\mathcal{F}_{14}; n, r) = \begin{cases} C^{(14)}(56n - r^2) & \text{for } r \equiv \pm 1, \pm 3 \text{ or } \pm 5 \pmod{28}, \\ -C^{(14)}(56n - r^2) & \text{for } r \equiv \pm 9, \pm 11, \text{ or } \pm 13 \pmod{28}, \end{cases} \quad (9.29)$$

$$c(\mathcal{F}_{15}; n, r) = \begin{cases} C^{(15)}(60n - r^2) & \text{for } r \equiv \pm 1, \pm 2, \pm 4, \text{ or } \pm 7 \pmod{30}, \\ -C^{(15)}(60n - r^2) & \text{for } r \equiv \pm 8, \pm 11, \pm 13, \text{ or } \pm 14 \pmod{30}, \end{cases} \quad (9.30)$$

with coefficients  $C^{14}(\Delta)$  and  $C^{15}(\Delta)$  given for small  $\Delta$  by



$\Delta$	-1	31	47	55	87	103	111	143	159	167	199	215	223
$C^{(14)}(\Delta)$	-1	3	10	15	15	30	42	20	51	65	51	85	120

$\Delta$	-1	11	44	56	59	71	104	116	119	131	164	176	179
$C^{(15)}(\Delta)$	-1	2	10	4	14	9	28	24	29	26	40	28	54

This pattern continues: as we continue to higher values, we find that for each index  $M$  which is a product of an even number of distinct primes (i.e., for which  $\mu(M) = 1$ , where  $\mu(\cdot)$  denotes the Möbius function), we need to introduce a new function  $\mathcal{Q}_M$  that is anti-invariant under  $W_p$  for all  $p$  dividing  $M$ , but that these functions and their images under Hecke-like operators then suffice to give possible choices for our special mock Jacobi forms in all indices, in the sense that the function

$$\Phi_{2,m} = \frac{1}{2^{\omega(m)-1}} \sum_{\substack{M|m \\ \mu(M)=+1}} \mathcal{Q}_M | \mathcal{V}_{2,m/M}^{(M)} \quad (\omega(m) := \#\{p \mid p \text{ prime}, p|M\}) \quad (9.31)$$

belongs to the coset  $\Phi_{k,m}^{\text{stand}} + \tilde{J}_{2,m}$  for all  $m \geq 1$ . (The factor  $2^{1-\omega(m)}$  will be explained below.) In fact more is true. The functions  $\mathcal{Q}_M$  in (9.31) are simply the  $(-1, \dots, -1)$  eigencomponents of  $\Phi_{2,M}$  for some choice of these latter functions within their defining cosets, and the various terms in (9.31) are the eigencomponents of  $\Phi_{2,m}$  for the involutions  $W_{m_1}$  ( $m_1 || m$ ). In other words, we have the following theorem:

**Theorem 9.1.** *For any choice of the mock Jacobi forms  $\Phi_{2,m} \in \Phi_{2,m}^{\text{stand}} + \tilde{J}_{2,m}$  ( $m \in \mathbb{N}$ ), the eigencomponents  $\Phi_{2,m}^{\varepsilon_1, \dots, \varepsilon_r}$  ( $m = \prod_i p_i^{\nu_i}$ ,  $\varepsilon_i = \pm 1$ ,  $\prod \varepsilon_i = +1$ ) with respect to the decomposition<sup>18</sup> (4.41) are given by*

$$\frac{1}{2} \Phi_{2,m} \Big| \prod_{i=1}^r (1 + \varepsilon_i W_{p_i^{\nu_i}}) = \mathcal{Q}_M | \mathcal{V}_{2,m/M}^{(M)} \pmod{(J_{2,m}^1)^{\varepsilon_1, \dots, \varepsilon_r}}, \quad (9.32)$$

where  $M$  is the product of the  $p_i$  with  $\varepsilon_i = -1$ ,  $\mathcal{Q}_M$  is defined by

$$\mathcal{Q}_M = \frac{1}{2} \Phi_{2,M} \Big| \prod_{p|M} (1 - W_p) = 2^{\omega(M)-1} \Phi_{2,M}^{-, \dots, -} \quad (\mu(M) = +1), \quad (9.33)$$

and  $\mathcal{V}_{2,m/M}^{(M)}$  is the Hecke-like operator defined in (4.38). In particular, the decomposition (9.31) is always true modulo  $\tilde{J}_{2,m}$ , and if we choose the functions  $\Phi_{2,M}$  and consequently  $\mathcal{Q}_M$  arbitrarily for all  $M$  with  $\mu(M) = 1$ , then the function defined by (9.31) gives an admissible choice of  $\Phi_{2,m}$  for all indices  $m \geq 1$ .

---

<sup>18</sup>or rather, its analogue for elliptic forms

**Examples.** The functions  $\mathcal{Q}_M$  for the first few values of  $M$  with  $\mu(M) = 1$  are

$$\mathcal{Q}_1 = -\mathcal{H}, \quad \mathcal{Q}_6 = -\frac{\mathcal{F}_6}{12}, \quad \mathcal{Q}_{10} = -\frac{\mathcal{F}_{10}}{6}, \quad \mathcal{Q}_{14} = -\frac{\mathcal{F}_{14}}{4}, \quad \mathcal{Q}_{15} = -\frac{\mathcal{F}_{15}}{3}, \quad \dots, \quad (9.34)$$

where  $\mathcal{H}, \mathcal{F}_6, \dots, \mathcal{F}_{15}$  are the functions introduced in §7.2, §9.2, and this subsection. (The reasons for the various normalizations, and for having two symbols  $\mathcal{F}_M$  and  $\mathcal{Q}_M$  to denote proportional functions, will be discussed later.)

We will prove Theorem 9.1 towards the end of this section by calculating the shadows of the mock Jacobi forms on the two sides of (9.32), since two mock Jacobi forms differ by a true Jacobi form if and only if they have the same shadow.

**Remarks. 1.** We wrote  $(J_{2,m}^!)^{\varepsilon_1, \dots, \varepsilon_r}$  rather than  $\tilde{J}_{2,m}^{\varepsilon_1, \dots, \varepsilon_r}$  in the theorem because of the fact mentioned in §4.4 that the involutions  $W_{m_1}$  do not preserve the space of weak Jacobi forms (or weak elliptic forms), so that the left-hand side of eq. (9.32) for an arbitrary choice of  $\Phi_{2,m}$  may contain monomials  $q^n y^r$  with  $n < 0$ . In fact, it will follow from the results given below that the functions  $\Phi_{2,M}$  can be chosen in such a way that the eigencomponents  $\Phi_{2,M}^{\dots, -}$  do not contain any negative powers of  $q$ , in which case the expression on the right-hand side of (9.31) really does lie in  $\Phi_{k,m}^{\text{stand}} + \tilde{J}_{2,m}$  as asserted, and not merely in  $\Phi_{k,m}^{\text{stand}} + J_{2,m}^!$ .

**2.** An exactly similar statement is true for the functions  $\Phi_{1,m}$ , except that in that case we have  $\prod_i \varepsilon_i = -1$  (because in odd weight  $W_m$  acts as multiplication by  $-1$ ), so that the  $M$  that occur in the decomposition of  $\Phi_{1,m}$  are the divisors of  $m$  with  $\mu(M) = -1$ , and the functions  $\mathcal{Q}_M$  that occur in the decomposition of  $\Phi_{1,M}$  are defined by

$$\mathcal{Q}_M = \frac{1}{2} \Phi_{1,M} \left| \prod_{p|M} (1 - W_p) = 2^{\omega(M)-1} \Phi_{2,M}^{\dots, -} \quad (\mu(M) = -1) \quad (9.35)$$

instead of equation (9.33). The reason for the factor  $2^{\omega(m)-1}$  in these two equations is now apparent: it is the minimal factor making  $\mathcal{Q}_M$  an integral linear combination of images of  $\Phi_{k,m}$  under Atkin-Lehner involutions. (We could remove one factor 2 because for each decomposition  $M = M_1 M_2$ , the two terms  $\mu(M_1) \Phi_{k,m} | W_{M_1}$  and  $\mu(M_2) \Phi_{k,m} | W_{M_2}$  are equal.)

Theorem 9.1 explains the first part of our observations, those connected with the eigencomponent decomposition of the mock Jacobi forms  $\Phi_{2,m}$  and the fact that the collection of all of these functions can be expressed in terms of the special functions (9.34) and their images under Hecke-like operators. But it does not say anything about the orders of the poles of the functions in question, which was our original criterion for optimality. On the other hand, from the tables of Fourier coefficients of the functions  $\mathcal{Q}_1, \dots, \mathcal{Q}_{15}$  that we have given we see that, although none of these functions except  $\mathcal{Q}_1$  is strongly holomorphic, each of them satisfies the only slightly weaker discriminant bound  $\Delta := 4mn - r^2 \geq -1$ . This property will turn out to play a prominent role. We will call Jacobi or elliptic forms satisfying it **forms with optimal**

**growth**<sup>19</sup>, because their Fourier coefficients grow more slowly than forms that do not obey this discriminant bound. We observe that any Jacobi or elliptic form having optimal growth is automatically a weak form, since  $4nm - r^2 \geq -1$  implies  $n \geq 0$ .

As we continued our computations to higher values of  $M$ , we were surprised to find that this “optimal growth” property continued to hold for each  $M$  of the form  $pq$  ( $p$  and  $q$  distinct primes) up to 200. It followed that the full function  $\Phi_{2,M}$  could also be chosen to have optimal growth in these cases, since  $\Phi_{2,M} = \mathcal{Q}_1 | \mathcal{V}_{2,M}^{(1)} + \mathcal{Q}_M$  and the image of the strongly holomorphic form  $\mathcal{Q}_1$  under  $\mathcal{V}_{2,M}^{(1)}$  is strongly holomorphic. This made it particularly interesting to look at the case  $\mathbf{m} = \mathbf{210}$ , the first product of four distinct primes, because in this case, even if the function  $\mathcal{Q}_{210} = \Phi_{2,210}^-$  can be chosen to have optimal growth, then the six middle terms in the decomposition (9.31), which here takes the form

$$\begin{aligned} 8 \Phi_{2,210} &= \mathcal{Q}_1 | \mathcal{V}_{2,210}^{(1)} + \mathcal{Q}_6 | \mathcal{V}_{2,35}^{(6)} + \mathcal{Q}_{10} | \mathcal{V}_{2,21}^{(10)} + \mathcal{Q}_{14} | \mathcal{V}_{2,15}^{(14)} \\ &\quad + \mathcal{Q}_{15} | \mathcal{V}_{2,14}^{(15)} + \mathcal{Q}_{21} | \mathcal{V}_{2,10}^{(21)} + \mathcal{Q}_{35} | \mathcal{V}_{2,6}^{(35)} + \mathcal{Q}_{210} , \end{aligned} \quad (9.36)$$

will not be. (Even if  $\mathcal{Q}_M$  is chosen to have optimal growth, its image under the operator  $\mathcal{V}_{2,t}^M$  will in general satisfy only  $\Delta \geq -t^2$ . For  $M$  of the form  $pq$  we had no problems with this because  $\mathcal{Q}_1$ , and hence also its image under  $\mathcal{V}_{2,M}^{(1)}$ , is actually holomorphic.) The corresponding computation was somewhat forbidding since the space  $\tilde{\mathcal{J}}_{2,210}$  of forms by which we can change the initial choice  $\Phi_{2,210}^{\text{stand}}$  has dimension 3815, and the conditions  $C(\Delta, r) = 0$  for  $\Delta < -1$  give 3813 linear constraints, so that we have to find the kernel of a roughly square matrix of size nearly 4000. Moreover, the coefficients of this matrix are gigantic, so that we have to work modulo some large prime. Nevertheless, this computation could be carried out (using PARI/GP), and it turned out that the function  $\mathcal{Q}_{210}$  could again be chosen to have optimal growth! This solution is given by  $\mathcal{Q}_{210} = -\mathcal{F}_{210}$ , where  $\mathcal{F}_{210} = \sum_{\ell \pmod{420}} h_\ell \vartheta_{210,\ell}$  with the Fourier coefficients of the mock theta functions  $h_\ell$  given in Table 4. In this table, we have listed the values of  $\ell$  with  $(\ell, 210) = 1$  and  $0 \leq \ell < 210$  (since  $h_{420n \pm \ell} = h_\ell$  and  $h_\ell = 0$  for  $(\ell, 210) > 1$ ) in groups of 8, with  $h_\ell = h_{\ell_0}$  for the first four and  $h_\ell = -h_{\ell_0}$  for the second four, where  $\ell_0$  (shown in boldface) belongs to  $\{1, 11, 13, 17, 19, 23\}$  and  $h_{\ell_0} = \sum_{n \geq 0} C(n) q^{n - \ell_0^2/840}$ . Thus, the third line of the table means that  $h_{\pm 13} = h_{\pm 43} = \dots = -h_{\pm 167} = -h_{\pm 197} = q^{-169/840} (6q + 17q^2 + 17q^3 + \dots)$ . A striking property of these coefficients is that they are so small: for instance, the largest value of  $c(n, \ell)$  in the table is  $c(12, 1) = 121$ , with corresponding discriminant  $\Delta = 4 \cdot 210 \cdot 12 - 1^2 = 10079$ , while for the Mathieu form  $h^{(2)}$  the table following eq. (7.31) gave the much larger coefficient 132825

---

<sup>19</sup>An alternative name would be “forms with only simple poles” because the forms  $h_\ell(\tau) = \sum_{\Delta} C(\Delta, \ell) q^{\Delta/4m}$  have no singularity worse than  $q^{-1/4m}$  as  $q \rightarrow 0$ , i.e., they have at most simple poles with respect to the local parameter  $q^{1/4m}$  at  $\tau = i\infty$ . However, this terminology would be dangerous because it could also suggest simple poles with respect to  $z$ . Another reasonable name would be “nearly holomorphic,” but this terminology also carries a risk of confusion since “almost holomorphic” is already a standard notation for modular forms which are polynomials in  $1/\tau_2$  with holomorphic coefficients.

$\ell$ with $h_\ell = h_{\ell_0}$	$\ell$ with $h_\ell = -h_{\ell_0}$	$n = 0$	1	2	3	4	5	6	7	8	9	10	11	12
1, 29, 41, 71	139, 169, 181, 209	-2	13	28	34	49	46	71	59	83	77	102	87	121
11, 31, 59, 101	109, 151, 179, 199	0	9	21	27	36	41	51	44	75	62	62	82	104
13, 43, 83, 97	113, 127, 167, 197	0	6	17	17	35	20	49	31	57	36	77	32	94
17, 53, 67, 73	137, 143, 157, 193	0	4	12	16	22	19	43	17	40	50	41	27	87
19, 61, 79, 89	121, 131, 149, 191	0	3	11	12	23	14	37	17	43	28	45	30	77
23, 37, 47, 103	107, 163, 173, 187	0	1	7	4	20	-1	32	3	30	10	50	-16	63

**Table 4:** Fourier coefficients of the function  $\mathcal{F}_{210}$

for the much smaller discriminant  $4 \cdot 2 \cdot 9 - 1^2 = 71$ . This behavior is a direct consequence of the “optimal growth” property together with the fact that the index is large (cf. Theorem 9.3 below and the discussion in §9.5).

In fact, in this example even more happens: not only the function  $\mathcal{Q}_{210} = \Phi_{2,210}^{-----}$ , but the whole function  $\Phi_{2,M}$  (and hence each of its eigencomponents  $\Phi_{2,210}^{\pm\pm\pm\pm}$ ) can be chosen to have optimal growth. It turns out that this statement holds for all square-free indices  $m$ , whether they have an even or an odd number of prime factors. If  $m$  is not square-free, then this is not true in general,<sup>20</sup> but it becomes true after a very small modification of the functions in question:

**Theorem 9.2.** *The functions  $\{\Phi_{2,m}\}_{m \geq 1}$  can be chosen to have the form*

$$\Phi_{2,m}(\tau, z) = \sum_{d^2|m} \Phi_{2,m/d^2}^0(\tau, dz) \quad (9.37)$$

where  $\{\Phi_{2,m}^0\}_{m \geq 1}$  are primitive mock Jacobi forms having optimal growth.

**Corollary.** *The functions  $\{\mathcal{Q}_M\}_{M > 1, \mu(M)=1}$  can be chosen to have optimal growth.*

(Here we have excluded the case  $M = 1$  because the function  $\mathcal{Q}_1 = -\mathcal{H}$  has already been chosen to have no poles at all.) Tables of Fourier coefficients of the functions  $\mathcal{Q}_M$  appearing in the corollary for  $M \leq 50$  are given in Appendix A.1.

If we denote by  $\mathcal{E}_m^{\text{OG}}$  the space of elliptic forms of index  $m$  (cf. §4.4) having optimal growth, then we have the sequence of inclusions

$$\mathcal{E}_m^0 \subset \mathcal{E}_m \subset \mathcal{E}_m^{\text{OG}} \subset \tilde{\mathcal{E}}_m \subset \mathcal{E}_m^! \quad \left( \varphi \in \mathcal{E}_m^{\text{OG}} \underset{\text{DEF}}{\iff} C_\varphi(\Delta, \ell) = 0 \text{ if } \Delta < -1 \right)$$

of spaces of different types of elliptic forms (cusp, holomorphic, optimal growth, weak, weakly holomorphic) with different orders of growth. Theorem 9.2 says that we can represent all of our

<sup>20</sup>It is true if and only if  $m$  is the product of a square-free number and a prime power.

functions in terms of new, “primitive,” forms  $\Phi_m^0 \in \mathcal{E}_m^{\text{OG}}$  for all  $m$ . The function  $\Phi_m^0$  is unique up to the addition of an element of the space  $J_{2,m}^{\text{OG}} = \mathcal{E}_m^{\text{OG}} \cap \tilde{J}_{2,m}$  of Jacobi forms of weight 2 and index  $m$  of optimal growth. We will study this latter space in the next subsection.

Theorem 9.2 has an immediate consequence for the growth of the Fourier coefficients of the optimal choice of  $\Phi_{2,m}$ , since the Hardy-Ramanujan circle method implies that the Fourier coefficients of a weakly holomorphic Jacobi (or mock Jacobi) form grow asymptotically like  $e^{\pi\sqrt{\Delta|\Delta_0|}/m}$ , where  $\Delta_0$  is the minimal value of the discriminants occurring in the form. We give a more precise version of this statement in the special case of forms of weight 2 having optimal growth.

**Theorem 9.3.** *Let  $\varphi$  be a weak or mock Jacobi form of weight 2 and index  $m$  with optimal growth. Then the Fourier coefficients of  $\varphi$  satisfy the asymptotic formula*

$$C_\varphi(\Delta, r) = \kappa_r e^{\pi\sqrt{\Delta}/m} + O(e^{\pi\sqrt{\Delta}/2m}) \quad (9.38)$$

as  $\Delta \rightarrow \infty$ , where the coefficient  $\kappa_r = \kappa(\varphi; r)$  is given by

$$\kappa_r = - \sum_{\substack{\ell \pmod{2m} \\ \ell^2 \equiv 1 \pmod{4m}}} C_\varphi(-1, \ell) \cos\left(\frac{\pi\ell r}{m}\right). \quad (9.39)$$

*Proof.* We only sketch the proof, since it is a standard application of the circle method. We write the coefficient  $c_\varphi(\Delta, r)$  as a Fourier integral  $\int_{-\frac{1}{2}+i\varepsilon}^{\frac{1}{2}+i\varepsilon} h_r(\tau) \mathbf{e}(-\Delta\tau/4m) d\tau$ , with  $h_\ell$  defined by (4.9) and (4.10). Its value is unchanged if we replace  $h_r(\tau)$  by its completion  $\widehat{h}_r(\tau)$ , since they differ only by terms with negative discriminant. The main contribution to the integral comes from a neighbourhood of  $\tau = 0$ , where we can estimate  $\widehat{h}_r(\tau)$  to high accuracy as

$$\widehat{h}_r(\tau) = -\frac{(i/\tau)^{3/2}}{\sqrt{2m}} \sum_{\ell \pmod{2m}} \mathbf{e}\left(\frac{r\ell}{2m}\right) \widehat{h}_\ell\left(-\frac{1}{\tau}\right) = -\frac{\kappa_r}{\sqrt{2m}} (i/\tau)^{3/2} \mathbf{e}\left(\frac{1}{4m\tau}\right) + O(\tau^{1/2}),$$

the first equation holding by the  $S$ -transformation equation ([52], p. 59) and the second because the “optimal growth” assumption implies that  $\widehat{h}_\ell(-1/\tau) = C_\varphi(-1, \ell) \mathbf{e}(1/4m\tau) + O(1)$  as  $\tau \rightarrow 0$  and the contribution from the Eichler integral of the shadow of  $\widehat{h}_\ell$  is at most  $O(1/\tau)$ . The rest follows by a standard calculation, the only point worthy of note being that since the  $h_\ell$ ’s have weight  $3/2$  the Bessel functions that typically arise in the circle method are simple exponentials here.  $\square$

It is interesting to compare the statement of Theorem 9.3 with the tables given in the Appendix. This will be discussed in more detail in §9.5.

## 9.4 Optimal choice of the function $\Phi_{2,m}$

As we already saw, Theorem 9.2 fixes the choice of the mock Jacobi form  $\Phi_{2,m}^0$  up to the addition of an element of the space  $J_{2,m}^{\text{OG}}$  of Jacobi forms of weight 2 and index  $m$  having optimal growth. The structure of this latter space is described by the following theorem, proved in §10.

**Theorem 9.4.** *The quotient  $J_{2,m}^{\text{OG}}/J_{2,m}$  has dimension 1 for every  $m$ . An element  $\mathcal{K}_m \in J_{2,m}^{\text{OG}}$  representing a non-trivial element of this quotient space can be chosen to satisfy*

$$\mathcal{K}_m(\tau, z) = y - 2 + y^{-1} + O(q) \in \mathbb{C}[y, y^{-1}][[q]],$$

and then has polar coefficients given by

$$C(\mathcal{K}_m; \Delta, r) = \begin{cases} 1 & \text{if } \Delta = -1, r^2 \equiv 1 \pmod{4m}, \\ 0 & \text{if } \Delta < -1. \end{cases} \quad (9.40)$$

The function  $\mathcal{K}_m$  can be chosen to be primitive and invariant under all the operators  $W_{m_1}$  ( $m_1 \parallel m$ ).

The choice of  $\mathcal{K}_m$  described in the theorem is unique up to the addition of a primitive holomorphic Jacobi form invariant under all  $W_m$  operators. This first occurs for  $m = 25$ , so  $\mathcal{K}_m$  is unique for small values of  $m$ . Because of Theorem 9.4, the function  $\Phi_m^0$  has only a very small amount of choice: we can change it without disturbing the “optimal growth” property by adding an arbitrary multiple of  $\mathcal{K}_m$  and an arbitrary holomorphic Jacobi form, but this is the only freedom we have. In particular, we have three special choices of “optimal”  $\Phi_{2,m}^0$ , each unique for  $m < 25$ , corresponding to the three that we already encountered for  $m = 6$  and  $m = 10$  in §9.2:

- (I) The  $(+, \dots, +)$ -eigencomponent of  $\Phi_{2,m}^{0,\text{I}}$  is strongly holomorphic.
- (II)  $\Phi_{2,m}^{0,\text{II}}$  has  $c(0, 1) = 0$ .
- (III)  $\Phi_{2,m}^{0,\text{III}}$  has  $c(0, 0) = 0$ .

Each of these three choices has certain advantages over the other ones. In case (I) we can choose  $\Phi_{2,m}^{0,\text{I}}$  within its class modulo  $J_{2,m}^0$  so that its  $(+, \dots, +)$ -eigencomponent is precisely equal to  $2^{1-\omega(m)}\mathcal{Q}_1|\mathcal{V}_{2,m}^{(1)}$  and hence has Fourier coefficients expressible in terms of class numbers, like in the examples for small  $m$  that we already saw. Also,  $\Phi_{2,m}^{0,\text{I}}$  is strongly holomorphic when  $m$  is a prime power, so that this is certainly the best of the three choices in that special case. The function occurring in (II) has a Fourier expansion beginning  $\frac{1}{12}(m+1) + O(q)$ , and the corresponding  $\psi_{m+1}^{0,\text{II}}$  satisfies (9.15) and belongs to the family of the weight zero, index  $m$  Jacobi forms discussed in [60]. The choice (III) looks at first sight like the least natural of the three,

since it might seem strange to insist on killing a  $\Delta = 0$  term when there are  $\Delta = -1$  terms that are more singular, but in fact it is the nicest in some ways, since it can be further improved by the addition of a holomorphic Jacobi form to have *all* of its  $\Delta = 0$  coefficients equal to zero, and its polar coefficients are then given by an elegant formula, which we now state.

**Theorem 9.5.** *The mock Jacobi form  $\Phi_{2,m}^{0,\text{III}}$  in Theorem 9.2 can be chosen, uniquely up to the addition of a Jacobi cusp form of weight 2 and index  $m$ , so that all its Fourier coefficients with  $\Delta = 0$  vanish. Its polar coefficients are then given by*

$$C(\Phi_{2,m}^{0,\text{III}}; -1, r) = \frac{1}{24} \sum_{d^2|m} \mu(d) \left( \left( \frac{r-1}{2}, \frac{m}{d^2} \right) + \left( \frac{r+1}{2}, \frac{m}{d^2} \right) \right) \quad \text{for } r^2 \equiv 1 \pmod{4m}. \quad (9.41)$$

Moreover, the function  $\Phi_{2,m}^{0,\text{III}}$  can be chosen to be primitive without affecting these properties.

This also gives a multiplicative formula for the polar coefficients of all eigencomponents of  $\Phi_{2,m}^{0,\text{III}}$ :

**Corollary.** *Write  $m = \prod_{i=1}^s P_i$ , where  $P_i = p_i^{\nu_i}$  with distinct primes  $p_i$  and exponents  $\nu_i \geq 1$ . Then for any  $\varepsilon_i \in \{\pm 1\}$  with  $\varepsilon_1 \cdots \varepsilon_s = 1$ , we have*

$$C\left(\Phi_{2,m}^{0,\text{III}} \mid \prod_{i=1}^s (1 + \varepsilon_i W_{P_i}); -1, r\right) = \frac{\pm 1}{12} \prod_{i=1}^s \begin{cases} (p_i + \varepsilon_i) & \text{if } \nu_i = 1, \\ (p_i^{\nu_i} - p_i^{\nu_i-2}) & \text{if } \nu_i \geq 2, \end{cases} \quad (9.42)$$

where the sign is the product of the  $\varepsilon_i$  for all  $i$  for which  $r \equiv -1 \pmod{2P_i}$ . In particular, for the functions  $\mathcal{Q}_M$  ( $M > 1$ ,  $\mu(M) = 1$ ), chosen by the Corollary to Theorem 9.2 to have optimal growth, the Fourier coefficients with non-positive discriminant are given by

$$C(\mathcal{Q}_M; \Delta, r) = \begin{cases} 0 & \text{if } \Delta = 0, \\ \pm \frac{\varphi(M)}{24} & \text{if } \Delta = -1, \end{cases} \quad (9.43)$$

where in the second line  $r^2 \equiv 1 \pmod{4M}$  and the sign is determined by  $\mu\left(\left(\frac{r+1}{2}, M\right)\right) = \pm 1$ .

The final statement of the corollary can be compared with the examples in (9.34), in which each of the  $\mathcal{F}_M$ 's occurring (for  $M > 1$ ) was normalized with  $C(\mathcal{F}_M; -1, 1) = -1$ , and the numerical factor relating  $\mathcal{Q}_M$  and  $\mathcal{F}_M$  equals  $-\varphi(M)/24$  in each case. To give a feeling for the nature of these functions in general, and as an illustration of the formulas in Theorem 9.5 and its corollary, we give in Table 5 below the first few Fourier coefficients  $C(\Phi; \Delta, \ell)$  for these and some related functions  $\Phi$  for  $m = 25$  (the first index where  $J_{2,m} \neq \{0\}$  and hence  $\mathcal{K}_m$  is not unique),  $m = 37$  (the first index where  $J_{2,m}^0 \neq \{0\}$  and hence  $\Phi_{2,m}^{0,\text{III}}$  is not unique),  $m = 50$  (the first case where (9.42) is non-trivial), and  $m = 91$  (the first case where where  $J_{2,m}^{0,-} \neq \{0\}$  and hence  $\mathcal{Q}_m$  is not unique). The free coefficients “ $a$ ”, “ $b$ ” and “ $c$ ” in the table arise because

$\Delta$	-1	0		4	11	16	19	24	31	
$\pm\ell \pmod{50}$	1	0	10	20	14	17	22	9	24	13
$\mathcal{K}_{25}$	1	-2	$a$	$1-a$	$1-a$	$1+2a$	$2+3a$	$2-4a$	$5-2a$	$3+2a$
$\Phi_{2,25}^{0,\text{III}}$	1	0	0	0	0	1	2	-1	2	1
$\Phi_{2,25}^{0,\text{I}}$	0	2	$c$	$-1-c$	$-1-c$	$2c$	$3c$	$-3-4c$	$-3-2c$	$-2+2c$

$\Delta$	-1	0	3	4	7	11	12	16	27	28
$\pm\ell \pmod{74}$	1	0	21	12	17	27	32	24	11	34
$\mathcal{K}_{37}$	1	-2	$a$	$a$	$1-a$	$1+a$	$2-a$	$2-2a$	$1-3a$	$2+3a$
$\Phi_{2,37}^{0,\text{III}}$	$\frac{19}{12}$	0	$b$	$-\frac{1}{6}+b$	$\frac{1}{4}-b$	$\frac{11}{12}+b$	$\frac{3}{2}-b$	$1-2b$	$-\frac{3}{4}-3b$	$\frac{13}{6}+3b$
$\Phi_{2,37}^{0,\text{I}}$	0	$\frac{19}{6}$	$c$	$-\frac{1}{6}+c$	$-\frac{4}{3}-c$	$-\frac{2}{3}+c$	$-\frac{5}{3}-c$	$-\frac{13}{6}-2c$	$-\frac{7}{3}-3c$	$-1+3c$

$\Delta$	-1	0		4	16	24	31		36	39		
$\pm\ell \pmod{100}$	1	49	0	20	40	14	28	24	13	37	42	19
$\mathcal{K}_{50}$	1	1	-2	$a$	$1-a$	$-a$	$1+a$	$1-2a$	$1+2a$	$1+2a$	$2+3a$	$-2a$
$\Phi_{2,50}^{0,\text{III}} (1+W_2)$	3	3	0	0	0	-2	2	-2	3	3	8	-1
$\Phi_{2,50}^{0,\text{III}} (1-W_2)$	1	-1	0	0	0	0	0	0	-1	1	0	-1

$\Delta$	-1	3	12	27	40	48	55	68	75
$\pm\ell \pmod{182}$	$1/27$	$19/33$	$38/66$	$57/83$	$18/60$	$50/76$	$71/85$	$32/46$	$17/87$
$\mathcal{Q}_{91}$	3	$a$	$-1+a$	$-1-2a$	$-1-3a$	$-2+2a$	$3a$	$1+3a$	$-5+2a$

**Table 5:** Examples of non-unique Jacobi and mock Jacobi forms

$\dim J_{2,25} = \dim J_{2,37}^0 = \dim J_{2,50} = \dim J_{2,91}^0 = 1$ . In the case of  $m = 91$  we show the two  $\ell$ -values corresponding to each value of  $\Delta$  in the format  $\ell/\ell^*$  and give in the following line the Fourier coefficient  $C(\mathcal{Q}_{91}; \Delta, \ell) = -C(\mathcal{Q}_{91}; \Delta, \ell^*)$ . In the cases of  $m = 25$  and  $m = 37$ , we have given the (non-unique) holomorphic mock Jacobi form  $\Phi_{2,2m}^{0,\text{I}}$  as well as  $\Phi_{2,2m}^{0,\text{III}}$ .

In summary, the ‘‘optimal growth’’ property of  $\Phi_{2,m}^0$  and the near-uniqueness of the Jacobi form  $\mathcal{K}_m$  have permitted us to pin down both of these functions in many cases, but there still remains an additive ambiguity whenever there are Jacobi cusp forms. We do not know how to resolve this ambiguity in general, and state this as an open problem:

*Question:* Is there a canonical choice of the forms  $\mathcal{K}_m$  and  $\mathcal{Q}_M$  for all positive integers  $m$  and all positive square-free integers  $M$ ?

Should the answer to either of these questions be positive, the corresponding forms would be of considerable interest. In particular, one can speculate that the canonical choice of  $\mathcal{Q}_M$ , if it exists, might be canonically associated to the quaternion algebra of discriminant  $M$ , since there is a natural bijection between positive square-free integers and quaternion algebras over  $\mathbb{Q}$ , with



the latter being indefinite or indefinite according as  $\mu(M) = 1$  (corresponding to our  $k = 2$  forms and to mock theta functions of weight  $3/2$ ) or  $\mu(M) = -1$  (corresponding to  $k = 1$  and to mock theta functions of weight  $1/2$ ).

We end this subsection by describing one possible approach to answering the above question that we tried. This approach was not successful in the sense that, although it did produce a specific choice for  $\mathcal{K}_m$  and  $\Phi_{2,m}^0$  (or  $\Phi_{1,m}^0$ ), these did not seem to have arithmetic coefficients and therefore had to be rejected, but the method of calculation involves a result that is of interest in its own right and may be useful in other contexts. The idea is to fix the choice of the Jacobi forms (or mock Jacobi forms) in question by demanding that they (or their completions) be orthogonal to Jacobi cusp forms with respect to the Petersson scalar product. (As an analogy, imagine that one wanted to determine the position of  $E_{12} = 1 + \dots$  in  $M_{12} = \mathbb{C}E_4^3 + \mathbb{C}\Delta$  but did not know how to compute the Fourier expansion of  $\sum(m\tau + n)^{-12}$ . Then one could simply write  $E_{12} = E_4^3 + a\Delta$  and use the orthogonality of Eisenstein series and cusp forms to compute the number  $a = -432000/691$  numerically as  $-(E_4^3, \Delta)/(\Delta, \Delta)$ , where  $(\ , \ )$  denotes the Petersson scalar product.) To apply this idea, we need a good way to compute the Petersson scalar product of Jacobi forms. This is the result of independent interest referred to above. For simplicity, we state it only in the case  $k = 2$ .

**Theorem 9.6.** *Let  $\varphi_1$  and  $\varphi_2$  be two Jacobi cusp forms of weight 2 and index  $m$ , and set*

$$A(\Delta) = \sum_{\ell \pmod{4m}} C_{\varphi_1}(\Delta, \ell) \overline{C_{\varphi_2}(\Delta, \ell)}, \quad H(t) = \sum_{\Delta=1}^{\infty} \frac{A(\Delta)/\sqrt{\Delta}}{\exp(2\pi\sqrt{t\Delta/m}) - 1}. \quad (9.44)$$

*Then the function  $\frac{tH(t) - H(1/t)}{t - 1}$  ( $t \neq 1$ ) has a constant value  $c_H$  which is proportional<sup>21</sup> to the Petersson scalar product  $(\varphi_1, \varphi_2)$ .*

*Sketch of proof:* We refer to [52] for the definition of  $(\varphi_1, \varphi_2)$  (eq. (13), p. 27) and for its expression as a simple multiple of  $\sum_{\ell}(h_{\ell}^{(1)}, h_{\ell}^{(2)})$ , where  $h_{\ell}^{(1)}$  and  $h_{\ell}^{(2)}$  are the coefficients in the theta expansions of  $\varphi_1$  and  $\varphi_2$  (Theorem 5.3, p. 61). Calculating these weight  $3/2$  scalar products by the Rankin-Selberg method as explained in §3.1, we find that  $(\varphi_1, \varphi_2)$  can be expressed as a simple multiple of the residue at  $s = 3/2$  of the Dirichlet series  $L(s) = \sum_{\Delta>0} A(\Delta)\Delta^{-s}$ . The Rankin-Selberg method also shows that the function  $\tilde{L}(s) = 2(2\pi)^{-s}m^s\Gamma(2s)\zeta(2s)L(s + \frac{1}{2})$  has a meromorphic continuation to all  $s$ , with poles only at  $s = 0$  and  $s = 1$ , and is invariant under  $s \rightarrow 1 - s$ . But  $\tilde{L}(s)$  is simply the Mellin transform of  $H(t)$ , so this implies the proposition by a standard argument.  $\square$

A rather surprising aspect of this theorem is that it is actually *easier* to compute Petersson scalar products for Jacobi forms than it is for ordinary modular forms of integral weight. In

<sup>21</sup>The exact constant of proportionality plays no role for our purposes, since we will only be concerned with ratios of Petersson scalar products. If the scalar product is normalized as in [52], its value is  $1/2$ .

that case the analogue of  $H(t)$  would have an infinite sum of  $K$ -Bessel functions multiplying each coefficient  $A(\Delta)$ , whereas here the Bessel functions are replaced by simple exponentials (because the modular forms  $h_\ell^{(i)}$  have half-integral weight, so that the usual gamma factor  $\Gamma(s)\Gamma(k+s-1)$  in the Rankin-Selberg method becomes a single gamma function  $\Gamma(2s)$ ) and at the same time the infinite sum becomes a geometric series that can be summed explicitly.

To apply the proposition to our situation in the case of  $\mathcal{K}_{37}$ , we start with an initial choice (like the one given in Table 5) and replace it by  $\mathcal{K}_{37} + a\varphi_{37}$ , where  $\varphi_{37}$  is the generator of  $J_{2,37}^0$  (normalized by  $C(\varphi_{37}; 3, 21) = 1$ ) and where  $a \in \mathbb{R}$  is chosen to make this function orthogonal to  $\varphi_{37}$ , i.e.,  $a = -(\mathcal{K}_{37}, \varphi_{37})/(\varphi_{37}, \varphi_{37})$ , in the hope that this  $a$  will be an integer or simple rational number. (Of course  $\mathcal{K}_{37}$  is not a cusp form, but the integral defining the Petersson scalar product still converges,<sup>22</sup> the sum defining  $H(t)$  converges for  $t > \frac{1}{148}$  by Theorem 9.3, and Theorem 9.6 still holds.<sup>23</sup>) The numerical calculation for the case  $\varphi_1 = \varphi_2 = \varphi_{37}$  gives the value  $c_H = c_1 = -0.95284748097797917403$  (a value that can be checked independently, since up to a factor  $-\pi$  it must be equal to the even period of the elliptic curve  $y^2 - y = x^3 - x$ , as calculated by GP-PARI as `ellinit([0,0,1,-1,0])[15]`). The corresponding number for  $\varphi_1 = \varphi_{37}$  and  $\varphi_2 = \mathcal{K}_{37}$  (with  $\mathcal{K}_{37}$  chosen as in Table 5, with  $a = 0$ ) is  $c_2 = 0.26847600706319893417$ , which is not a simple multiple of  $c_1$ , but when we calculate again with  $\varphi_2 = \Phi_{2,37}^{0,III}$  (again as in Table 5, with  $b = 0$ ) we find a value  $c_3 = 0.107471184190738587618$  that is related to the two previous values by  $12c_3 - 19c_2 = 4c_1$  numerically to high precision. Thus in this case we have indeed found a form orthogonal to cusp forms and with rational coefficients. This at first looks promising, but in fact has a simple and disappointing explanation: the linear combination  $\Phi_{2,37}^{0,III} - \frac{19}{12}\mathcal{K}_{37} - \frac{1}{3}\varphi_{37} = \Phi_{2,37}^{0,I} - \frac{1}{3}\varphi_{37}$  is nothing other than  $\mathcal{Q}_1|V_{37}$  (as one can see in Table 5, where setting  $c = -\frac{1}{3}$  gives entries equal to  $-H(\Delta)$ ), and since the function  $\mathcal{Q}_1 = -\mathcal{H}$  is constructed from an Eisenstein series, albeit a non-holomorphic one, it is automatically orthogonal to all cusp forms. And indeed, when we look at the case of  $\mathcal{Q}_{91}$  and  $\varphi_{91} \in J_{2,91}^{0,-}$ , where there are no Eisenstein series in sight, then the calculation of the quotient  $(\widehat{\mathcal{Q}}_{91}, \varphi_{91})/(\varphi_{91}, \varphi_{91})$  failed to yield a simple number. (The non-holomorphy of  $\widehat{\mathcal{Q}}_{91}$  does not affect the application of Theorem 9.6, since its non-holomorphic part has Fourier coefficients only for  $\Delta < 0$ , where the coefficients of  $\varphi_{91}$  vanish.) As an incidental remark, we mention that the cusp form  $\varphi_{37}$ , whose first few

<sup>22</sup>The precise condition needed for the convergence of the Petersson scalar product of two weakly holomorphic Jacobi forms  $\varphi_1$  and  $\varphi_2$  is that  $\text{ord}_\infty(h_\ell^{(1)}) + \text{ord}_\infty(h_\ell^{(2)}) > 0$  for all  $\ell \in \mathbb{Z}/2m\mathbb{Z}$ . This is amply satisfied in the case  $\varphi_1 = \varphi_{37}$ ,  $\varphi_2 = \mathcal{K}_{37}$ , since the only negative value of  $\text{ord}_\infty(h_\ell^{(2)})$  is  $-1/148$  for  $\ell \equiv \pm 1 \pmod{74}$ , and there  $\text{ord}_\infty(h_\ell^{(1)}) = 147/148$ .

<sup>23</sup>The proof given above breaks down, since the numbers  $A(\Delta)$  grow exponentially in  $\sqrt{\Delta}$ , so that the Dirichlet series  $\sum_\Delta A(\Delta)/\Delta^s$  does not converge for any value of  $s$ , but the function  $f(y) = y^{3/2} \sum_\Delta A(\Delta)e^{-\pi\Delta y/m}$ , which is the constant term of an  $SL(2, \mathbb{Z})$ -invariant function that is small at infinity, still has a Mellin transform with a meromorphic continuation to all  $s$ , and if we define  $\widetilde{L}(s)$  as the product of this Mellin transform with  $\pi^{-s}\Gamma(s)\zeta(2s)$ , then  $\widetilde{L}(s)$  has the same analytic properties as before and its inverse Mellin transform  $H(t)$  is still given by the sum in (9.44) for  $t$  sufficiently large and satisfies the same functional equation.

Fourier coefficients were obtained only with difficulty in [52] (pp. 118–120 and 145), can now be obtained very easily using the “theta blocks” of [63] as

$$\varphi_{37}(\tau, z) = \frac{1}{\eta(\tau)^6} \prod_{i=1}^{10} \vartheta(\tau, a_i z), \quad (a_1, \dots, a_{10}) = (1, 1, 1, 2, 2, 2, 3, 3, 4, 5),$$

and similarly  $\varphi_{2,91}$  can be obtained easily as the difference of the two similarly defined theta blocks for the 10-tuples  $(a_1, \dots, a_{10}) = (1, 1, 2, 3, 4, 4, 5, 5, 6, 7)$  and  $(1, 1, 2, 2, 3, 3, 4, 5, 7, 8)$ .

## 9.5 Observations on the weight one family, integrality, and positivity

In the last two subsections we discussed experimental properties of the family of weight 2 mock Jacobi forms that were introduced in §9.1 and formulated a number of general results that will be proved in §10. In this subsection we report on the experimental data for the weight 1 family and discuss various properties of both families that play a role in connection with other investigations in mathematics and physics (like “Umbral Moonshine”) or that seem to be of independent interest.

The discussion of the weight 1 family is quite different in nature from that of the weight 2 case, because here there do not seem to be general structural results as there were there, but only a small number of special functions for small values of the index having nice properties. In particular, the analogue of Theorem 9.2 no longer holds, and we find instead that (at least up to  $m = 100$ , and presumably for all  $m$ ) the appropriately defined primitive mock Jacobi forms  $\Phi_{1,m}^0$  can be chosen to have optimal growth only for the indices

$$m = 2, 3, 4, 5, 6, 7, 8, 9, 10, 13, 16, 18, 25. \quad (9.45)$$

Similarly, although the analogue of Theorem 9.1 is still valid in the weight 1 case (now with the special mock Jacobi forms  $\mathcal{Q}_M$  labelled by square-free integers  $M$  having an odd rather than an even number of prime factors), the functions  $\mathcal{Q}_M$  can apparently be chosen to have optimal growth only for

$$M = 2, 3, 5, 7, 13, 30, 42, 70, 78. \quad (9.46)$$

(This list has a precise mathematical interpretation; see §3. below.) Thus here a small number of special examples is singled out. Interestingly enough, these turn out to include essentially all of the special mock theta functions that have been studied in the literature in the past.

For convenience, we divide up our discussion into several topics.

1. *Optimal choices for the weight one family*

As in the weight 2 case, we begin our discussion by looking at small values of  $m$ . We will consider the original forms  $\Phi_{1,m}$  first, starting with the “standard” choices (9.2) and trying to modify them by the addition of weak Jacobi forms to attain the property of having optimal growth. Actually, it is better to look at the primitive mock Jacobi forms, now defined by

$$\Phi_{1,m}^0 \equiv \sum_{d^2|m} \frac{\mu(d)}{d} \Phi_{1,m/d^2}|U_d, \quad \Phi_{1,m} \equiv \sum_{d^2|m} \frac{1}{d} \Phi_{1,m/d^2}^0|U_d \pmod{\tilde{J}_{1,m}}, \quad (9.47)$$

instead of (9.37) and (10.52), for the same reason for which we introduced the functions  $\Phi_{2,m}^0$  in §9.3: residue arguments similar to those that we will give in §10 for the weight 2 case show here that  $\Phi_{1,m}$  can never have optimal growth unless it agrees with  $\Phi_{1,m}^0$  (which happens if and only if  $m$  is square-free or the square of a prime number), so that the latter class is certain to be more productive. Finally, we will look at the forms  $\mathcal{Q}_M$  where  $M$  is a product of an odd number of distinct prime factors, since according to Theorem 9.1 all of the forms  $\Phi_{1,m}$  can be constructed from these. Unlike the situation for  $k = 2$ , here we do not have to worry about any choices, because of the following theorem (to be compared with Lemma 2.3 of [21] and to be proved, like all its companions, in §10) and corollary. The only issue is therefore whether the “optimal growth” condition can be fulfilled at all.

**Theorem 9.7.** *There are no weight one weak Jacobi forms of optimal growth.*

**Corollary.** *If any of the mock Jacobi forms  $\Phi_{1,m}$ ,  $\Phi_{1,m}^0$ , or  $\mathcal{Q}_M$  ( $\mu(M) = -1$ ) can be chosen to have optimal growth, then that choice is unique.*

We now look at some small individual indices. For  $\mathbf{m}=\mathbf{1}$  there is nothing to discuss, since the function  $\Phi_{1,1}^{\text{st}} = (C/2A)^F$  vanishes identically, as already explained in §8.5 (Example 1). (This, by the way, is the reason why  $\Phi_{1,m} = \Phi_{1,m}^0$  when  $m$  is the square of a prime.) The case of  $\mathbf{m}=\mathbf{2}$  has also already been treated in the same place (Example 2), where we saw that

$$\varphi_{1,2}^{\text{opt}} = \frac{B}{24A} C, \quad \Phi_{1,2}^{\text{opt}} = -\frac{1}{24} \mathcal{F}_2, \quad (9.48)$$

with  $\mathcal{F}_2$  the mock modular form defined in §7 and related to the Mathieu moonshine story. For  $\mathbf{m}=\mathbf{3}$ , we find that the (unique) choice of  $\Phi_{1,m}$  having optimal growth is given by

$$\varphi_{1,3}^{\text{opt}} = \frac{B^2 - E_4 A^2}{288A} C, \quad \Phi_{1,3}^{\text{opt}} = \varphi_{1,3}^{\text{opt}} - \mathcal{A}_{1,3} = \frac{1}{12} \mathcal{F}_3, \quad (9.49)$$

where the normalizing factor  $1/12$  has been chosen to give  $\mathcal{F}_3$  integral Fourier coefficients. These coefficients are determined by virtue of their periodicity and oddness by the values  $C^{(3)}(\Delta) = C(\mathcal{F}_3; \Delta, r)$  for  $r \in \{1, 2\}$ , the first few values of  $C^{(3)}(\Delta)$  being given by

$\Delta$	-1	8	11	20	23	32	35	44	47	56	59
$C^{(3)}(\Delta)$	-1	10	16	44	55	110	144	280	330	572	704

(A more extensive table of these coefficients, listed separately for the two values  $r = 1$  and  $r = 2$ , is given in Appendix A.2, where we also give the corresponding values for the other forms treated here, so that we no longer will give tables of coefficients in the text.) For  $\mathbf{m}=4$  and  $\mathbf{m}=5$ , the functions  $\Phi_{1,m}$  can again be chosen (of course uniquely, by the corollary above) to have optimal growth, the corresponding meromorphic Jacobi forms  $\varphi_{1,4}^{\text{opt}}$  and  $\varphi_{1,5}^{\text{opt}}$  being given by

$$\frac{B^3 - 3E_4A^2B + 2E_6A^3}{2 \cdot 12^2 A} C \quad \text{and} \quad \frac{B^4 - 6E_4A^2B^2 + 8E_8A^3B - 3E_4^2A^4}{2 \cdot 12^3 A} C, \quad (9.50)$$

respectively. Note that in the case of  $m = 4$ , according to the decomposition result given in Theorem 9.1, we could also make the choice  $\mathcal{Q}_2|\mathcal{V}_{1,2}^{(2)}$  for the function  $\Phi_{1,4}$ , but then it would have the worse discriminant bound  $\Delta_{\min} = -4$ . We also observe that, although 4 is not square-free, we do not have to distinguish between  $\Phi_{1,m}$  and  $\Phi_{1,m}^0$  here, because they differ by the function  $\Phi_{1,1}^0|U_2$  and the function  $\Phi_{1,1}$  vanishes identically.

It is now very striking that the numerators of the fractions in (9.49) and (9.50) are the *same* as the expressions that we already saw in eqs. (9.10), (9.11) and (9.12) in §9.2, meaning that for these values of  $m$  we have

$$\varphi_{1,m}^{\text{opt}} = \frac{C}{A} \psi_{0,m-1}^{\text{opt}} = C \varphi_{2,m-2}^{\text{opt}} \quad (9.51)$$

with the same weight 0 weak Jacobi modular forms  $\psi_{0,m-1}^{\text{opt}}$  as were tabulated in Table 2. The same thing happens for several other small values of the index. We do not know the deeper reason (if any) for this coincidence, but merely remark that the weight 0 weak Jacobi forms obtained from the weight 1 meromorphic Jacobi forms in our family by dividing by  $C$  also play a prominent role in the recent work of Cheng, Duncan and Harvey [21] on “umbral moonshine,” and also that the forms  $\psi_{0,m}^{\text{opt}}$  occur in a somewhat different context in earlier work of Gritsenko[60] related to Borcherds products, as was already mentioned in §9.3 when we introduced the “second standard choice”  $\Phi_{2,m}^{0,\text{II}}$  of the primitive mock Jacobi form  $\Phi_{2,m}^0$ .

Continuing to larger values, we look first at primes, since for  $m$  prime the three functions of interest  $\Phi_{1,m}$ ,  $\Phi_{1,m}^0$  and  $\mathcal{Q}_m$  all coincide. For  $\mathbf{m}=7$  we find that  $\mathcal{Q}_7 = \Phi_{1,7}^{\text{opt}}$  again has optimal growth and again satisfies (9.51) with  $\varphi_{0,6}^{\text{opt}}$  as given in Table 2. However, for  $\mathbf{m}=11$  the function  $\mathcal{Q}_M$  *cannot* be chosen to be of optimal growth; the best possible choice, shown in the table A.3 in the appendix, has minimal discriminant  $-4$ . For  $\mathbf{m}=13$  the function  $\mathcal{Q}_{13}$  again has optimal growth and is again given by (9.51), but for larger primes this never seems to happen again, the minimal value of  $\Delta$  for the optimal choices of  $\mathcal{Q}_M$  for the first few prime indices being given by

$M$	2	3	5	7	11	13	17	19	23	29	31	37
$\Delta_{\min}$	1	1	1	1	4	1	4	4	8	5	8	4

The next simplest case is  $m = p^2$ , since here  $\Phi_{1,m}$  is still primitive. For  $\mathbf{m}=9$  and  $\mathbf{m}=25$  this function can be chosen to have optimal growth, but apparently not after that. The following

case to look at is  $m = p^\nu$  with  $\nu \geq 2$ , since here the eigendecomposition (9.31) of  $\Phi_{1,m}$  is trivial. (The eigenvalue of  $W_m$  must be  $-1$  because the weight is odd, so  $\Phi_{1,m} = \Phi_{1,m}^- = \mathcal{Q}_p | \mathcal{V}_{p^{\nu-1}}^{(p)}$ .) Here we find optimal growth only for  $\mathbf{m}=8$  and  $\mathbf{m}=16$ . Next, for  $M$  square-free but not prime, where the decomposition (9.2) is trivial, the only three cases giving optimal growth seem to be  $\mathbf{m}=6$ ,  $\mathbf{m}=10$  and  $\mathbf{m}=18$ . And finally, for  $M$  neither square-free nor a power of a prime we found no examples at all. Altogether this leads to the list (9.45) given above. The first few coefficients of the forms  $\Phi_{1,m}^0$  for all of these values of  $m$  are tabulated in Appendix A.2.

However, this is not the end of the story. If  $\Phi_{1,m}$  has optimal growth for some  $m$ , then of course all of its eigencomponents with respect to the Atkin-Lehner operators also do, but the converse is not true, so there still can be further square-free values of  $M$  for which the function  $\mathcal{Q}_M = 2^{1-\omega(M)} \Phi_{1,M}^{-,\dots,-}$  has optimal growth, and this indeed happens. Since primes have already been treated and products of five or more primes are beyond the range of our ability to compute, we should consider products of three primes. Here we find four examples, for the indices

$$30 = 2 \cdot 3 \cdot 5, \quad 42 = 2 \cdot 3 \cdot 7, \quad 70 = 2 \cdot 5 \cdot 7, \quad 78 = 2 \cdot 3 \cdot 13,$$

but for no other values less than 200 (and probably for no other values at all), giving the list (9.46). Again the Fourier coefficients of the corresponding forms, as well as those for all other  $M < 150$  of the form  $p_1 p_2 p_3$ , have been listed in the appendix (part A.3). These examples are of particular interest, for a reason to which we now turn.

## 2. Relationship to classical mock theta functions

Recall from §7.1 that a mock modular form is called a mock theta function if its shadow is a unary theta series, and that this definition really does include all of Ramanujan's original mock theta functions that began the whole story. The theta expansion coefficients  $h_\ell(\tau)$  of our special mock Jacobi forms  $\Phi_{1,m}$  and  $\Phi_{2,m}$  are always mock theta functions, since their shadows are multiples of the unary theta series  $\varphi_{m,\ell}^1$  and  $\varphi_{m,\ell}^0$ , respectively. In fact, it turns out that many of our examples actually coincide with the mock theta functions studied by Ramanujan and his successors. This is the "particular interest" just mentioned.

Let us consider first the case  $\mathbf{m}=30$ , the first of our  $\mathcal{Q}_M$  examples with composite  $M$ . Because it is so highly composite, there are very few orbits of the group of Atkin-Lehner involutions on the group of residue classes modulo  $2m$  and hence very few components in the theta decomposition of  $\mathcal{Q}_{30}$ . In fact there are only two, so that up to sign we get only two distinct coefficients  $h_1$  and  $h_7$  and  $\mathcal{Q}_{30}$  has a decomposition

$$\begin{aligned} -3 \mathcal{Q}_{30} = & h_1 (\vartheta_{30,1} + \vartheta_{30,11} + \vartheta_{30,19} + \vartheta_{30,29} - \vartheta_{30,31} - \vartheta_{30,41} - \vartheta_{30,49} - \vartheta_{30,59}) \\ & + h_7 (\vartheta_{30,7} + \vartheta_{30,13} + \vartheta_{30,17} + \vartheta_{30,23} - \vartheta_{30,37} - \vartheta_{30,43} - \vartheta_{30,47} - \vartheta_{30,53}). \end{aligned}$$

(The normalizing factor  $-3$  is included for convenience: the function  $-3\mathcal{Q}_{30}$  is the function  $\mathcal{F}_{30}$  defined in 9.52 below.) The mock theta functions  $h_1$  and  $h_7$  have Fourier expansions beginning

$$\begin{aligned} h_1(\tau) &= q^{-1/120} (-1 + q + q^2 + 2q^3 + q^4 + 3q^5 + \cdots), \\ h_7(\tau) &= q^{71/120} (1 + 2q + 2q^2 + 3q^3 + 3q^4 + 4q^5 + \cdots). \end{aligned}$$

These expansions are easily recognized: up to trivial modifications they are two of the “mock theta functions of order 5” in the original letter of Ramanujan to Hardy, namely

$$\begin{aligned} 2 + q^{1/120} h_1(\tau) &= \chi_1(q) = \sum_{n=0}^{\infty} \frac{q^n}{(1 - q^{n+1}) \cdots (1 - q^{2n})}, \\ q^{-71/120} h_2(\tau) &= \chi_2(q) = \sum_{n=0}^{\infty} \frac{q^n}{(1 - q^{n+1}) \cdots (1 - q^{2n+1})}. \end{aligned}$$

(Actually, Ramanujan gave five pairs of mock theta functions of “order 5” in his letter, any two differing from one another by weakly holomorphic modular forms, but it was pointed out in [118] that of the five associated pairs of mock modular forms only  $(q^{-1/120}(2 - \chi_1(q)), q^{71/120}\chi_2(q))$  has the property that its completion transforms as a vector-valued modular form for the full modular group.) For  $\mathbf{m}=\mathbf{42}$ , exactly the same thing happens, and the three distinct components  $h_1, h_5$  and  $h_{11}$  of  $\mathcal{Q}_{42}$ , whose initial coefficients are tabulated in Appendix A.3, turn out to be, up to powers of  $q$  and signs, identical with Ramanujan’s three “order 7” mock theta functions  $\mathcal{F}_{7,i}$  ( $i = 1, 2, 3$ ) that were given as our first examples of mock modular forms in §7.1. What’s more, this behavior continues: in [118] it was mentioned that one could construct mock theta functions of arbitrary prime<sup>24</sup> “order”  $p$  generalizing these two examples as quotients by  $\eta(\tau)^3$  of certain weight 2 indefinite theta series, and an explicit example

$$M_{11,j}(\tau) = \frac{1}{\eta(\tau)^3} \sum_{\substack{m > 2|n|/11 \\ n \equiv j \pmod{11}}} \binom{-4}{m} \binom{12}{n} \left(m \operatorname{sgn}(n) - \frac{n}{6}\right) q^{m^2/8 - n^2/264} \quad (j \in \mathbb{Z}/11\mathbb{Z})$$

was given for  $p = 11$ . Computing the beginnings of the Fourier expansions of these five mock theta functions (only five rather than eleven because  $M_{11,j} = -M_{11,-j}$ ) and comparing with the table in Appendix A.3, we find perfect agreement with the coefficients in the theta expansion of  $\mathcal{Q}_{66}$ , the first example in that table of a form that is not of optimal growth.

This discussion thus clarifies the precise mathematical meaning of the number designated by Ramanujan as the “order” of his mock theta functions, but which he never defined: it is indeed related to the *level* of the completions of the corresponding mock modular forms, as was clear from the results of Zwegers and other authors who have studied these functions, but it

---

<sup>24</sup>actually, only  $(p, 6) = 1$  is needed



is even more closely related to the *index* of the mock Jacobi forms that have these mock theta functions as the coefficients of its theta expansion.

Of course the above cases are not the only ones where there is a relationship between the special mock Jacobi forms  $\mathcal{Q}_M$  and classically treated mock theta functions, but merely the ones where the relationship between the “order” and the index is clearest. Without giving any further details, we say only that Ramanujan’s order 3 functions are related in a similar way<sup>25</sup> to our  $\mathcal{Q}_3$  and that his order 10 functions (not given in his letter to Hardy, but contained in the so-called “Lost Notebook”) are related to  $\mathcal{Q}_5$ . We also refer once again to the beautiful recent work of Cheng, Duncan and Harvey [21] on “umbral moonshine,” in which many of these same functions are discussed in detail in connection with representations of special finite groups.

Finally, we should mention that all of Ramanujan’s functions were  $q$ -hypergeometric series but that in general no  $q$ -hypergeometric expression for the theta expansion coefficients of the functions  $\Phi_{k,m}$  or  $\mathcal{Q}_M$  is known. This seems to be an interesting subject for further research. One could look in particular for representations of this sort for the functions  $\mathcal{Q}_{13}$  and  $\Phi_{1,25}^0$  as given in Table A.2, since these have particularly small and smoothly growing coefficients that suggest that such a representation, if it exists, might not be too complicated. It would also be reasonable to look at  $\mathcal{Q}_{70}$  and  $\mathcal{Q}_{78}$ , the only two other known weight 1 examples with optimal growth.

### 3. Integrality of the Fourier coefficients

In Subsections 9.2 and 9.3 and above, we encountered several examples of functions  $\mathcal{Q}_M$  of optimal growth that after multiplication by a suitable factor gave a function  $\mathcal{F}_M$  having all or almost all of the following nice arithmetic properties:

- (A) All Fourier coefficients  $C(\mathcal{F}_M; \Delta, r)$  are integral.
- (B) The polar Fourier coefficient  $C(\mathcal{F}_M; -1, 1)$  equals  $-1$ .
- (C) The non-polar Fourier coefficients  $C(\mathcal{F}_M; \Delta, r)$  have a sign depending only on  $r$ .
- (D)  $C(\mathcal{F}_M; \Delta, r)$  is always positive for  $r = r_{\min}$ , the smallest positive number whose square equals  $-\Delta \pmod{4m}$ .

In particular, for  $k = 2$  this occurred for  $M = 6, 10, 14$  and  $15$  (compare eqs. (7.28), (9.22), (9.29) and (9.30), which describe the coefficients  $C^{(M)}(\Delta) = C(\mathcal{F}_M; \Delta, r_{\min})$  for these indices) and for  $k = 1$ , for the values  $M = 2, 3, 5, 7, 13$  and  $30$  (with “non-negative” instead of “positive” in (C) in the case of  $M = 13$ ), as well as for the functions  $\Phi_{1,m}^0$  for the prime powers  $M = 4, 8, 9, 16$  and  $25$  (again with “positive” replaced by “non-positive,” and also allowing half-integral values for the Fourier coefficients with  $\Delta = 0$ ). It is natural to ask for what values

---

<sup>25</sup>but not “on the nose,” for instance, our function  $h_{3,1}$  differs from Ramanujan’s 3rd order mock theta function  $-3q^{1/12}f(q^2)$  by an eta product.



of  $M$  these properties can be expected to hold. We will discuss properties (A) and (B) here and properties (C) and (D) in point 4. below.

Clearly either (A) or (B) can always be achieved by multiplying  $\mathcal{Q}_M$  by a suitable scalar factor. We define a renormalized function  $\mathcal{F}_M$  in all cases by setting

$$\mathcal{F}_M(\tau, z) = -c_M \mathcal{Q}_M(\tau, z), \tag{9.52}$$

where  $c_M$  is the unique positive rational number such that  $\mathcal{F}_M$  has integer coefficients with no common factor. Then (A) is always true and (B) and (C) will hold in the optimal cases. This normalization is also convenient for the presentation of the coefficients in tables, since there are no denominators and very few minus signs. Tables of the first few Fourier coefficients of all  $\mathcal{F}_M$  up to  $M = 50$  in the weight 2 case (i.e., for  $M$  with  $\mu(M) = 1$ ) and of all known  $\mathcal{F}_M$  having optimal growth in the weight 1 case (i.e., for  $M$  with  $\mu(M) = -1$ ) are given in A.1 and A.3 of the Appendix, respectively. Looking at these tables, we see that the values of  $M$  for which all three properties above hold are  $M = 6, 10, 14, 15, 21$  and  $26$  in the case  $k = 2$  and  $M = 2, 3, 5, 7, 13, 30$  and  $42$  in the case  $k = 1$ , and that in the  $k = 2$  case we also get  $\mathcal{F}_M$  satisfying (C) but not (B) for  $M = 22$  and  $\mathcal{F}_M$  satisfying (B) but not (C) for  $M = 35$  and  $39$ . This list is not changed if we look at a larger range of  $M$  (we have calculated up to  $M = 210$  in the weight 2 case and  $M = 100$  in the weight 1 case) or  $\Delta$  (in each case we have calculated about twice as many Fourier coefficients as those shown in the tables), so that it seems quite likely that it is complete.

For the weight 2 functions, the corollary of Theorem 9.41 tells us that  $C(\mathcal{F}_M; -1, 1)$  will always be negative with the normalization (9.52) and that we have

$$c_M = t \cdot \text{denom}\left(\frac{\varphi(M)}{24}\right), \quad C(\mathcal{F}_M; -1, 1) = -t \cdot \text{numer}\left(\frac{\varphi(M)}{24}\right) \tag{9.53}$$

for some positive rational number  $t$ . In all cases we have looked at,  $t$  turned out to be an integer, and in fact usually had the value 1. For instance, the second statement is true for all values of  $M$  tabulated in Appendix A, and also for  $M = 210$ , where Table 4 in §9.3 shows that  $c_M = 1$  and hence  $t = 1$ , but the table below giving all values up to  $M = 100$  shows that it fails for  $M = 65$  and  $M = 85$ .

$M$	6	10	14	15	21	22	26	33	34	35	38	39	46
$c_M$	12	6	4	3	2	12	2	6	3	1	4	1	12
$C(\mathcal{F}_M; -1, 1)$	1	1	1	1	1	5	1	5	2	1	3	1	11
$t$	1	1	1	1	1	1	1	1	1	1	1	1	1

51	55	57	58	62	65	69	74	77	82	85	86	87	91	93	94	95
3	3	2	6	4	2	6	2	2	3	6	4	3	1	2	12	1
4	5	3	7	5	4	11	3	5	5	16	7	7	3	5	23	3
1	1	1	1	1	2	1	1	1	1	2	1	1	1	1	1	1

If it is true that  $t$  is integral, then equation (9.53) shows that the above list of  $\mathcal{F}_M$  satisfying property (B) is complete, because we can then only have  $|C(\mathcal{F}_M; -1, 1)| = 1$  if  $\varphi(M)$  divides 24, which in turn implies that  $M$  is at most 90 and hence within the range of our calculations. We do not know whether the integrality is true in general, but in any case  $12^{M+1}t$  must be an integer, since correcting the initial choice  $\Phi_{2,M}^{\text{stand}} = 12^{-M-1}B^{M+1}A^{-1} - \mathcal{A}_{2,M}$  by weak Jacobi forms cannot make its denominator any worse.

Exactly similar comments apply in the weight 1 case. Here a residue calculation similar to the ones described in the next section (but with the residue  $\mathcal{R}$  as defined in (10.17) below replaced by its weight 1 analogue as defined in (10.59)) tells us that in the case of optimal growth eq. (9.43) still holds, so that we find that eq. (9.53) still holds for some positive rational number  $t$ . If this  $t$  turns out to be an integer, it again follows that property (B) can only hold if  $\varphi(M)|24$ , and the list of all integers  $M$  with  $\mu(M) = -1$  satisfying this latter condition coincides exactly with the list (9.46). Thus, although we cannot yet prove that that list is complete, we can at least understand where it comes from.

#### 4. Positivity of the Fourier coefficients

We now turn to properties (C) and (D). Here we can use the asymptotic formula given in Theorem 9.3 to understand the question of the signs and asymptotics of the Fourier coefficients. (We will discuss only the weight 2 case in detail, but a similar theorem and similar discussion could equally well be given for the case  $k = 1$ .) This theorem implies that (C) “almost holds” in the sense that for any fixed value of  $r$  the Fourier coefficients  $C(\Delta, r)$  eventually have a constant sign, so that there are at most finitely many exceptions to the statement. We therefore have to consider (D) and also the question of which values of  $r$  are most likely to give rise to the exceptional signs.

For the function  $\mathcal{Q}_M$  (which we remind the reader differs from  $\mathcal{F}_M$  by a negative proportionality factor, so that the expected behavior (D) would now say that  $C(\Delta, r)$  is always negative when  $r = r_{\min}$  is the smallest positive square root of  $-\Delta$ ), the value of the constant  $\kappa_r$  appearing in that theorem as given according to (9.39) and (9.43) by

$$\kappa(\mathcal{Q}_M, r) = -\frac{\varphi(M)}{24} \cdot 2 \sum_{\substack{0 < \ell < M \\ \ell^2 \equiv 1 \pmod{4M}}} \mu\left(\left(\frac{\ell+1}{2}, M\right)\right) \cos\left(\frac{\pi \ell r}{M}\right). \quad (9.54)$$

A similar formula applies in the weight 1 case with “cos” resplaced by “sin.” In the case when  $M$  is a prime, this immediately gives the positivity of  $\kappa(\mathcal{Q}_M, r_{\min})$ , and in the case when  $M$  has three prime factors, the tables in A.3 show that (D) usually fails. We therefore consider the weight 2 case, where  $\mu(M) = +1$ .

If  $M$  (which we always assume to be larger than 1) is a product of two primes, and in particular for all  $M < 330$  with  $\mu(M) = 1$  except 210, the sum on the right-hand side has only

two terms and reduces to simply  $\cos(\pi r/M) - \cos(\pi r^*/M)$ , and since  $0 < r_{\min} < r_{\min}^* < 2M$  and  $\cos(\pi x)$  is decreasing on the interval  $[0, \pi]$ , the value of  $-\kappa(\mathcal{Q}_M, r_{\min})$  is indeed always strictly negative. It then follows from Theorem 9.3 that almost all  $C(\mathcal{F}_M; \Delta, r_{\min})$  are positive, but there can be, and in general are, finitely many exceptions. However, for small  $M$  these are fairly rare, e.g., the following table lists all triples  $(M, r = r_{\min}, n)$  within the range of Appendix A for which  $c(\mathcal{F}_M; n, r)$  is negative.

$M$	6–22	33	34	35	38	39	46
$r$	—	8	15	2	17	10	21
$n$	—	1, 5, 7	5, 11	1, 5	5, 7, 11	5	5, 7, 11, 13

(The corresponding  $r$ -values are marked with a star in the appendix, and a search up to a much larger  $n$ -limit yields no further negative coefficients for these values of  $M$ .) Here again the asymptotic formula (9.38) helps us to understand this behavior: the negative values are much more likely to occur when  $\kappa_r$  is small. For instance, for  $M = 33$  the table

$r (= r_{\min})$	1	2	4	5	7	8	10	13	16	19
$\cos(\frac{\pi r}{66}) - \cos(\frac{\pi r^*}{66})$	1.58	1.31	1.71	0.94	1.71	0.49	1.58	1.31	0.94	0.49

shows that the residue classes  $r_{\min} \pmod{2M}$  most likely to give some negative Fourier coefficients are 8 and 19, for which the values of  $\kappa_r$  (which are the same; this duplication always occurs for odd  $M$ ) are minimal, and indeed the class  $r = 8$  did lead to three negative coefficients. The class  $r = 19$  did not contain any negative values of  $c(n, r)$ , but instead achieved its small initial average by the initial zeros forced by the optimal growth condition, which says that  $c(n, r) = 0$  for  $0 \leq n \leq (r^2 - 2)/4m$ .

For values of  $M$  for more than two prime factors on the other hand, it does not have to be true that  $\kappa(\mathcal{F}_M, r_{\min}) > 0$ . It is true for the first such index,  $M = 210$ , and even in a very strong form: not only is  $\kappa_r$  positive for each of the six possible values of  $r_{\min}$  (which are nothing other than the values  $\ell_0$  printed in boldface in Table 4), but of the seven “partners” of each  $r_{\min}$ , i.e., the seven other values of  $r \in [0, 210]$  with  $r^2 \equiv r_{\min}^2 \pmod{840}$ , the three that have the same sign of  $C(\Delta, r)$  as for  $C(\Delta, r_{\min})$  (i.e., which differ from  $r_{\min} \pmod{2p}$  for exactly two of the four prime divisors  $p$  of 210) all belong to the first half of the interval  $[0, 210]$ , while the four that have the opposite sign (i.e., which differ from  $r_{\min} \pmod{2p}$  for one or three of these  $p$ ) belong to the second half, so that all eight terms in the sum in (9.54) are positive. The value of  $\kappa(\mathcal{F}_M, r_{\min})$  is again positive for all  $r_{\min}$  for the next two values  $M = 330$  and  $M = 390$ , but for  $M = 462 = 2 \cdot 3 \cdot 7 \cdot 11$  and  $r (= r_{\min}) = 31$  we have  $\kappa(\mathcal{F}_M, r) < 0$ , so that in this case all but

finitely many coefficients  $C(\mathcal{F}_M; \Delta, r)$  will be negative rather than positive. What does hold for any  $M$  and any  $r \pmod{2M}$  is that the values of  $C(\mathcal{F}_M; \Delta, r)$  eventually all have the same sign, because the coefficient  $\kappa(\mathcal{F}_M, r)$  in (9.38) can never vanish, since the primitive  $(2M)^{\text{th}}$  roots of unity are linearly independent over  $\mathbb{Q}$ .

## 9.6 Higher weights

So far we have studied two families of meromorphic Jacobi forms  $\varphi_{1,m}(\tau, z)$  and  $\varphi_{1,m}(\tau, z)$  with particularly simple singularities, and the corresponding mock Jacobi forms. We now discuss in much less detail the analogous families for higher weights. The starting point is a meromorphic Jacobi form  $\varphi(\tau, z) = \varphi_{k,m}(\tau, z)$  of weight  $k$  and arbitrary index  $m \in \mathbb{N}$  having as its unique singularity (modulo the period lattice  $\mathbb{Z}\tau + \mathbb{Z}$ ) a pole of order  $k$  at  $z = 0$ , which we normalized by requiring the leading term of  $\varphi$  to be  $(k-1)!/(2\pi iz)^k$ . By subtracting a linear combination of functions  $\varphi_{k',m}$  with  $k' < k$  we can then fix the entire principal part of  $\varphi$  at  $z = 0$  to be

$$\varphi(\tau, z) = \frac{(k-1)!}{(2\pi iz)^k} \sum_{0 \leq 2j < k} \left( \frac{m E_2(\tau)}{12} \right)^j \frac{(2\pi iz)^{2j}}{j!} + O(1) \quad \text{as } z \rightarrow 0.$$

We can make a standard choice, generalizing the previously defined forms (9.2) and (9.1), by taking  $\varphi$  to be the unique polynomial in  $A^{-1}$  and  $B$  (resp.  $C$  times such a polynomial if  $k$  is odd) with modular coefficients of the appropriate weight having this principal part, the next three of these being

$$\varphi_{3,m}^{\text{stand}} = \frac{B^m C}{12^m A^2}, \quad \varphi_{4,m}^{\text{stand}} = \frac{6 B^{m+2}}{12^{m+2} A^2}, \quad \varphi_{5,m}^{\text{stand}} = \frac{(B^2 - m E_4 A^2 / 2) B^{m-1} C}{12^m A^3}.$$

Any other allowed choice of  $\varphi_{k,m}$  will differ from this standard one by a weak Jacobi form of weight  $k$  and index  $m$ , and any two choices will have the same polar part. To write it down, we introduce Euler's functions

$$\mathcal{E}_k(y) = \text{Av}_{\mathbb{Z}} \left[ \frac{(k-1)!}{(2\pi iz)^k} \right] = \left( -y \frac{d}{dy} \right)^{k-1} \text{Av}_{\mathbb{Z}} \left[ \frac{1}{2\pi iz} \right] = \left( -y \frac{d}{dy} \right)^{k-1} \left( \frac{1}{2} \frac{y+1}{y-1} \right) \in \mathbb{Q} \left[ \frac{1}{y-1} \right]$$

(which for  $k = 1$  and  $k = 2$  coincide with the functions  $\mathcal{R}_0$  and  $\mathcal{R}_0^{(2)}$  from §8) and their index  $m$  averages

$$\mathcal{A}_{k,m}(\tau, z) = \text{Av}_{\mathbb{Z}\tau + \mathbb{Z}}^{(m)} \left[ \frac{(k-1)!}{(2\pi iz)^k} \right] = \text{Av}^{(m)} [\mathcal{E}_k(y)] = \sum_{s \in \mathbb{Z}} q^{ms^2} y^{2ms} \mathcal{E}_k(q^s y)$$

generalizing (9.3). These functions all have wall-crossing, with Fourier expansions given by

$$\mathcal{E}_k(y) = \begin{cases} \frac{1}{2} \delta_{k,1} + \sum_{t>0} t^{k-1} y^{-t} & \text{if } |y| > 1 \\ -\frac{1}{2} \delta_{k,1} - \sum_{t<0} t^{k-1} y^{-t} & \text{if } |y| < 1 \end{cases} = \sum_{t \in \mathbb{Z}} \frac{\text{sgn}(t) - \text{sgn}(z_2)}{2} t^{k-1} y^{-t}$$

and

$$\mathcal{A}_{k,m}(\tau, z) = \sum_{s,t \in \mathbb{Z}} \frac{\operatorname{sgn}(t) - \operatorname{sgn}(s + z_2/\tau_2)}{2} t^{k-1} q^{ms^2-st} y^{2ms-t}. \quad (9.55)$$

(Note that in the latter expression the coefficient of  $y^r$  is a Laurent polynomial in  $q$  for every  $r$  and that the discriminant  $4nm - r^2$  is the negative of a perfect square for each monomial  $q^n y^r$  occurring.) Then the polar part of  $\varphi_{k,m}$  is given by

$$\varphi_{k,m}^P(\tau, z) = \sum_{0 \leq 2j < k} \frac{(k-1)!}{(k-1-2j)! j!} \left( \frac{m E_2(\tau)}{12} \right)^j \mathcal{A}_{k-2j,m}(\tau, z).$$

(To see this, observe that by (9.55) each of the functions  $\mathcal{A}_{k,m}$ , and hence any linear combination of them with coefficients depending only on  $\tau$ , is in the kernel of the operator  $\varphi \mapsto h_\ell$  defined by (8.2) for all  $\ell \in \mathbb{Z}$ .) Its finite part  $\Phi_{k,m} = \varphi_{k,m}^F = \varphi_{k,m} - \varphi_{k,m}^P$  is a weakly holomorphic elliptic form that depends on the particular choice of  $\varphi_{k,m}$ . From the results of §10 (specifically, Theorem 10.1) it follows that for  $k \geq 3$  the function  $\varphi_{k,m}$  can always be chosen so that  $\Phi_{k,m}$  is strongly holomorphic, this choice then obviously being unique up to the addition of a strongly holomorphic Jacobi form.

As an example, consider  $k = 3$  and  $m = 2$ . (This is the smallest possibility since index 1 cannot occur in odd weight.) The finite part of our standard choice is

$$\Phi_{3,2}^{\text{stand}} = \frac{B^2 C}{144 A^2} - \mathcal{A}_{3,2} - \frac{E_2}{3} \mathcal{A}_{3,1} = h_{3,2}^{\text{stand}} (\vartheta_{2,1} - \vartheta_{2,3})$$

with

$$h_{3,2}^{\text{stand}} = \frac{1}{144} q^{-1/8} (1 + 51q + 2121q^2 + 14422q^3 + 60339q^4 + 201996q^5 + 588989q^6 + \dots).$$

This is weakly holomorphic and has large Fourier coefficients, but by subtracting a multiple of the unique weak Jacobi form  $E_4 C$  of this weight and index we get a new choice

$$\Phi_{3,2}^{\text{opt}} = \Phi_{3,2}^{\text{stand}} - \frac{E_4}{144} C = h_{3,2}^{\text{opt}} (\vartheta_{2,1} - \vartheta_{2,3})$$

that is strongly holomorphic and has much smaller coefficients

$$h_{3,2}^{\text{opt}} = h_{3,2}^{\text{stand}} - \frac{E_4}{144 \eta^3} = -\frac{4}{3} q^{-1/8} (q + 4q^2 + 5q^3 + 11q^4 + 6q^5 + 21q^6 + \dots),$$

in accordance with the theorem just cited.

However, this choice, although optimal from the point of view of the order of growth of its Fourier coefficients, is not the most enlightening one. By subtracting a different multiple of  $E_4 C$  from  $\Phi_{3,2}^{\text{stand}}$ , we get a third function

$$\Phi_{3,2} = \Phi_{3,2}^{\text{stand}} + \frac{7E_4}{144} C = \Phi_{3,2}^{\text{opt}} + \frac{E_4}{18} C = h_{3,2} (\vartheta_{2,1} - \vartheta_{2,3})$$

with

$$h_{3,2} = h_{3,2}^{\text{stand}} + \frac{7E_4}{144\eta^3} = h_{3,2}^{\text{opt}} + \frac{E_4}{18\eta^3} = \frac{1}{18} q^{-1/8} (1 + 219q + 2793q^2 + 15262q^3 + \dots).$$

This function can be identified: it is given by

$$h_{3,2} = \frac{1}{3} \partial_{1/2}^{\text{RS}}(h_2),$$

where  $h_2(\tau) = q^{-1/8}(-1 + 45q + 231q^2 + \dots)$  is the mock theta function defined in (7.16) (the one occurring in the Mathieu moonshine story) and  $\partial_{1/2}^{\text{RS}}$  is the Ramanujan-Serre derivative as defined in (3.7). We can also write this at the level of mock Jacobi forms as

$$\Phi_{3,2} = \frac{1}{24} \mathcal{L}_{1,2}(\mathcal{F}_2),$$

where  $\mathcal{F}_2$  is the mock Jacobi form defined in (7.32) and  $\mathcal{L}_{1,2}$  the modified heat operator defined in (4.12).

An analogous statement holds in general, e.g., for weight 3 we have

$$\Phi_{3,m} = -\mathcal{L}_{1,m}(\Phi_{1,m}) \pmod{\tilde{J}_{3,m}}$$

for any index  $m$  and any choice of the functions  $\Phi_{3,m}$  and  $\Phi_{1,m}$  in their defining classes, and similarly for higher weight, the specific formulas for  $k \in \{3, 4\}$  and the standard choices being

$$\begin{aligned} \Phi_{3,m}^{\text{stand}} + \mathcal{L}_{1,m}(\Phi_{1,m}^{\text{stand}}) &= -\frac{m-1}{12^m} \left( 4(m-2)E_6A + (2m+3)E_4B \right) B^{m-3}C, \\ \Phi_{4,m}^{\text{stand}} + \mathcal{L}_{2,m}(\Phi_{2,m}^{\text{stand}}) &= -\frac{m+1}{72 \cdot 12^m} \left( 4mE_6A + (2m-3)E_4B \right) B^{m-1}. \end{aligned}$$

Thus in one sense, the higher-weight families are better than the weight 1 and weight 2 families (because they can be made strongly holomorphic by subtracting an appropriate weak Jacobi form), but in another, they are much less interesting (because they are always combinations of derivatives of lower-weight examples and ordinary Jacobi forms, so that they do not produce any essentially new functions).

## 10. Structure theorems for Jacobi and mock Jacobi forms

In the last section we introduced two special families of meromorphic Jacobi forms and the associated mock Jacobi forms and described various observations concerning them. In this section we try to give theoretical explanations for some of these. We begin by studying in §10.1 how the holomorphic Jacobi forms of weight  $k$  and index  $m$  lie in the space of weak Jacobi

forms of the same weight and index, or equivalently, what the relations are among the polar Fourier coefficients (i.e., the coefficients of  $q^n y^r$  with  $4nm < r^2$ ) of weak Jacobi forms. The results described here are relatively easy consequence of results in the literature but are, as far as we know, new. The remaining two subsections describe the application of these results to mock Jacobi forms in general and to our two special families in particular, providing fairly complete explanations of the experimentally found properties of the weight two family and partial explanations in the weight one case.

### 10.1 Relations among the polar coefficients of weak Jacobi forms

The definitions and main properties of holomorphic and weak Jacobi forms were reviewed in §4, following the more detailed treatment in [52]. Here we investigate the relationship of these two spaces in more detail.

By definition, a weak Jacobi form has coefficients  $c(n, r)$  which are non-zero only for  $n \geq 0$ , whereas a holomorphic one has coefficients that vanish unless  $4nm \geq r^2$ . By virtue of the periodicity property (4.4), this stronger condition need only be checked for  $r$  running over a set of representatives for  $\mathbb{Z}/2m\mathbb{Z}$ , say  $-m < r \leq m$ , and since condition (4.1) with  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$  also implies that  $c(n, -r) = (-1)^k c(n, r)$  and consequently  $C(\Delta, -r) = (-1)^k C(\Delta, r)$ , it in fact suffices to consider the coefficients with  $0 \leq r \leq m$  if  $k$  is even and those with  $0 < r < m$  if  $k$  is odd. In other words, for all positive integers  $k$  and  $m$  we have an exact sequence

$$0 \longrightarrow J_{k,m} \longrightarrow \tilde{J}_{k,m} \xrightarrow{P} \mathbb{C}^{\mathcal{N}_{\pm,m}} \quad ((-1)^k = \pm 1), \quad (10.1)$$

where

$$\begin{aligned} \mathcal{N}_{+,m} &= \{(r, n) \in \mathbb{Z}^2 \mid 0 \leq r \leq m, 0 \leq n < r^2/4m\}, \\ \mathcal{N}_{-,m} &= \{(r, n) \in \mathbb{Z}^2 \mid 0 < r < m, 0 \leq n < r^2/4m\}, \end{aligned}$$

and where the map  $P$  (“polar coefficients”) associates to  $\varphi \in \tilde{J}_{k,m}$  the collection of its Fourier coefficients  $c_\varphi(n, r)$  for  $(n, r) \in \mathcal{N}_{\pm,m}$ . In particular we have

$$\dim J_{k,m} \geq j(k, m) := \dim \tilde{J}_{k,m} - |\mathcal{N}_{\pm,m}| \quad ((-1)^k = \pm 1), \quad (10.2)$$

with equality if and only if the map  $P$  in (10.1) is surjective. (Cf. [52], p. 105.)

Using the description of  $\tilde{J}_{*,*}$  given in §4 (eqs. (4.25) and (4.35)) and the definition of  $\mathcal{N}_{\pm,m}$ , we can write this lower bound on the dimension explicitly as

$$j(k, m) = \begin{cases} \sum_{0 \leq j \leq m} (\dim M_{k+2j} - \lceil j^2/4m \rceil) & \text{if } k \text{ is even,} \\ \sum_{0 < j < m} (\dim M_{k+2j-1} - \lceil j^2/4m \rceil) & \text{if } k \text{ is odd,} \end{cases} \quad (10.3)$$

where  $\lceil x \rceil$  denotes the “ceiling” of a real number  $x$  (= smallest integer  $\geq x$ ). This formula is computable and elementary since  $\dim M_k = k/12 + O(1)$  with the  $O(1)$  term depending only



on  $k \pmod{12}$ , but is nevertheless not a good formula because the two terms  $\sum \dim M_{k+2j}$  and  $\sum [j^2/4m]$  are each equal to  $m^2/12 + O(m)$ , and  $j(k, m)$  is only of the order of  $m$ . In [52], pp. 122–125 it was shown that, somewhat surprisingly, the numbers  $j(k, m)$  can be given by a completely different-looking formula involving class numbers of imaginary quadratic fields. This formula has the form

$$j(k, m) = \frac{2k-3}{24}(m+1) + \frac{b}{4} - \frac{1}{4} \sum_{d|4m} h(-d) + O(1), \quad (10.4)$$

where  $m = ab^2$  with  $a$  square-free and where the  $O(1)$  term depends only on  $m$  and  $k$  modulo 12. In particular,  $j(k, m) = \frac{2k-3}{24}m + O(m^{1/2+\varepsilon})$  as  $m \rightarrow \infty$  with  $k$  fixed. The formula (10.4) has an interesting interpretation as the dimension of a certain space of classical modular forms of level  $m$  and weight  $2k-2$ , discussed in detail on pp. 125–131 of [52].

In [52], it was shown that the inequality (10.2) is an equality for  $k \geq m$  and it was conjectured that this holds for all  $k \geq 3$ . This was later proved in [104] as a consequence of a difficult trace formula calculation. In fact, a slightly stronger statement was proved there: It is clear that the exact sequence (10.1) remains valid if one replaces  $J_{k,m}$  by  $J_{k,m}^0$  and the map  $P : \tilde{J}_{k,m} \rightarrow \mathbb{C}^{\mathcal{N}_{\pm,m}}$  by the analogously defined map  $P^0 : \tilde{J}_{k,m} \rightarrow \mathbb{C}^{\mathcal{N}_{\pm,m}^0}$ , where  $\mathcal{N}_{\pm,m}^0$  is defined exactly like  $\mathcal{N}_{\pm,m}$  but with the strict inequality  $n < r^2/4m$  replaced by  $n \leq r^2/4m$ . Then the dimension formula proved in [104] says that  $\dim J_{k,m}^0 = \dim \tilde{J}_{k,m} - |\mathcal{N}_{\pm,m}^0|$  and hence that the map  $P^0$  is surjective, which is a stronger statement than the surjectivity of  $P$  since the set  $\mathcal{N}_{\pm,m}^0$  is larger than  $\mathcal{N}_{\pm,m}$ . In summary, the results of [52] and [104] together give the following complete description of the “polar coefficients” maps for both holomorphic and cuspidal Jacobi forms when  $k \geq 3$ :

**Theorem 10.1.** *For  $k > 2$  and all  $m \geq 1$  the map  $P^0 : \tilde{J}_{k,m} \rightarrow \mathbb{C}^{\mathcal{N}_{\pm,m}^0}$  is surjective and we have a commutative diagram of short exact sequences*

$$\begin{array}{ccccccc} 0 & \longrightarrow & J_{k,m}^0 & \longrightarrow & \tilde{J}_{k,m} & \xrightarrow{P^0} & \mathbb{C}^{\mathcal{N}_{\pm,m}^0} \longrightarrow 0 \\ & & \cap & & \parallel & & \downarrow \\ 0 & \longrightarrow & J_{k,m} & \longrightarrow & \tilde{J}_{k,m} & \xrightarrow{P} & \mathbb{C}^{\mathcal{N}_{\pm,m}} \longrightarrow 0 \end{array} \quad (10.5)$$

We now turn to the case  $k = 2$ , the one of main interest for us. Here formula (10.4) takes the explicit form

$$j(2, m) = \frac{1}{2} \left( \left\lfloor \frac{m+1}{3} \right\rfloor + \left\lfloor \frac{m+1}{3} \right\rfloor \right) - \frac{m-b-1}{4} - \frac{1}{4} \sum_{d|4m} h(-d), \quad (10.6)$$

but for this value of  $k$  it was observed in [52] (p. 131) that equality could not always hold in (10.2) (indeed, the right-hand side is often negative) and that, for reasons explained in detail



there, one should replace the number  $j(k, m)$  by the modified number

$$j^*(2, m) := j(2, m) + \delta(m), \quad \text{where} \quad \delta(m) := \sum_{\substack{d \leq \sqrt{m} \\ d|m, d^2 \nmid m}} 1. \quad (10.7)$$

We note that  $\delta(m)$  is a linear combination of three multiplicative functions of  $m$ , namely

$$\delta(m) = \frac{1}{2}\sigma_0(m) + \frac{1}{2}\delta_{a,1} - \sigma_0(b) \quad (10.8)$$

(with  $m = ab^2$ ,  $a$  square-free as before), or equivalently in terms of a generating Dirichlet series

$$\sum_{m=1}^{\infty} \frac{\delta(m)}{m^s} = \frac{1}{2}\zeta(s)^2 + \frac{1}{2}\zeta(2s) - \zeta(s)\zeta(2s). \quad (10.9)$$

We also note that  $\delta(m)$  depends only on the ‘‘shape’’ of  $m$  (= the collection of exponents  $\nu_i$  if  $m$  is written as  $\prod p_i^{\nu_i}$ ), the first three cases being

$$\delta(p^\nu) = 0, \quad (10.10)$$

$$\delta(p_1^{\nu_1} p_2^{\nu_2}) = n_1 n_2, \quad (10.11)$$

$$\delta(p_1^{\nu_1} p_2^{\nu_2} p_3^{\nu_3}) = 3n_1 n_2 n_3 + \varepsilon_1 n_2 n_3 + \varepsilon_2 n_1 n_3 + \varepsilon_3 n_1 n_2, \quad (10.12)$$

in which we have written each  $\nu_i$  as  $2n_i + \varepsilon_i - 1$  with  $n_i \geq 1$  and  $\varepsilon_i \in \{0, 1\}$ .

The conjecture made in [52] that  $\dim J_{k,m} = j^*(2, m)$  for all  $m \geq 1$  was proved in [104], but here we want to give a sharper statement describing the cokernel of the map  $P$  explicitly. To do this, we follow [52], pp. 132, where an elementary argument (containing a small error which will be corrected below) was given to show that  $\dim J_{2,m} \geq j^*(2, m)$  for square-free  $m$ . Actually, it is more convenient to work with cusp forms, since the corresponding statement here is

$$\dim J_{2,m}^0 = \dim \tilde{J}_{2,m} - |\mathcal{N}_{+,m}^0| + \delta^0(m) \quad (10.13)$$

with

$$\delta^0(m) = \sum_{d|m, d \leq \sqrt{m}} 1 = \sum_{\substack{m=m_1 \cdot m_2 \\ \text{up to order}}} 1, \quad (10.14)$$

and  $\delta^0(m)$  is a slightly simpler function than  $\delta(m)$ . (The third term in (10.8) is missing.) We want to explain formula (10.13) by finding an explicit map

$$\beta : \mathbb{C}^{\mathcal{N}_{+,m}^0} = \bigoplus_{\substack{0 \leq r \leq m \\ 0 \leq n \leq r^2/4m}} \mathbb{C} \longrightarrow \Delta^0(m) := \bigoplus_{\substack{m=m_1 \cdot m_2 \\ \text{up to order}}} \mathbb{C} \quad (10.15)$$

such that the sequence

$$0 \longrightarrow J_{2,m}^0 \longrightarrow \tilde{J}_{2,m} \xrightarrow{P^0} \mathbb{C}^{\mathcal{N}_{+,m}^0} \xrightarrow{\beta} \Delta^0(m) \longrightarrow 0. \quad (10.16)$$

is exact.

To give a natural definition of the map  $\beta$ , we work, not with  $\tilde{J}_{2,m}$  or even with  $J_{2,m}^!$ , but with the larger vector spaces  $\tilde{\mathcal{E}}_{+,m}$  and  $\mathcal{E}_{+,m}^!$  of even weak or weakly holomorphic elliptic forms of index  $m$ , as defined at the end of §4.4. We define a “residue map”  $\mathcal{R} : \mathcal{E}_{+,m}^! \rightarrow \mathbb{C}$  by

$$\varphi = \sum_{n \geq 0, r \in \mathbb{Z}} c(n, r) q^n y^r \quad \mapsto \quad \mathcal{R}[\varphi] = \operatorname{Res}_{\tau=\infty}(\varphi(\tau, 0) d\tau) = \sum_{r \in \mathbb{Z}} c(0, r). \quad (10.17)$$

Then  $\mathcal{R}[\varphi] = 0$  if  $\varphi \in J_{2,m}^!$  because in that case  $\varphi(\cdot, 0)$  belongs to  $M_2^!$  and the  $q^0$  term of any weakly holomorphic modular form  $f$  of weight two is zero by the residue theorem applied to the  $SL(2, \mathbb{Z})$ -invariant 1-form  $f(\tau)d\tau$ . (The map  $\mathcal{R}$  was already used in [52], but it was erroneously stated there that for  $\varphi \in \tilde{J}_{2,m}$  the function  $\varphi(\tau, 0)$  itself, rather than just its residue at infinity, vanishes.) Now for each decomposition  $m = m_1 \cdot m_2$  (up to order, i.e., without distinguishing the splittings  $m_1 \cdot m_2$  and  $m_2 \cdot m_1$ ) we define  $R_{m_1, m_2} : \mathcal{E}_{+,m}^! \rightarrow \mathbb{C}$  by setting  $R_{m_1, m_2}(\varphi) = \mathcal{R}[\varphi|W_{m_1}]$  if  $(m_1, m_2) = 1$ , where  $W_{m_1}$  is the involution defined in §4.4, while if  $(m_1, m_2) = t > 1$  then we define  $R_{m_1, m_2}(\varphi)$  as  $R_{m_1/t, m_2/t}(\varphi|u_t)$ , where  $u_t$  is the map from  $\mathcal{E}_{+,m}^!$  to  $\mathcal{E}_{+,m/t^2}^!$  defined in §4.4. (It was pointed out there that both  $W_{m_1}$  for  $(m_1, m/m_1) = 1$  and  $u_t$  for  $t^2|m$  make sense on all of  $\mathcal{E}_{+,m}^!$ , and not just on the subspace of holomorphic Jacobi forms, but that this is *not* true for weak elliptic or weak Jacobi forms, which is why we are now working with weakly holomorphic forms.) We put all of these maps together to a single map  $R : \mathcal{E}_{+,m}^! \rightarrow \Delta^0(m)$  and then define the desired map  $\beta$  by the commutative diagram

$$\begin{array}{ccc} \tilde{\mathcal{E}}_{+,m} & \xrightarrow{P^0} & \mathbb{C}^{\mathcal{N}_{+,m}^0} \\ \cap & & \cap \\ \mathcal{E}_{+,m}^! & \longrightarrow & \mathbb{C}^{\mathcal{N}_{+,m}^{!,0}} \\ R \searrow & & \swarrow \beta \\ & \Delta^0(m) & \end{array} \quad (10.18)$$

with  $\mathbb{C}^{\mathcal{N}_{+,m}^{!,0}}$  defined as in (10.23) below. (To see that such a map  $\beta$  exists, observe that the coefficients  $c(0, r)$  occurring in (10.17) all have discriminant  $\Delta = -r^2 \leq 0$ , that the operators  $W_{m_1}$  and  $u_t$  can be defined purely in terms of the coefficients  $C_\varphi(\Delta, r)$  and preserve the condition  $\Delta \leq 0$ , and finally that the coefficients  $C_\varphi(\Delta, r)$  for  $\varphi \in \tilde{\mathcal{E}}_{+,m}$  can all be expressed in terms of the coefficients  $c_\varphi(n, r)$  with  $(n, r) \in \mathcal{N}_{+,m}$  by virtue of the periodicity and evenness conditions defining  $\tilde{\mathcal{E}}_{+,m}$ .) The discussion above shows that the composition  $\beta \circ P^0$  in (10.16) vanishes, since  $\mathcal{R}$  vanishes on weakly holomorphic and hence in particular on weak Jacobi forms, and combining this statement with the dimension formula (10.13) proved in [104] we obtain the main result describing strong and weak Jacobi forms of weight 2 and arbitrary index:

**Theorem 10.2.** *The sequence (10.16) with  $\beta$  defined by (10.18) is exact for all integers  $m \geq 1$ .*

The two theorems above give a complete description of the space of Jacobi forms of all weights  $k \geq 2$ . We end this subsection by discussing the case of weight one. Here the description of Jacobi forms is very simple to state (though its proof is quite deep), since we have the following theorem of Skoruppa ([100], quoted as Theorem 5.7 of [52]):

**Theorem 10.3.** *We have  $J_{1,m} = \{0\}$  for all  $m \geq 1$ .*

Combining this with the discussion above, we find that in this case we have an exact sequence

$$0 \longrightarrow \tilde{J}_{1,m} \longrightarrow \bigoplus_{\substack{0 < r \leq m \\ 0 \leq n < r^2/4m}} \mathbb{C} \longrightarrow \mathbb{C}^{|j(1,m)|} \longrightarrow 0. \quad (10.19)$$

with

$$|j(1,m)| = -j(1,m) = \frac{m}{24} + O(m^{1/2+\varepsilon}).$$

We have not been able to find a natural description of the last space, or explicit description of the last map, in (10.19), analogous to the results given for  $k = 2$ .

## 10.2 Choosing optimal versions of weak elliptic forms

In this subsection, we will apply the theorem just explained to show how to make “optimal versions” of elliptic forms by counting actual Jacobi forms to get poles of small order.

Let  $\tilde{\mathcal{E}}_{+,m}$  be as in the previous subsection and  $\tilde{\mathcal{E}}_{-,m}$  be the similarly defined space with  $c(n, -r) = -c(n, r)$ , and let  $\mathcal{E}_{\pm,m}$  and  $\mathcal{E}_{\pm,m}^0$  be the corresponding spaces with the condition (4.7) replaced by (4.5) and (4.6), respectively. The discussion of §10.1 can be summarized as saying that we have a commutative diagram of short exact sequences,

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{E}_{\pm,m} & \longrightarrow & \tilde{\mathcal{E}}_{\pm,m} & \xrightarrow{P} & \mathbb{C}^{\mathcal{N}_{\pm,m}} \longrightarrow 0 \\ & & \cup & & \cup & & \parallel \\ 0 & \longrightarrow & J_{k,m} & \longrightarrow & \tilde{J}_{k,m} & \longrightarrow & \mathbb{C}^{\mathcal{N}_{\pm,m}} \longrightarrow 0 \end{array} \quad (10.20)$$

for  $k > 2$ ,  $(-1)^k = \pm 1$ , and similarly with  $J_{k,m}$  and  $\mathcal{E}_{\pm,m}$  replaced by  $J_{k,m}^0$  and  $\mathcal{E}_{\pm,m}^0$ ,  $\mathcal{N}_{\pm,m}$  by  $\mathcal{N}_{\pm,m}^0$  and  $P$  by  $P^0$ . For  $k = 2$ , we have instead:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{E}_{+,m}^0 & \longrightarrow & \tilde{\mathcal{E}}_{+,m} & \xrightarrow{P^0} & \mathbb{C}^{\mathcal{N}_{+,m}^0} \longrightarrow 0 \\ & & \cup & & \cup & & \parallel \\ 0 & \longrightarrow & J_{2,m}^0 & \longrightarrow & \tilde{J}_{2,m} & \longrightarrow & \mathbb{C}^{\mathcal{N}_{+,m}^0} \xrightarrow{\beta} \Delta^0(m) \longrightarrow 0 \end{array} \quad (10.21)$$

in the cuspidal case, where  $\Delta^0(m)$  was defined in (10.15) and  $\beta$  is defined by  $\beta \circ P^0 = R$ . As a consequence of these diagrams and of the vanishing of  $\delta(p^\nu)$  (cf. equation (10.10)), which implies that the diagram (10.20) also holds for  $k = 2$  when  $m$  is a prime power, we obtain:

**Theorem 10.4.** *Let  $k \geq 2$ , and  $\varphi \in \tilde{\mathcal{E}}_{\pm, m}$ , where  $(-1)^k = \pm 1$ . Then  $\varphi$  is the sum of an element of  $\tilde{\mathcal{J}}_{k, m}$  and an element of  $\mathcal{E}_{\pm, m}^0$  if and only if either*

- (i)  $k > 2$ , or
- (ii)  $k = 2$  and  $R(\varphi) (= \beta(P^0(\varphi))) = 0$ ;

*If  $k = 2$  and  $m$  is a prime power, then  $\varphi$  is the sum of an element of  $\tilde{\mathcal{J}}_{k, m}$  and an element of  $\mathcal{E}_{\pm, m}$ .*

Theorem 10.4 for  $k = 2$  makes a negative statement: if  $\varphi \in \tilde{\mathcal{E}}_{+, m}$  has  $\mathcal{R}[\varphi] \neq 0$ , and  $m$  is not a prime power, then  $\varphi$  cannot in general be decomposed as the sum of a weak Jacobi form of weight 2 and a holomorphic elliptic form. However, it turns out that we can come very close to this, writing  $\varphi$  as a sum of images under Hecke-like operators of forms that “have optimal growth” in the sense of §9.3. We now explain this in two steps. We first use the decomposition (4.44), and its variants for weakly holomorphic Jacobi or elliptic forms, to reduce to the case of primitive forms. We then show that any primitive even elliptic form can be corrected by the addition of a weakly holomorphic weight 2 Jacobi form to give a new elliptic form whose non-zero Fourier coefficients  $c(n, r)$  all have discriminant  $4mn - r^2 \geq -1$ .

Since the decomposition (4.44), as was already observed in §4.3 fails for weak Jacobi forms but holds for weakly holomorphic ones, we must first extend the above results to weakly holomorphic forms.

**Proposition 10.1.** *All statements in Theorem 10.4 remain true mutatis mutandis if all  $\tilde{\mathcal{E}}$ ,  $\tilde{\mathcal{J}}$  are replaced by  $\mathcal{E}^!$ ,  $J^!$ .*

The proposition is equivalent to saying that the analog of (10.21) still holds with weakly holomorphic rather than weak forms, i.e. we have a diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathcal{E}_{+, m}^0 & \longrightarrow & \mathcal{E}_{+, m}^! & \xrightarrow{P^0} & \mathbb{C}^{\mathcal{N}_{+, m}^{0, !}} \longrightarrow 0 \\
& & \cup & & \cup & & \parallel \\
0 & \longrightarrow & J_{2, m}^0 & \longrightarrow & J_{2, m}^! & \longrightarrow & \mathbb{C}^{\mathcal{N}_{+, m}^{0, !}} \xrightarrow{\beta} \Delta^0(m) \longrightarrow 0
\end{array} \tag{10.22}$$

where

$$\mathcal{N}_{+, m}^{0, !} = \{(r, n) \in \mathbb{Z}^2 \mid 0 \leq r \leq m, -\infty < n \leq r^2/4m\}. \tag{10.23}$$

We again prove this by comparing dimensions, even though the middle spaces in each row in the diagram (10.22) are infinite-dimensional. Since any elliptic form  $\varphi$  has a minimal value of  $n$  with  $c_\varphi(n, r) \neq 0$  for some  $r$ , it suffices to prove the exactness of the rows in (10.22) with  $\mathcal{E}_{+, m}^!$ ,  $J_{2, m}^!$ , and  $\mathbb{C}^{\mathcal{N}_{+, m}^{0, !}}$  replaced by the spaces  $\mathcal{E}_{+, m}^{\geq N}$ ,  $J_{2, m}^{\geq N}$ , and  $\mathbb{C}^{\mathcal{N}_{+, m}^{0, \geq N}}$ , of forms with non-zero coefficients only for  $n \geq -N$  for some  $N \geq 0$ . The only non-trivial part is the exactness at  $\mathbb{C}^{\mathcal{N}_{+, m}^{0, \geq -N}}$ , and since the composition of the maps  $P^0$  and  $\beta$  vanishes, we only need to check that the alternating sum of dimensions is zero. For this, we observe that the dimensions of  $J_{2, m}^{\geq -N}$

and  $\mathbb{C}^{\mathcal{N}_{+,m}^{0,\geq-N}}$  are both larger by  $(m+1)N$  than their values at  $N=0$ : for  $\mathbb{C}^{\mathcal{N}_{+,m}^{0,\geq-N}}$ , this is obvious since  $\mathcal{N}_{+,m}^{0,\geq-N}$  differs from  $\mathcal{N}_{+,m}^0$  by the addition of  $N$  rows of length  $m+1$ , and for  $J_{2,m}^{\geq-N}$  it follows from the computation

$$\begin{aligned} \dim J_{2,m}^{\geq-N} &= \dim \tilde{J}_{12N+2,m} \quad (\text{because } J_{2,m}^{\geq-N} = \Delta(\tau)^{-N} \tilde{J}_{12N+2,m}) \\ &= \sum_{j=0}^m \dim M_{k+12N+2j} = \dim \tilde{J}_{2,m} + (m+1)N. \end{aligned} \quad (10.24)$$

This completes the proof of the proposition.  $\square$

Now we can apply the decomposition (4.44) to the whole diagram (10.22) to find that everything can be reduced to primitive forms (and their images under the  $U_t$  operators). The key point is that although  $U_t$  and  $u_t$  change the discriminant  $\Delta = 4mn - r^2$  by multiplying or dividing it by  $t^2$ , respectively, they do not change its sign, and therefore all of the spaces in our diagram can be decomposed as direct sums of primitive parts of lower index and their images under  $U_t$ . This reduces everything to case of primitive forms.

For primitive forms our diagram becomes

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{E}_{+,m}^{0,\text{prim}} & \longrightarrow & \mathcal{E}_{+,m}^{!,\text{prim}} & \xrightarrow{P^0} & \mathbb{C}^{\mathcal{N}_{+,m}^{0!,\text{prim}}} & \longrightarrow & 0 \\ & & \cup & & \cup & & \parallel & & \\ 0 & \longrightarrow & J_{2,m}^{0,\text{prim}} & \longrightarrow & J_{2,m}^{!,\text{prim}} & \longrightarrow & \mathbb{C}^{\mathcal{N}_{+,m}^{0!,\text{prim}}} & \xrightarrow{\beta} & \Delta^{0,\text{prim}}(m) & \longrightarrow & 0 \end{array} \quad (10.25)$$

where

$$\Delta^{0,\text{prim}}(m) = \bigoplus_{\substack{m=m_1 \cdot m_2 \\ (m_1, m_2)=1 \\ \text{up to order}}} \mathbb{C}. \quad (10.26)$$

The map  $\beta$  in (10.25) is again defined by the requirement that  $\beta \circ P^0 = R$  on  $\mathcal{E}_{+,m}^{!,\text{prim}}$ , and lands in  $\Delta^{0,\text{prim}}(m)$  because  $\varphi|u_t = 0$  for  $\varphi$  primitive and  $t > 1$ . The space  $\Delta^{0,\text{prim}}(m)$  has dimension  $\delta^{0,\text{prim}}(m) = 2^{s-1}$  if  $m > 1$ , where  $s$  is the number of primes dividing  $m$ . (The failure of this formula for  $m=1$  corresponds to the second term in (10.8).)

We also have the non-cusp-form analogs of the diagrams (10.22) and (10.25), where all superscripts “0” have been dropped and where  $\mathcal{N}_{+,m}^!$  is defined as in (10.23) but with the condition  $4mn - r^2 \leq 0$  replaced by the strict inequality  $4mn - r^2 < 0$ , while  $\Delta^{\text{prim}}(m)$  is a space of dimension  $\delta^{\text{prim}}(m) = \delta^{0,\text{prim}}(m) - 1 = 2^{s-1} - 1$  (compare equation (10.8)). This space is the quotient of the space in (10.26) by the one-dimensional subspace of vectors with all components equal. This is essentially the idea of the proof of Theorem 9.4 given below.

Having completed these preparations, we now proceed to the proofs of our main results. We consider the case of square-free index first and then give a brief description of the modifications needed to extend them to primitive forms of arbitrary index.

For  $m > 1$  square-free, we will show that any  $\varphi \in \tilde{\mathcal{E}}_{+,m}$  can be corrected by a weak Jacobi form of weight 2 to get another elliptic form whose non-zero Fourier coefficients  $c(n, r)$  all have discriminant  $4mn - r^2 \geq -1$ , i.e., that we have a decomposition

$$\tilde{\mathcal{E}}_{+,m} = \tilde{J}_{2,m} + \mathcal{E}_{+,m}^{\text{OG}} \quad (m \text{ square-free}), \quad (10.27)$$

with  $\mathcal{E}_{+,m}^{\text{OG}}$  defined as in §9.3. Actually, this is still not quite the statement we want, since it is slightly wasteful: the codimension of  $\mathcal{E}_{+,m}^0$  in  $\mathcal{E}_{+,m}^{\text{OG}}$  is  $2^{s-1} + 1$ , where  $s$  as before is the number of prime factors of  $m$  (there are  $2^{s-1}$  pairs  $(r, n) \in \mathcal{N}_{+,m}^0$  with  $\Delta := 4nm - r^2$  equal to  $-1$ , and one pair  $(0,0)$  with  $\Delta = 0$ ), and there are only  $\dim \Delta^0(m) = 2^{s-1}$  constraints  $R_{m_1, m_2}(\varphi) = 0$  coming from diagram (10.21). Correspondingly, the intersection of the two spaces on the right-hand side of (10.27) contains  $J_{2,m}$  ( $= J_{2,m}^0$  in this case) as a subspace of codimension 1. To get a sharp statement, we should therefore replace  $\mathcal{E}_{+,m}^{\text{OG}}$  by a suitable codimension one subspace. There are three natural ways to do this, each of which leads to a sharpening of (10.27), corresponding exactly (in the square-free case) to the various choices of  $\Phi_{2,m}^0$  introduced in the discussion preceding Theorem 9.5. What's more, all statements remain true verbatim for arbitrary  $m > 1$  if we restrict to primitive forms. We state all three variants in a single theorem.

**Theorem 10.5.** *Suppose that  $m > 1$  is square-free and let  $\mathcal{E}_m^{\text{OG},\star}$  ( $\star = \text{I, II, III}$ ) denote the codimension one subspace of  $\mathcal{E}_{+,m}^{\text{OG}}$  defined by one of the three conditions*

- (I)  $\sum_{0 < r < m, r^2 \equiv 1 \pmod{4m}} C_\varphi(-1; r) = 0,$
- (II)  $c_\varphi(0, 1) = 0,$
- (III)  $c_\varphi(0, 0) = 0,$

respectively. Then we have an exact sequence

$$0 \longrightarrow J_{2,m} \xrightarrow{\text{diag}} \tilde{J}_{2,m} \oplus \mathcal{E}_m^{\text{OG},\star} \xrightarrow{-} \tilde{\mathcal{E}}_{+,m} \longrightarrow 0, \quad (10.28)$$

i.e., any even weak elliptic form of index  $m$  can be decomposed as the sum of a weak Jacobi form of weight 2 and index  $m$  and an element of  $\mathcal{E}_m^{\text{OG},\star}$ , and this decomposition is unique up to a holomorphic Jacobi form (which can be added to one term and subtracted from the other).

For  $m > 1$  the same statement remains true if we replace  $\tilde{\mathcal{E}}_{+,m}$  by its subspace  $\tilde{\mathcal{E}}_{+,m}^{\text{res-prim}}$  of forms  $\varphi$  satisfying  $R(\varphi|u_t) = 0$  for all  $t > 1$  and  $\mathcal{E}_m^{\text{OG},\star}$  by its intersection with  $\tilde{\mathcal{E}}_{+,m}^{\text{res-prim}}$ .

*Proof.* By the exactness of the top line of diagram (10.21), we can find functions  $\psi_{m;(r,n)} \in \tilde{\mathcal{E}}_{+,m}$ , unique up to the addition of cusp forms, mapping to the canonical basis  $\delta_{(r,n)}$  of  $\mathbb{C}^{\mathcal{N}_{+,m}^0}$ . Clearly, the space  $\mathcal{E}_{+,m}^{\text{OG,III}}/\mathcal{E}_{+,m}^0$  is spanned by  $\psi_{m;(r,n)}$  with  $0 < r < m$  and  $4mn - r^2 = -1$ , and has dimension  $2^{s-1}$ . (There are exactly two square roots of 1 modulo  $2P$  for each exact prime power divisor  $P$  of  $m$  and these can be combined arbitrarily by the Chinese remainder theorem. This gives  $2^s$  residue classes  $r \pmod{2m}$  with  $r^2 \equiv 1 \pmod{4m}$ , and if we choose the representatives of smallest absolute values then half of them are positive.) Moreover, the

basis elements can be chosen to be a single orbit of the group  $\{W_{m_1}\}_{m_1|m}$  introduced in §4.4, whose cardinality is the same number  $2^{s-1}$ . Namely, if  $A_m$  is any choice of  $\psi_{m;(1,0)}$ , i.e. any function in  $\tilde{\mathcal{E}}_{+,m}$  whose unique non-zero coefficient in  $\mathcal{N}_{+,m}^0$  is  $c(0,1) = 1$ , then the other functions  $\psi_{m;(r,n)}$  with  $(r,n) \in \mathcal{N}_{+,m}^0$  and  $4mn - r^2 = -1$ , can be chosen to be the images of  $A_m$  under the Atkin-Lehner involutions  $W_{m_1}$ ,  $m = m_1 m_2$ . More explicitly, with this choice  $\psi_{m;(r,n)}$  equals  $A_m|W_{m_1}$  with  $m_1 = \gcd(\frac{r+1}{2}, m)$ ,  $m_2 = \gcd(\frac{r-1}{2}, m)$ , and conversely  $A_m|W_{m_1}$  for any decomposition  $m = m_1 m_2$  equals  $\psi_{m;(|r_0|, (r_0^2-1)/4m)}$  where  $r_0$  ( $= 1^*$  in the notation of §4.4) is the unique  $r_0$  satisfying

$$r_0 \equiv -1 \pmod{2m_1}, \quad r_0 \equiv 1 \pmod{2m_2}, \quad |r_0| < m. \quad (10.29)$$

On the other hand, for any  $\varphi \in \mathcal{E}_{+,m}^{\text{OG,III}}$ , we have

$$R_{1,m}(\varphi) = \mathcal{R}[\varphi] = 2c_\varphi(0,1) \quad (10.30)$$

(because  $c_\varphi(0,-1) = c_\varphi(0,1)$  and  $c_\varphi(0,r) = 0$  for all  $r \neq \pm 1$  by the defining property of  $\mathcal{E}_{+,m}^{\text{OG,III}}$ ), and more generally,

$$R_{m_1,m_2}(\varphi) = 2c_\varphi(r_0, (r_0^2-1)/4m) \quad (10.31)$$

for each decomposition  $m = m_1 m_2$ , and with  $r_0$  as in (10.29). It follows that, for any two decompositions  $m = m_1 m_2$  and  $m = m'_1 m'_2$ , we have

$$R_{m_1,m_2}(A_m|W_{m'_1}) = \begin{cases} 2 & \text{if } m'_1 = m_1 \text{ or } m'_1 = m_2, \\ 0 & \text{otherwise.} \end{cases} \quad (10.32)$$

This shows that the map

$$\mathcal{E}_{+,m}^{\text{OG,III}}/\mathcal{E}_{+,m}^0 \xrightarrow{R} \bigoplus_{\substack{m=m_1 \cdot m_2 \\ \text{up to order}}} \mathbb{C} \quad (10.33)$$

is an isomorphism, and in combination with Theorem 10.4, completes the proof of the theorem in the case of square-free  $m$  and  $\star = \text{III}$ : any  $\varphi \in \tilde{\mathcal{E}}_{+,m}$  can be decomposed, uniquely up to Jacobi cusp forms, as the sum of  $\frac{1}{2} \sum_{m=m_1 m_2} R_{m_1,m_2}(\varphi) A_m|W_{m_1} \in \mathcal{E}_{+,m}^{\text{OG,III}}$  and an element of  $\tilde{\mathcal{J}}_{2,m}$ .

The proofs for  $\star = \text{I}$  and  $\star = \text{II}$  are almost identical, changing the basis  $\{\psi_{m;(r,n)} = A_m|W_{m_1}\}$  of  $\mathcal{E}_{+,m}^{\text{OG,III}}/\mathcal{E}_{+,m}^0$  in both cases in such a way as to satisfy the new defining condition without changing the residues, i.e., in the case  $\star = \text{II}$  by replacing the first basis element  $\psi_{m;(1,0)}$  by  $2\psi_{m;(0,0)}$ , and in the case  $\star = \text{I}$  by adding  $2\psi_{m;(0,0)} - 2^{-s+1} \sum_{r'} \psi_{m;(r',n)}$  to each basis element.

The proof of the statement for forms of arbitrary index  $m$  is very similar, since there are still  $2^{s-1}$  polar coefficients of a form in  $\mathcal{E}_{+,m}^{\text{OG}}$ , in one-to-one correspondence with the decompositions of  $m$  into two coprime factors. The details are left to the reader.  $\square$

Theorem 10.5 says that the first line in the diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & J_{2,m}^{\text{prim}} & \xrightarrow{\text{diag}} & \tilde{J}_{2,m}^{\text{prim}} \oplus \mathcal{E}_m^{\text{OG,prim},\star} & \xrightarrow{-} & \tilde{\mathcal{E}}_{+,m}^{\text{prim}} \longrightarrow 0 \\
& & \cap & & \cap & & \parallel \\
0 & \longrightarrow & J_{2,m}^{\text{OG,prim}} & \xrightarrow{\text{diag}} & \tilde{J}_{2,m}^{\text{prim}} \oplus \mathcal{E}_m^{\text{OG,prim}} & \xrightarrow{-} & \tilde{\mathcal{E}}_{+,m}^{\text{prim}} \longrightarrow 0
\end{array} \tag{10.34}$$

is exact. The exactness of the second line is a trivial consequence, and the fact that  $\mathcal{E}_{+,m}^{\text{OG,prim},\star}$  has codimension exactly one in  $\mathcal{E}_{+,m}^{\text{OG,prim}}$  implies that  $J_{2,m}^{\text{prim}}$  has codimension one in  $J_{2,m}^{\text{OG,prim}}$ . Moreover, the proof of the theorem also shows that  $J_{2,m}^{\text{OG,}\star} = J_{2,m}$ . (If  $\varphi \in J_{2,m}^{\text{OG}}$ , then the vanishing of the residues  $R_{m_1 m_2}(\varphi)$  for all decompositions with  $(m_1, m_2) = 1$  implies that the polar coefficients  $C_\varphi(-1, r)$  are all equal to one another and to  $-\frac{1}{2}c_\varphi(0, 0)$ , so that imposing any of the conditions (I), (II), (III) forces  $\varphi$  to be holomorphic. Conversely, if  $\varphi \in J_{2,m}$ , then the conditions OG, (I), (II) are trivial, and (III) holds because  $\mathcal{R}[\varphi] = 0$ .) It follows that the codimension of  $J_{2,m}$  in  $J_{2,m}^{\text{OG}}$  is also one.

### 10.3 The residues of mock Jacobi forms

We can apply the general theorems of the previous subsection to mock Jacobi forms. In particular, if we have any weak mock Jacobi form of weight  $k \geq 3$  and index  $m$ , it will be in  $\tilde{\mathcal{E}}_{\pm, m}$ , and can be corrected by a weak Jacobi form to get a holomorphic mock Jacobi form. In the case  $k = 2$ , there is an obstruction  $R(\varphi)$  to this being true, but, by the way we defined it, this obstruction can only depend on the shadow of  $\varphi$ . In this subsection, we show how to compute  $R(\varphi)$  explicitly in terms of the shadow. Since  $R(\varphi)$  is the vector with components  $\mathcal{R}[\varphi|u_{(m_1, m_2)}|W_{m_1/(m_1, m_2)}]$ , it suffices to compute  $R[\varphi]$  for a weak mock Jacobi form of weight 2 in terms of the shadow of  $\varphi$ .

Let  $\varphi(\tau, z)$  be a (strong or weak) mock Jacobi form of weight 2 and index  $m$ . Then, by definition,  $\varphi$  has a theta expansion (4.10) where each  $h_\ell$  is a (in general weakly holomorphic) mock modular form of weight  $3/2$  and has a completion  $\hat{h}_\ell = h_\ell + g_\ell^*$  for some modular form  $g_\ell$  of weight  $1/2$  such that the completion  $\hat{\varphi} = \sum_\ell \hat{h}_\ell \vartheta_{m, \ell} = \varphi + \sum_\ell g_\ell^* \vartheta_{m, \ell}$  transforms like a Jacobi form of weight 2 and index  $m$ .

As usual, we denote by  $\vartheta_{m, \ell}^0(\tau)$  the Thetanullwert  $\vartheta_{m, \ell}(\tau, 0)$ .

**Theorem 10.6.** *Let  $\varphi$  be as above. Then*

$$\mathcal{R}[\varphi] = \frac{\sqrt{\pi}}{6} \sum_{\ell \pmod{2m}} (\vartheta_{m, \ell}^0, g_\ell), \tag{10.35}$$

where  $(\cdot, \cdot)$  denotes the Petersson scalar product in weight  $1/2$ .



*Proof.* With the normalization of the Petersson scalar product as given in (3.14), we have

$$\sum_{\ell \pmod{2m}} (\vartheta_{m,\ell}^0, g_\ell) = \frac{3}{\pi} \int_{\mathcal{F}} F(\tau) d\mu(\tau), \quad (10.36)$$

where

$$F(\tau) = \tau_2^{1/2} \sum_{\ell \pmod{2m}} \overline{g_\ell(\tau)} \vartheta_{m,\ell}^0(\tau), \quad (10.37)$$

and  $\mathcal{F}$  denotes a fundamental domain for  $\Gamma \equiv SL_2(\mathbb{Z})$ . This makes sense since the function  $F(\tau)$  is  $\Gamma$ -invariant (the individual terms in (10.37) are not, which is why we had to be a little careful in the normalization of the scalar product) and is convergent because the weight of the modular forms  $g_\ell$  and  $\vartheta_{m,\ell}^0$  is less than 1. From eq. (7.1) it follows that the  $\bar{\tau}$ -derivative of the completion

$$\widehat{\varphi}(\tau, 0) = \varphi(\tau, 0) + \sum_{\ell \pmod{2m}} g_\ell^*(\tau) \vartheta_{m,\ell}^0(\tau), \quad (10.38)$$

of  $\varphi(\tau, 0)$  satisfies

$$\frac{\partial}{\partial \bar{\tau}} \widehat{\varphi}(\tau, 0) = \frac{1}{4i\sqrt{\pi}} \frac{F(\tau)}{\tau_2^2}. \quad (10.39)$$

The fact that  $\widehat{\varphi}(\tau, 0)$  transforms like a modular form of weight 2 on  $\Gamma$  implies that the differential one-form  $\omega = \widehat{\varphi}(\tau, 0) d\tau$  is  $\Gamma$  invariant, and equation (10.39) implies

$$d\omega = d(\widehat{\varphi}(\tau, 0) d\tau) = -\frac{\partial \widehat{\varphi}(\tau, 0)}{\partial \bar{\tau}} d\tau d\bar{\tau} = \frac{1}{2\sqrt{\pi}} F(\tau) d\mu(\tau). \quad (10.40)$$

Therefore, by Stokes's theorem, we have

$$\frac{\sqrt{\pi}}{6} \sum_{\ell \pmod{2m}} (\vartheta_{m,\ell}^0, g_\ell) = \int_{\mathcal{F}} d\omega = \int_{\partial \mathcal{F}} \omega = \lim_{T \rightarrow \infty} \int_{\partial \mathcal{F}(T)} \omega \quad (10.41)$$

where  $\mathcal{F}(T)$  is the truncated fundamental domain

$$\{\tau \in \mathcal{H} \mid |\tau| > 1, |\tau_1| < \frac{1}{2}, \tau_2 < T\}. \quad (10.42)$$

On the other hand, we have

$$\int_{\partial \mathcal{F}(T)} \omega = \int_{-\frac{1}{2}+iT}^{\frac{1}{2}+iT} \omega = \int_{-\frac{1}{2}+iT}^{\frac{1}{2}+iT} \varphi(\tau, 0) d\tau + O\left(\frac{1}{\sqrt{T}}\right). \quad (10.43)$$

The first equality holds because  $\omega$  is  $\Gamma$ -invariant, and all the edges of  $\mathcal{F}(T)$  except the top edge  $\tau_2 = T$  come in pairs which are  $\Gamma$ -equivalent, but of opposite orientation. The second equality follows from (10.38), and because  $\vartheta_{m,\ell}^0$  is  $O(1)$  and  $g_\ell^*(\tau)$  is  $O(T^{-1/2})$  by (7.2) with  $k = 3/2$ . The theorem then follows.  $\square$

The theorem immediately implies a more general formula for the residue  $R_{m_1, m_2}(\varphi) = \mathcal{R}[\varphi|W_{m_1}]$  when  $m = m_1 m_2$  with  $(m_1, m_2) = 1$  in terms of Petersson scalar products, namely,

$$R_{m_1, m_2}(\varphi) = \frac{\sqrt{\pi}}{6} \sum_{\ell \pmod{2m}} (\vartheta_{m, \ell}^0, g_{\ell^*}), \quad (10.44)$$

where  $\ell \mapsto \ell^*$  is the involution on  $(\mathbb{Z}/2m\mathbb{Z})^*$  appearing in the definition of  $W_{m_1}$ .

#### 10.4 Remaining proofs of theorems from §9

In this subsection, we apply the results proved so far to the special family studied in §9, namely, the meromorphic Jacobi forms  $\varphi_{2, m}$  having a pole  $1/(2\pi iz)^2$  at the origin and their associated mock Jacobi forms  $\Phi_{2, m}$ . Specifically, we shall prove Theorems 9.1, 9.2, 9.4, 9.5, and 9.7.

We begin with a preliminary remark. Two of the results we need to prove are Theorem 9.1 and the statement that the functions  $\Phi_{2, m}^0$  defined implicitly by (9.37) are primitive modulo weak Jacobi forms. Both of these have the form that a certain elliptic form is in fact a (weak) Jacobi form, i.e., that its shadow vanishes. Since the forms in question are defined using the Hecke-like operators defined in §4.4, we first have to explain how the action of these operators extends to non-holomorphic elliptic forms, since then we can apply them to the completions of our mock Jacobi forms and verify the necessary identities between shadows. The definitions of the Hecke-like operators were given in terms of the Fourier coefficients  $c(n, r)$  or  $C(\Delta, \ell)$  defined by (4.3) and (4.4). In the case of the completions of mock modular forms  $\Phi$ , when we apply equation (4.36) or (4.37) for the action of  $U_t$  or  $V_t$  with the  $\varphi$  there replaced by  $\Phi^C := \widehat{\Phi} - \Phi$  (the “correction term” discussed at the end of §8.3), we must interpret  $c(n, r)$  as the coefficient of  $\beta(|n|\tau_2) q^n y^r$  rather than simply  $q^n y^r$  as in the holomorphic case, where  $\beta(t) = \sqrt{\pi/4m} \operatorname{erfc}(2\sqrt{\pi t})$  in the weight 2 case, and similarly for equation (4.43) and  $C(\Delta, r \pmod{2m})$ . In view of (9.5) and (7.2), these coefficients in the case of  $\Phi_{2, m}$  are given by

$$C(\Phi_{2, m}^C; \Delta, \ell \pmod{2m}) = \sum_{\substack{\lambda^2 = -\Delta \\ \lambda \equiv \ell \pmod{2m}}} |\lambda|. \quad (10.45)$$

**Proof of Theorem 9.1:** We have to show that for  $m = P_1 \cdots P_r$  ( $P_i = p_i^{\nu_i}$ ,  $p_i$  distinct primes) we have

$$\Phi_{2, m} \Big| \prod_{i=1}^r (1 + \varepsilon_i W_{P_i}) \equiv \left( \Phi_{2, M} \Big| \prod_{p|M} (1 - W_p) \right) \Big| \mathcal{V}_{2, m/M}^{(M)}, \quad (10.46)$$

where  $M$  is the product of the primes  $p_i$  for which  $\varepsilon_i = -1$  and “ $\equiv$ ” means that the two sides of the equation differ by a weak Jacobi form. Because of the multiplicativity of the Hecke-like

operators  $U_d$ ,  $V_t$  and  $W_{m_1}$ , this formula can be rewritten as

$$\Phi_{2,m} \left| \prod_{i=1}^r (1 + \varepsilon_i W_{P_i}) \equiv \Phi_{2,M} \left| \prod_{\varepsilon_i=-1} (1 - W_{P_i}) \mathcal{V}_{P_i/P_i}^{(p_i)} \right| \prod_{\varepsilon_i=+1} \mathcal{V}_{P_i}^{(1)}, \quad (10.47)$$

so by induction on  $r$ , it suffices to show that if  $P = p^\nu || m$  then

$$\Phi_{2,m} \left| (1 + W_P) \equiv \Phi_{2,m/P} \left| \mathcal{V}_{2,P}^{(1)} \right., \quad \Phi_{2,m} \left| (1 - W_P) \equiv \Phi_{2,m/P} \left| (1 - W_P) \mathcal{V}_{2,P/p}^{(p)} \right. \right. \quad (10.48)$$

For this, it is enough to show that in each the difference of the left-hand side and the right-hand side has shadow zero.

For the first equation in (10.48), eq. (10.45) implies that the ‘‘correction term’’ for the left-hand side is given by

$$\begin{aligned} C(\Phi_{2,m}^C \left| (1 + W_P); \Delta, \ell \pmod{2m} \right.) &= C(\Phi_{2,m}^C; \Delta, \ell \pmod{2m}) + C(\Phi_{2,m}^C; \Delta, \ell^* \pmod{2m}) \\ &= \begin{cases} D(\delta_{D,\ell}^{(2m)} + \delta_{D,-\ell}^{(2m)} + \delta_{D,\ell^*}^{(2m)} + \delta_{D,-\ell^*}^{(2m)}) & \text{if } \Delta = -D^2, \text{ for some } D \in \mathbb{N} \\ 0 & \text{if } -\Delta \text{ is not a perfect square} \end{cases} \end{aligned} \quad (10.49)$$

where  $\ell^* \pmod{2m}$  is defined by  $\ell^* \equiv -\ell \pmod{2P}$ ,  $\ell^* \equiv +\ell \pmod{2m/P}$ , and the notation  $\delta_{a,b}^{(n)}$  means  $\delta_{a,b \pmod{n}}$ . The correction term for the right-hand side, on the other hand, is given (since  $\mathcal{V}_{2,P}^{(1)} = V_{2,P} - pV_{2,P/p^2}U_p$ , where the second term is omitted if  $p^2 \nmid P$ ) by

$$\begin{aligned} &\sum_{d \mid (\frac{\Delta+\ell^2}{4m}, \ell, P)} d C\left(\Phi_{2,\frac{m}{P}}^C; \frac{\Delta}{d^2}, \frac{\ell}{d} \pmod{2\frac{m}{P}}\right) - p \sum_{d \mid (\frac{\Delta+\ell^2}{4m}, \ell, \frac{P}{p^2})} d C\left(\Phi_{2,\frac{m}{P}}^C; \frac{\Delta}{p^2 d^2}, \frac{\ell}{pd} \pmod{2\frac{m}{P}}\right) \\ &= \sum_{d \mid (\frac{\Delta+\ell^2}{4m}, \ell, P)} \sum_{\lambda \equiv \ell \pmod{\frac{2m}{P}d}}^{\lambda^2 = -\Delta} |\lambda| - \sum_{d \mid (\frac{\Delta+\ell^2}{4m}, \ell, \frac{P}{p^2})} \sum_{\lambda \equiv \ell \pmod{\frac{2m}{P}d}}^{\lambda^2 = -\Delta} |\lambda|, \\ &= \sum_{\substack{d \in \{P, P/p\} \\ d \mid (\frac{\Delta+\ell^2}{4m}, \ell, P)}} \sum_{\lambda \equiv \ell \pmod{\frac{2m}{P}d}}^{\lambda^2 = -\Delta} |\lambda|, \end{aligned} \quad (10.50)$$

where the last line holds because the terms in the first sum in the second line with  $d|Pp^{-2}$  cancel with the terms in the second sum. The terms with  $d = P$  in the third line give the first two terms  $\sqrt{-D} = \pm\ell^*$  of the right hand side of (10.49), while the terms with  $d = P$  in the third line give the second two terms  $\sqrt{-D} = \pm\ell^*$ .

The correction term for the left-hand side of the second equation in (10.48) is given by the the right-hand side of (10.49) with the signs of the last two terms changed. This is now equal to zero unless  $(\ell, m) = 1$ , in which case it is equal to  $\sqrt{-\Delta}$  (resp.  $-\sqrt{-\Delta}$ ) if  $D \equiv \pm\ell \pmod{2m}$  (resp.  $D \equiv \pm\ell^* \pmod{2m}$ ). The correction term for the right-hand side, on the other hand, is

given by

$$\begin{aligned}
& \sum_{d|(\ell, \frac{P}{p}, \frac{\Delta+\ell^2}{4m})} d \left( C\left(\Phi_{2,mp/P}^C; \frac{\Delta}{d^2}, \frac{\ell}{d} \pmod{\frac{2mp}{P}}\right) - C\left(\Phi_{2,mp/P}^C; \frac{\Delta}{d^2}, \frac{\ell^*}{d} \pmod{\frac{2mp}{P}}\right) \right) \\
&= \sum_{d|(\ell, \frac{P}{p}, \frac{\Delta+\ell^2}{4m})} \left( \sum_{\substack{\lambda^2 = -\Delta \\ \lambda \equiv \frac{\ell}{d} \pmod{\frac{2mp}{P}}}} - \sum_{\substack{\lambda^2 = -\Delta \\ \lambda \equiv \frac{\ell^*}{d} \pmod{\frac{2mp}{P}}}} \right) |\lambda|, \tag{10.51}
\end{aligned}$$

where  $\ell^* \pmod{2mp/P}$  is now defined by  $\ell^* \equiv -\ell \pmod{2p}$ ,  $\ell^* \equiv +\ell \pmod{2m/P}$ . If  $p|\ell$ , then  $\ell^* = \ell$  and the expression vanishes, so that only  $d = 1$  contributes in the first summation. The condition  $4m|(\Delta + \ell^2)$  combined with  $\lambda \equiv \ell \pmod{2mp/P}$  implies  $\lambda \equiv \ell \pmod{2m}$  (and similarly for  $\ell^*$  since  $\ell^2 \equiv \ell^{*2} \pmod{2mp/P}$ ), thus proving the second equation in (10.48).  $\square$

**Proof of Theorem 9.2:** By Möbius inversion, the mock Jacobi forms  $\Phi_{2,m}$  associated to any choice of minimal meromorphic Jacobi forms  $\varphi_{2,m}$  can be represented in the form (9.37), where  $\Phi_{2,m}^0$  is defined by

$$\Phi_{2,m}^0 = \sum_{d^2|m} \mu(d) \Phi_{2,m/d^2} | U_d. \tag{10.52}$$

We have to show that  $\Phi_{2,m}^0$  can be chosen to be primitive and to have optimal growth.

To prove the second of these statements, it is enough to show that  $\Phi_{2,m}^0 | u_p$  has no shadow for all primes  $p$  with  $p^2|m$ . Then, by Theorem 10.5, the weak elliptic form  $\Phi_{2,m}^0$  can be chosen (by adding a weak Jacobi form) to have optimal growth. Writing the square divisors of  $m$  as  $d^2 p^{2i}$  with  $p \nmid d$  and noting that  $\mu(dp^i) = 0$  for  $i > 1$  and that  $U_{dp} = U_d U_p$ ,  $U_d u_p = u_p U_d$ , and  $U_p u_p = p$ , we find from (10.52) that

$$\begin{aligned}
\Phi_{2,m}^0 | u_p &= \sum_{d^2|m, p \nmid d} \mu(d) [\Phi_{2,m/d^2} | U_d - \Phi_{2,m/d^2 p^2} | U_d | U_p] | u_p \\
&= \sum_{d^2|m, p \nmid d} \mu(d) [\Phi_{2,m/d^2} | u_p - p \Phi_{2,m/d^2 p^2}] | U_d,
\end{aligned}$$

so it suffices to show that  $\Phi_{2,m} | u_p$  and  $p \Phi_{2,m/p^2}$  have the same shadow for any  $m$  with  $p^2|m$ . But the shadow of  $\Phi_{2,m} = \varphi_{2,m}^F$  is determined by  $\varphi_{2,m}^C = \widehat{\Phi}_{2,m} - \Phi_{2,m}$  (cf. comments at the end of §8.3), and by (10.45), we have:

$$C(\varphi_{2,m}^C; \Delta p^2, rp \pmod{2m}) = \begin{cases} p|\lambda| & \text{if } \Delta = -\lambda^2, \quad \lambda \equiv r \pmod{2m/p}, \\ 0 & \text{otherwise.} \end{cases} \tag{10.53}$$

Then, from (4.43), we have:

$$C(\varphi_{2,m}^C | u_p; \Delta, \ell \pmod{2m/p^2}) = \sum_{\substack{r \pmod{2m/p} \\ r \equiv \ell \pmod{2m/p^2}}} C(\varphi_{2,m}^C; \Delta p^2, rp \pmod{2m}),$$

$$\begin{aligned}
&= \begin{cases} p|\lambda| & \text{if } \Delta = -\lambda^2, \quad \lambda \equiv \ell \pmod{2m/p^2}, \\ 0 & \text{otherwise.} \end{cases} \\
&= pC(\varphi_{2,m/p^2}^C; \Delta, \ell).
\end{aligned} \tag{10.54}$$

This completes the proof of the fact that  $\Phi_{2,m}^0$  can be chosen to have optimal growth. Now, by formula (4.43), we have that  $\Phi_{2,m}^0|u_t$ , and hence also  $\Phi_{2,m}^0|u_t|U_t$ , is holomorphic for  $t > 1$ , so by replacing  $\Phi_{2,m}^0$  by  $\pi^{\text{prim}}(\Phi_{2,m}^0)$  as defined in (4.46) we can also assume that  $\Phi_{2,m}^0$  is primitive.  $\square$

**Proof of Theorem 9.4:** The proof of Theorem 10.5 given in §10.2 the following discussion already showed that  $J_{2,m}^{\text{OG}}/J_{2,m}$  is one-dimensional and that any representative  $\mathcal{K}_m$  of this quotient space, if we normalize it to have  $c(\mathcal{K}_m; 0, 0) = 2$ , has polar coefficients as given in the statement of Theorem 9.4. Since all the polar coefficients  $C(\mathcal{K}_m; -1, r)$  are equal, we have that  $\mathcal{K}_m|W_{m_1} - \mathcal{K}_m \in J_{2,m}$  for all  $m_1|m$ , so (by averaging) we can choose  $\mathcal{K}_m$  to be invariant under all Atkin-Lehner operators. Finally, we can make  $\mathcal{K}_m$  primitive by the argument used at the end of the proof of Theorem 9.2.  $\square$

**Remark.** For square-free  $m$ , since all holomorphic Jacobi forms are in fact cusp forms,  $\mathcal{K}_m$  is unique up to Jacobi cusp forms, and its coefficients with  $\Delta = 0$  are unique. In this case the only such coefficient, up to translation, is  $c(0, 0)$ . If  $m$  is not square-free, there is more than one coefficient  $c(n, r)$  such that  $\Delta = 0$ . Notice that  $r^2 \equiv 0 \pmod{4m} \Leftrightarrow (r, 2m) = 2m/t$  with  $t^2|m$ . For each  $t$  with  $t^2|m$ , we have

$$\sum_{j \pmod{t}} C(\mathcal{K}_m; 0, \frac{2mj}{t} \pmod{2m}) = \begin{cases} -2 & \text{if } t = 1, \\ 0 & \text{if } t > 1. \end{cases} \tag{10.55}$$

(For  $t = 1$ , the sum on the left reduces to the single term  $c(\mathcal{K}_m; 0, 0) = -2$ , and for  $t > 1$ , it equals  $C(\mathcal{K}_m|u_t; 0, 0)$ , which vanishes because  $\mathcal{K}_m|u_t \in J_{2,m/t^2}$ .) We can choose the  $\Delta = 0$  coefficients arbitrarily subject to this one constraint, and once we have made a choice, then  $\mathcal{K}_m$  is unique up to cusp forms even in the non-square-free case. For instance, at the expense of introducing denominators, we could fix  $\mathcal{K}_m$  uniquely up to cusp forms by requiring that  $C(\mathcal{K}_m; 0, r)$  depends only on the gcd of  $r$  and  $2m$ . (This would correspond to choosing  $a = \frac{1}{2}$  in the first and third tables at the end of §9.3.) Another choice, more similar to our “ $\star = \text{II}$ ” condition, would be to choose  $C(\mathcal{K}_m; 0, r) = 0$  for all  $r$  with  $4m|r^2$  and  $r \nmid 2m$ , in which case the remaining coefficients  $C(\mathcal{K}_m; 0, 2m/t)$  ( $t^2|m$ ) are uniquely determined by (10.55). (This would correspond to choosing  $a = 1$  in the first of the tables at the end of §9.3.) The reader can easily work out the complete formula for all of the coefficients  $C(\mathcal{K}_m; 0, r)$  for either of these two special choices.

Our next task is to prove Theorem 9.5. For this we need to calculate the residues of  $\Phi_{2,m}$ , using the results of the previous subsection. We therefore have to apply the formula (10.44) to

the mock modular form  $\Phi_{2,m} = \varphi_{2,m}^F$ . (Note that the residue of this form will be independent of the specific choice of  $\varphi_{2,m}$ , since any two choices differ by a Jacobi form and the residues of Jacobi forms vanish.) Since the completion of  $\Phi_{2,m}$  involves the weight  $1/2$  unary theta series  $\vartheta_{m,\ell}^0$ , the first step is to compute the scalar products of these theta series.

**Proposition 10.2.** *For  $m \in \mathbb{N}$ , and  $\ell_1, \ell_2 \in \mathbb{Z}/2m\mathbb{Z}$ , we have*

$$(\vartheta_{m,\ell_1}^0, \vartheta_{m,\ell_2}^0) = \frac{1}{2\sqrt{m}} (\delta_{\ell_1, \ell_2 \pmod{2m}} + \delta_{\ell_1, -\ell_2 \pmod{2m}}) \quad (10.56)$$

*Proof.* Formula (3.15) gives

$$(\vartheta_{m,\ell_1}^0, \vartheta_{m,\ell_2}^0) = \frac{1}{2} \operatorname{Res}_{s=\frac{1}{2}} \left( \sum_{r>0} \frac{(\delta_{r,\ell_1(2m)} + \delta_{r,-\ell_1(2m)})(\delta_{r,\ell_2(2m)} + \delta_{r,-\ell_2(2m)})}{(r^2/4m)^s} \right).$$

The proposition follows.  $\square$

**Proposition 10.3.** *Let  $\varphi_{2,m}$  be any meromorphic Jacobi form of weight 2 and index  $m$  with pole  $1/(2\pi iz)^2$  and its translates, and let  $\Phi_{2,m}$  be its finite part  $\varphi_{2,m}^F$ . Then*

$$R_{m_1, m_2}(\Phi_{2,m}) = \frac{m_1 + m_2}{12}. \quad (10.57)$$

*Proof.* By equation (9.5), the  $\ell^{\text{th}}$  component  $g_\ell$  of the shadow for  $\Phi_{2,m}$  is  $-\sqrt{\frac{m}{4\pi}} \vartheta_{m,\ell}^0$ . When  $(m_1, m_2) = 1$ ,  $R_{m_1, m_2}(\Phi_{2,m}) = \mathcal{R}[\Phi_{2,m} | W_{m_1}]$ . Theorem 10.6 then gives:

$$\begin{aligned} \mathcal{R}[\Phi_{2,m} | W_{m_1}] &= \frac{\sqrt{m}}{\sqrt{4\pi}} \frac{\sqrt{\pi}}{6} \sum_{\ell \pmod{2m}} (\vartheta_{m,\ell}^0, \vartheta_{m,\ell^*}^0) \\ &= \frac{1}{24} \sum_{\ell \pmod{2m}} (\delta_{\ell, \ell^* \pmod{2m}} + \delta_{\ell, -\ell^* \pmod{2m}}) \\ &= \frac{1}{12} (m_1 + m_2). \end{aligned}$$

In this calculation, we used Proposition 10.2 to get the second line. For the third line, recall that  $\ell^*$  is defined by  $(\ell^* \equiv -\ell \pmod{2m_1}, \ell^* \equiv \ell \pmod{2m_2})$ , so that  $\ell^* \equiv \ell \pmod{2m} \Leftrightarrow m_1 | \ell$  (which is true for  $2m_2$  values of  $\ell$  in  $\mathbb{Z}/2m\mathbb{Z}$ ), and similarly,  $\ell^* \equiv -\ell \pmod{2m} \Leftrightarrow m_2 | \ell$  (which is true for  $2m_1$  values of  $\ell$ ). When  $(m_1, m_2) = t > 1$ , we have  $R_{m_1, m_2}(\Phi_{2,m}) = R_{m_1/t, m_2/t}(\Phi_{2,m} | u_t)$ . As we have shown in the proof of Theorem 9.2, the shadow of  $\Phi_{2,m} | u_t$  is the same as the shadow of  $t\Phi_{2,m/t^2}$ , so  $R_{m_1/t, m_2/t}(\Phi_{2,m} | u_t) = tR_{m_1/t, m_2/t}(\Phi_{2,m/t^2}) = t(\frac{m_1}{t} + \frac{m_2}{t})/12 = (m_1 + m_2)/12$  by the special case already proved.  $\square$

The reader might find it helpful to compare the statement of this proposition with the table in §9.2 presenting the data for the case  $m = 6$ , in which the function  $\Phi = \Phi_{2,6}$  (in any of its three versions I, II, or III) has

$$\begin{aligned} R_{1,6}(\Phi) &= 2C(\Phi; -1, \pm 1 \pmod{12}) + C(\Phi; 0, 0 \pmod{12}) = 7/12, \\ R_{2,3}(\Phi) &= 2C(\Phi; -1, \pm 5 \pmod{12}) + C(\Phi; 0, 0 \pmod{12}) = 5/12. \end{aligned}$$

**Proof of Theorem 9.5:** We can now complete the proof of Theorem 9.5 and its corollary. For  $m$  square-free, any choice of  $\Phi = \Phi_{2,m}^0$  in  $\mathcal{E}_m^{\text{OG}}$ , and any  $r$  with  $r^2 \equiv 1 \pmod{4m}$ , we have

$$2C(\Phi; -1, \ell) + C(\Phi; 0, 0) = \mathcal{R}[\Phi | W_{m_1}] = \frac{m_1 + m_2}{12} \quad (10.58)$$

by Proposition 10.3, where  $m = m_1 m_2$  is the decomposition of  $m$  for which  $r^* = 1$ . Since  $C(\Phi^{\text{III}}; 0, 0) = 0$ , this proves Theorem 9.5 and also gives the corresponding formulas

$$C(\Phi^{\text{I}}; -1, r) = -\frac{(m_1 - 1)(m_2 - 1)}{12}, \quad C(\Phi^{\text{II}}; -1, r) = -\frac{1}{12} \left( \frac{m_1 + m_2}{2} - \prod_{p|m} \frac{p+1}{2} \right)$$

for  $\Phi^{\text{I}}$  and  $\Phi^{\text{II}}$ , as well as formula (9.43) for the polar coefficients of  $\mathcal{Q}_M$  when  $\mu(M) = 1$ . If  $m$  is not square-free, then the solutions  $r \pmod{2m}$  of  $r^2 \equiv 1 \pmod{4m}$  are still in 1:1 correspondence with the decompositions of  $m$  into coprime factors  $m_1$  and  $m_2$ . To compute the corresponding Fourier coefficient of  $\Phi_{2,m}^0$ , we look at each term in (10.52) separately, writing each  $d$  as  $d_1 d_2$  with  $d_i^2 | m_i$  and noting that  $\frac{m_1}{d_1^2} = \left(\frac{r-1}{2}, \frac{m}{d^2}\right)$ ,  $\frac{m_2}{d_2^2} = \left(\frac{r+1}{2}, \frac{m}{d^2}\right)$ , after which (9.41) follows from (10.57) and the fact that  $\mathcal{R}[\varphi | U_d] = \mathcal{R}[\varphi]$ .  $\square$

**Proof of Theorem 9.7:** It is enough to show that any weak Jacobi form of optimal growth is actually holomorphic since, as we have seen in Theorem 10.3, there are no holomorphic Jacobi forms of weight 1. To show this, we use the “residue operator on weight 1 forms” defined by

$$\mathcal{R}^{(1)}[\varphi] = \text{Res}_{\tau=\infty}(\varphi'(\tau, 0) d\tau) = \sum_{r \in \mathbb{Z}} r c_\varphi(0, r) \quad (10.59)$$

instead of (10.17), where, as usual,  $\varphi'(t, z) = \frac{1}{2\pi i} \frac{d}{dz} \varphi(t, z)$ . This residue vanishes if  $\varphi \in \tilde{J}_{1,m}$ , since then  $\varphi'(\cdot, 0) \in M_2^!$ . If, further,  $\varphi$  has optimal growth, then  $2c_\varphi(0, 1) = \mathcal{R}^{(1)}[\varphi] = 0$ , so  $C_\varphi(-1, 1)$  vanishes. By the same argument with  $\varphi$  replaced by  $\varphi$  acted upon by each of the Atkin-Lehner operators, one has that  $C_\varphi(-1, r) = 0$  for all  $r$  with  $r^2 \equiv -1 \pmod{4m}$ , and thus  $\varphi \in J_{1,m}$ .  $\square$

## 11. Quantum black holes and mock modular forms

We now have all the ingredients required to address the questions posed in the introduction. These answers follow from applying our results of §8, in particular theorem 8.3, to the quarter-BPS partition function of the the  $\mathcal{N} = 4$  theory. It turns out that in this case, many of the

mathematical objects in the theorem simplify. A reader who has not gone through the details of §8 could still fruitfully read this section to obtain the relevant results for this example.

In particular, we show in §11.1 that the immortal black hole degeneracies extracted from the asymptotic partition function (6.14) are Fourier coefficients of a (mixed) mock Jacobi form. The completion transforms as a true Jacobi form and obeys a first order partial differential equation (11.23) which can be understood as a holomorphic anomaly equation. In §11.2 we discuss the physical origin of the meromorphy of the asymptotic counting function in this example and how it is related to the noncompactness of the microscopic brane dynamics. In §11.3 we analyze the contour corresponding to the M-theory limit to embed our analysis in the context of  $AdS_3/CFT_2$  holography near the horizon of a black string. We conclude in §11.4 with comments and open problems.

### 11.1 Mock Jacobi forms, immortal black holes, and $AdS_2$ holography

We first discuss the consequences of the  $S$ -duality symmetry and its relation to the spectral flow symmetry for the moduli-independent immortal degeneracies  $d^*(n, \ell, m)$  defined by (6.14). Recall that under the S-duality transformations

$$\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \tag{11.1}$$

with integer  $b$ , the S-modulus (2.21) transforms as:

$$S \rightarrow S + b. \tag{11.2}$$

which means that the axion transforms as  $a \rightarrow a + b$ . Since the axion arises from a dimensional reduction of the 3-form field in the M-theory frame (discussed in §11.3), this subgroup (11.1) of  $S$ -duality has its origins in large gauge transformations of the 3-form field. When the axion transforms as above, the effective theta angle changes and as a result the charge vector of the dyon transforms as  $(Q, P) \rightarrow (Q + Pb, P)$  due to the Witten effect [111]. Consequently, the charge-invariants transform as

$$n \rightarrow n + mb^2 + b\ell, \quad \ell \rightarrow \ell + 2mb, \quad m \rightarrow m. \tag{11.3}$$

which are nothing but the spectral flow transformations. Note that  $m$  is left fixed by this subgroup of S-duality whereas other elements of the S-duality group would change  $m$ . Thus, only this subgroup is expected to act on the sector with a fixed  $m$ . The covariance of the spectrum under these  $S$ -duality transformations implies

$$d(n, \ell, m)|_{S, \mu} = d(n + mb^2 + b\ell, \ell + 2mb, m)|_{S+b, \mu}. \tag{11.4}$$



Since the degeneracies of immortal black holes are independent of the moduli by definition, they are actually invariant under  $S$ -duality:

$$d^*(n + mb^2 + b, \ell + 2mb, m) = d^*(n, \ell, m). \quad (11.5)$$

From the mathematical point of view, this is precisely the action of the elliptic part of the Jacobi transformations (4.2) on the Fourier coefficients of a Jacobi (or mock Jacobi) form.

The immortal degeneracies  $d^*(n, \ell, m)$  are computed using the attractor contour (6.13), for which the imaginary parts of the potentials are proportional to the charges and scale as

$$\text{Im}(\sigma) = 2n/\varepsilon, \quad \text{Im}(\tau) = 2m/\varepsilon, \quad \text{Im}(z) = -\ell/\varepsilon, \quad (11.6)$$

with  $\varepsilon$  very small and positive. In other words,

$$|p| = \lambda^{2n}, \quad |q| = \lambda^{2m}, \quad |y| = \lambda^{-\ell}, \quad \text{with } \lambda = \exp(-2\pi/\varepsilon) \rightarrow 0 \quad (11.7)$$

on the attractor contour. We assume that  $n > m$  without loss of generality, otherwise one can simply exchange  $\sigma$  and  $\tau$ . Moreover, using the spectral flow symmetry (11.5) we can always bring  $\ell$  to the window  $0 \leq \ell < 2m$  for the immortal degeneracies.

To extract the Fourier coefficient from the product representation (5.13) of the counting function (5.14) using (11.7), we need to expand a typical term in the product of the form

$$\frac{1}{(1 - p^r q^s y^t)^{2C_0(4rs - t^2)}}. \quad (11.8)$$

where  $2C_0$  is the Fourier coefficient of the elliptic genus of a single copy of  $K3$ . Depending on whether  $|p^r q^s y^t|$  is less than one or greater than one, we expand the denominator using

$$\frac{1}{1 - x} = \begin{cases} 1 + x + x^2 + \dots & \text{for } |x| < 1, \\ -(x^{-1} + x^{-2} + x^{-3} + \dots) & \text{for } |x| > 1. \end{cases} \quad (11.9)$$

We see that it is not *a priori* obvious that expanding first in  $p$  is the same as using the attractor contour because if the powers of  $t$  are sufficiently large and positive, the attractor contour may correspond to expanding some terms effectively in  $p^{-1}$  instead of in  $p$ .

To address this question we now show that for the terms that appear in the product representation,  $|p^r q^s y^t|$  is always less than one using the fact that

$$C_0(\Delta) = 0 \quad \text{for } \Delta < -1. \quad (11.10)$$

To begin with, the  $p$ -dependent terms with  $\Delta = 0, -1$  arise only for  $s = 0$  and are of the form

$$\frac{1}{(1 - py)^2 (1 - py^{-1})^2 (1 - p^r)^{20}} \quad (11.11)$$

These involve only single powers  $y$  or  $y^{-1}$  and positive powers of  $p$ . The absolute value of  $py$  or  $py^{-1}$  for the attractor contour goes as  $\lambda^{2m\pm l}$  which is always much less than one for  $0 \leq |l| < 2m$ . The remaining  $p$ -dependent terms with  $s > 0$  have  $\Delta = 4rs - t^2 > 0$  and  $r > 0$ . The absolute value of  $p^r q^s y^t$  goes as

$$\lambda^{2nr+2ms-lt}. \quad (11.12)$$

The exponent is most negative when  $\ell$  takes its maximum positive value  $\ell_{max} = \sqrt{2mn}$  and  $t$  takes its maximum positive value  $t_{max} = \sqrt{2rs}$ . For black hole configurations with a smooth horizon,  $4mn - \ell^2 > 0$  because the classical area of the horizon is the square-root of this quantity. Hence the most negative value of the exponent would be

$$2nr + 2ms - 2\sqrt{mnr s} = 2(\sqrt{nr} - \sqrt{ms})^2 > 0, \quad (11.13)$$

and the absolute value of  $p^r q^s y^t$  is always less than or equal to

$$\lambda^{2(\sqrt{nr} - \sqrt{ms})^2} \quad (11.14)$$

which is always much less than one. This implies that to extract the attractor Fourier coefficients, we expand all terms in the product in the small  $p$  expansion using (11.9) thus giving us the Jacobi form  $\psi_m$  of (1.1) as a partition function in the remaining two chemical potentials.

The degeneracies extracted from  $\psi_m(\tau, z)$  still experience many wall-crossings, and we should now use the rest of the attractor contour to analyze the immortal part of the degeneracies of  $\psi_m$ . We do this in two steps by first performing the inverse Fourier transform in  $z$ , and then in  $\tau$ . For the first step, we need to specify a contour for the imaginary part of  $z$  relative to that of  $\tau$ . The conditions (11.6) on the imaginary parts of  $(\tau, z)$  implies that  $\text{Im}(z) = -\frac{\ell}{2m}\text{Im}(\tau)$ . Since by moving the contour in a purely horizontal direction we do not cross any poles, we can add an arbitrary real part to  $z$ , which we can choose so that  $z = -\ell\tau/2m$ . This gives the result of the first integral to be:

$$f_{m,\ell}^*(\tau) = e^{-\pi i \ell^2 \tau / 2m} \int_P^{P+1} \psi_m(\tau, z) e^{-2\pi i \ell z} dz, \quad P = -\ell\tau/2m. \quad (11.15)$$

We then have to perform a  $\tau$  integral with the contour specified by the attractor values (11.6), but since the integrand  $f_{m,\ell}^*(\tau)$  is holomorphic in  $\tau$ , the answer is independent of the contour. We thus get the degeneracies  $d^*(n, \ell, m)$  to be:

$$d^*(n, \ell, m) = \int e^{\pi i \ell^2 \tau / 2m} f_{m,\ell}^*(\tau) d\tau, \quad (11.16)$$

where the integral is over an interval of length 1 for the fixed imaginary value (11.6) of  $\tau$ .

To extend to other values of  $\ell$ , we use the spectral flow invariance (11.5) and sum over all values of  $\ell$  to get a canonical partition function in the two variables  $(\tau, z)$ . Because of

the factor  $e^{\pi i \ell^2 \tau / 2m}$  in (11.16), the sum over spectral-flow images yields the function  $\vartheta_{m,\ell}(\tau, z)$ . Putting all this together, and comparing the above equations (11.15), (11.16) with our definition (8.3) of the finite part of a meromorphic Jacobi form, we see that the single centered black hole degeneracies are Fourier coefficients of precisely the *finite* part of  $\psi_m$  as defined in §8:

$$\sum_{\ell \bmod (2m)} f_{m,\ell}^*(\tau) \vartheta_{m,\ell}(\tau, z) = \psi_m^F(\tau, z). \quad (11.17)$$

We see that the seemingly unconventional charge-dependent choice for the contour in §8 for defining the finite part is completely natural from the point of view of the attractor mechanism (indeed, it was motivated there by these physical considerations). According to our analysis in §8, this finite part  $\psi_m^F$  is a *mock* Jacobi form. This is one of the main conclusions of our analysis, so we summarize the entire discussion by the following concise statement:

*The degeneracies of single centered (immortal) black holes with magnetic charge invariant  $M^2/2 = m$  are Fourier coefficients of a mock Jacobi form of index  $m$ .*

We stress here that the converse is not true. The single centered black hole solution exists only when the discriminant  $4mn - \ell^2$  is positive, and its degeneracy grows exponentially for large values of the discriminant. On the other hand, as we saw in §4, any Jacobi form with exponential growth necessarily has negative discriminant states. This means that the partition function for single centered black holes alone with fixed magnetic charge and varying electric charge cannot have good modular properties. Indeed the partition function  $\psi_m^F$  does contain negative discriminant states for example for  $n < m$ . These low-lying states do not correspond to single centered black holes [27, 97].

We now turn to the physical interpretation of the polar part  $\psi_m^P(\tau, z)$  of the meromorphic Jacobi form defined in (8.44). As mentioned in the introduction, the only multi-centered gravitational configurations that contribute to the supersymmetric index of quarter-BPS dyons in our  $\mathcal{N} = 4$  string theory have exactly two centers [33] each of which is a half-BPS state. At a wall, one of these two-centered configurations decays. A basic decay at a wall-crossing is when one center is purely electric and the other is purely magnetic. Other decays are related to this basic decay by duality [94]. For this two-centered configuration, with one electric and one magnetic center, the indexed partition function after summing over the electric charges is:

$$p_{24}(m+1) \cdot \frac{1}{\eta^{24}(\tau)} \cdot \sum_{\ell > 0} \ell y^\ell, \quad (11.18)$$

where the first factor is the degeneracy of the half-BPS magnetic center, the second factor is the partition function of the electric center [34], and the third factor is the partition function counting the dimensions of the  $SU(2)$  multiplets with angular momentum  $(\ell - 1)/2$  contained

in the electromagnetic field generated by the two centers [41] for which  $N \cdot M = \ell$ . The third factor is an expansion of the function

$$\frac{y}{(1-y)^2} \quad (11.19)$$

in the range  $|q| < |y| < 1$  (see equation (9.7)). Here we already see the basic wall-crossing of the theory because different ways of expanding this meromorphic function for  $|y| < 1$  and  $|y| > 1$  will give different degeneracies. Other wall-crossings are related to this basic wall-crossing by S-duality which as we have explained is nothing but spectral flow for a given fixed  $m$ . Thus the full partition function that captures all wall-crossings is obtained by ‘averaging’ over the spectral-flow images of (11.19). This averaging is nothing but the averaging operation defined in (9.3) and (9.4)

$$\text{Av}^{(m)} \left[ \frac{y}{(y-1)^2} \right] = \frac{q^{ms^2+s} y^{2ms+1}}{(1-q^s y)^2} \quad (11.20)$$

It is now easy to see that this averaging over spectral-flow images gives exactly the polar part  $\psi_m^{\text{P}}$  of the meromorphic Jacobi form  $\psi_m$ :

$$\frac{p_{24}(m+1)}{\eta^{24}(\tau)} \sum_{s \in \mathbb{Z}} \frac{q^{ms^2+s} y^{2ms+1}}{(1-q^s y)^2} = \psi_m^{\text{P}}(\tau, z). \quad (11.21)$$

This follows from the fact that the poles of  $\psi_m(\tau, z)$  are exactly at  $z \in \mathbb{Z}\tau + \mathbb{Z}$  and the residue at  $z = 0$  is

$$\frac{p_{24}(m+1)}{\eta^{24}(\tau)}.$$

as mentioned in (1.5).

We thus see that the decomposition theorem (8.3) as applied to our dyon partition function  $\psi_m(\tau, z)$  has a natural physical interpretation. It is simply the statement that the full partition function is a sum of its immortal and decaying constituents:

$$\psi_m(\tau, z) = \psi_m^{\text{F}}(\tau, z) + \psi_m^{\text{P}}(\tau, z). \quad (11.22)$$

The nontrivial part is of course the implication of such a decomposition for modularity. By separating part of the function  $\psi_m$ , we have evidently broken modularity, and  $\psi_m^{\text{F}}(\tau, z)$  is not a Jacobi form any more. However, the decomposition theorem (8.3) guarantees that  $\psi_m^{\text{F}}$  still has a very special modular behavior in that it is a mock Jacobi form. The mock behavior can be summarized by the following partial differential equation obeyed by its completion:

$$\tau_2^{3/2} \frac{\partial}{\partial \bar{\tau}} \widehat{\psi}_m^{\text{F}}(\tau, z) = \sqrt{\frac{m}{8\pi i}} \frac{p_{24}(m+1)}{\Delta(\tau)} \sum_{\ell \bmod (2m)} \overline{\vartheta_{m,\ell}(\tau, 0)} \vartheta_{m,\ell}(\tau, z). \quad (11.23)$$

Note that  $\psi_m^P$  also admits a completion  $\widehat{\psi}_m^P$  which is modular. However, the function  $\widehat{\psi}_m^P$ , unlike  $\widehat{\psi}_m^F$ , is meromorphic, and therefore has wall-crossings. We refer the reader to the summary at the end of §8.3 for a detailed discussion of the relationship between meromorphy (in  $z$ ), holomorphy (in  $\tau$ ) and modularity.

Let us now discuss the implications of these results for  $AdS_2/CFT_1$  holography. The near horizon geometry of a supersymmetric black hole is a two-dimensional anti de Sitter space  $AdS_2$  and the dual theory is a one-dimensional conformal field theory  $CFT_1$ . The partition function of this  $CFT_1$  is nothing but the integer  $d^*(n, \ell, m)$  which gives the number of microstates of the black hole [95, 96]. Our results show that that these integers are the Fourier coefficients of a mock modular form and as a result there is a hidden modular symmetry. Such a modular symmetry would make it possible to use powerful techniques such as the Hardy-Ramanujan-Rademacher expansion to process the integers  $d^*(n, \ell, m)$  into a form that can be more readily identified with a bulk partition function of string theory in  $AdS_2$ . There has been recent progress in evaluating the bulk partition function using localization techniques which makes it possible to study the subleading nonperturbative corrections in the Rademacher expansion in a systematic way [30, 31]. It would be interesting to see if the additional terms in the Rademacher expansion arising from the ‘mock’ modular nature can be given a physical bulk interpretation.

## 11.2 Meromorphy, mock modularity, and noncompactness

From the mathematical point of view, we have seen how the mock modularity of  $\psi_m^F$  is a consequence of meromorphy of the Jacobi form  $\psi_m$ . In the  $\mathcal{N} = 4$  example above, we can understand the physical origin of this meromorphy as coming from the non-compactness of the underlying brane dynamics. To see this more clearly, we write the meromorphic Jacobi form  $\psi_m(\tau, z)$  as

$$\psi_m(\tau, z) = \frac{1}{\varphi_{-2,1}(\tau, z)} \frac{1}{\Delta(\tau)} \chi(\tau, z; \text{Sym}^{m+1}(K3)). \quad (11.24)$$

The three factors in the above equations have the following physical interpretation [37]. The function  $\psi_m(\tau, z)$  can be viewed as an indexed partition function of the following (0, 4) SCFT

$$\sigma(\text{TN}) \times \sigma_L(\widetilde{K}-n) \times \sigma(\text{Sym}^{m+1}(K3)). \quad (11.25)$$

Each CFT factor has a natural interpretation from the microscopic derivation of the dyon counting formula [37] using the 4d-5d lift [55]. The  $\sigma_L(\widetilde{K}-n)$  factor is the purely left-moving bosonic CFT associated with  $\widetilde{K}-n$  bound states of the Kaluza-Klein monopole with momentum. The  $\sigma(\text{TN})$  factor is a (4, 4) SCFT associated with the motion of the center of mass of motion in the Kaluza-Klein background in going from five dimensions to four dimensions in the 4d-5d lift. The  $\sigma(\text{Sym}^{m+1}(K3))$  is the (4, 4) SCFT that is associated with the five-dimensional Strominger-Vafa  $Q1-Q5-n$  system. Note that the SCFT  $\sigma(\text{TN})$  has a *noncompact* target space and the double pole of  $1/\varphi_{-2,1}(\tau, z)$  can be traced back to this noncompactness.

Going over from the partition function to the Fourier coefficients corresponds to going over from the canonical ensemble with fixed  $z$  to the micro-canonical ensemble with fixed  $\ell$ . The Fourier coefficient is supposed to count the number of right-moving ground-states in this fixed charge sector. Right-moving states with nonzero energy cancel in pairs in this index. Now, counting ground states for a compact theory is a well-defined operation. However, when the target-space is noncompact, the wavefunctions are not properly normalized without putting an infrared regulator in the target space. In the present context, the essential physics of the noncompactness is captured by the point particle motion of the center of mass in the Taub-NUT geometry. A simple regulator can be introduced by turning on the axion field which according to (2.22) corresponds in the Type-IIB frame to taking the modulus  $U_B = U_{1B} + iU_{2B}$  to have a small nonzero real part  $U_{1B}$  [37]. This introduces a nonzero potential on the Taub-NUT geometry proportional to  $U_{1B}^2 U_{2B}$  that regulates the infrared behavior to give a well defined counting problem [57, 89]. However, the index thus computed depends on the sign of  $U_{1B}$  and jumps as  $U_{1B}$  goes from being slightly negative to slightly positive. Indeed the partition function for these modes is precisely the function

$$\frac{y}{(1-y)^2} \tag{11.26}$$

which can be expanded either in power of  $y$  or in powers of  $y^{-1}$  depending on whether  $U_{1B}$  is positive or negative. For  $\ell$  positive, there are  $\ell$  normalizable states when  $U_{1B} < 0$  and none when  $U_{1B} > 0$ . This is the essential source of the wall-crossing behavior in this context. We should emphasize that even though turning on  $U_{1B}$  appears to make the problem effectively compact, this is true only up to some maximal  $\ell_{max}$  which scales as  $U_{1B}^2 U_{2B}$  [57, 89]. For a given  $\ell$  positive, one can always make the problem well-defined with a compact target space by choosing the moduli so that  $\ell_{max}$  is larger than the  $\ell$  of interest. The partition function in that case is well-defined without any poles and goes as

$$y + 2y^2 + \dots + \dots + \ell_{max} y^{\ell_{max}} . \tag{11.27}$$

However, if one wishes to compute the canonical partition function that sums over all  $\ell$ , one has to essentially take  $U_{1B}^2 U_{2B}$  to go to infinity which is what leads to the pole at  $y = 1$  in (11.26). This in essence is the origin of the meromorphy of the canonical counting function of the asymptotic states, and therefore indirectly also of the mock modularity of the immortal near-horizon counting function.

As we have seen, this meromorphy of the asymptotic counting function  $\psi_m$  implies that the counting function of the near-horizon immortal states cannot be both modular and holomorphic at the same time. One can either have the holomorphic function  $\psi_m^F$  which is not quite modular, or a modular function  $\widehat{\psi}_m^F$  which is not quite holomorphic. From this point of view,

the shadow can be regarded as a ‘modular anomaly’ in  $\psi_m^F$  or as a ‘holomorphic anomaly’ in the completion  $\widehat{\psi}_m^F$ .

A natural question that arises from this discussion is if the mock Jacobi form and the holomorphic anomaly can be given a direct physical interpretation rather than arrive at it via a meromorphic Jacobi form as we have done. To address this question, it is useful to view the problem from the perspective of  $AdS_3$  holography in M-theory as we now discuss.

### 11.3 The M-Theory limit and $AdS_3$ holography

From the perspective of  $AdS_2/CFT_1$  holography discussed above, there is no *a priori* reason why the black hole degeneracy should have anything to do with modularity. It is only when we can view the black hole as an excitation of a black string that we can have an *a priori* physical expectation of modularity. This is because the near horizon geometry of a black string is a three-dimensional anti de Sitter space  $AdS_3$ . The Euclidean thermal  $AdS_3$  has a conformal torus as a boundary whose complex structure parameter is  $\tau$ . The partition function of the dual boundary conformal field theory  $CFT_2$  depends on  $\tau$ . In this framework the modular symmetry is identified with the  $SL(2, \mathbb{Z})$  mapping class group of the boundary torus [78], whereas the elliptic symmetry is identified with large gauge transformations of the 3-form field [39] as we have already discussed in §11.1.

We now describe how the quarter-BPS black holes considered thus far can be regarded as excitations of a black string in an appropriate M-theory frame. This will enable us to make contact with  $AdS_3/CFT_2$  holography. The M-theory framework is natural also from the point of view of generalizing these considerations to BPS black holes in  $\mathcal{N} = 2$  compactifications on a general Calabi-Yau manifold  $X_6$ .

To go to the M-theory frame, we consider the  $T^2$  to be a product of two circles  $S^1 \times \tilde{S}^1$  and consider the charge configuration (2.15) in the Type-IIB frame. We first  $T$ -dualize the  $\tilde{S}^1$  circle to the  $\hat{S}^1$  circle to go to the Type-IIA frame and then use mirror symmetry of  $K3$  to map the Type-IIB configuration (2.15) to a Type-IIA configuration of D4-F1-n-NS5 charges. We then lift it to M-theory to obtain a charge configuration consisting of M5-M2 bound states carrying momentum along  $S^1$ . This configuration now has M5-branes wrapping  $D \times S^1$  where  $D$  is a divisor in  $K3 \times \hat{T}^2$ , an M2-brane wrapping the  $\hat{T}^2 = \hat{S}^1 \times S_m^1$ , with some momentum along  $S^1$ . This is a precisely a configuration of the kind considered by Maldacena, Strominger, and Witten [79]. Since all M5-branes wrap  $S^1$  and the momentum is along  $S^1$ , it is natural flip  $S_m^1$  and  $S^1$  and regard the  $S^1$  as the new M-circle. In the limit when the radius  $R$  of the new M-circle  $S^1$  is very large, the low-lying excitations of this M5-brane are described by the MSW string wrapping the M-circle. To write the final configuration in the M-theory frame, let  $C^1$  to be the homology 2-cycle of the  $\hat{T}^2$ , and let  $C^2$  and  $C^3$  be two 2-cycles in  $K3$  with intersection



matrix

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (11.28)$$

The intersection number of these three 2-cycles is

$$\int_{K3 \times \hat{T}^2} C^1 \wedge C^2 \wedge C^3 = 1. \quad (11.29)$$

Let  $\{D_a\}$  be the 4-cycles dual to  $\{C^a\}$ ,  $D_a \cap C^b = \delta_a^b$ . In this basis, the M5-brane is wrapping the 4-cycle

$$D = \sum_{a=1}^3 p^a D_a$$

with  $p^1 = \tilde{K}$ ,  $p^2 = Q_1$  and  $p^3 = Q_5$ . The M2-brane charges are  $q_2 = q_3 = 0$  and  $q_1 = \tilde{n}$ , and M-momentum or equivalently the D0-brane charge from the Type-IIA perspective is  $q_0 = n$ . Using this notation we see that

- The electric charges in (2.15) map to  $n$  units of momentum along the M-circle  $S^1$ , and  $\tilde{K}$  M5-branes wrapping  $D_1 \times S^1$ .
- The magnetic charges in (2.15) map to  $Q_1$  M5-branes wrapping  $D_2 \times S^1$ ,  $Q_5$  M5-branes wrapping  $D_3 \times S^1$ , and  $\tilde{n}$  M2-branes wrapping  $\hat{T}^2$ .

In summary, the charge configuration of our interest is

$$\Gamma = \begin{bmatrix} N \\ M \end{bmatrix} = \begin{bmatrix} 0, & n; & 0, & \tilde{K} \\ Q_1, & \tilde{n}; & Q_5, & 0 \end{bmatrix}_B = \begin{bmatrix} 0, & q_0; & 0, & p^1 \\ p^2, & q_1; & p^3, & 0 \end{bmatrix}_M \quad (11.30)$$

Reduction of this charge configurations along  $S^1$  gives a configuration in Type-IIA frame consisting of only D4-D2-D0-branes.

Since the effective MSW string wraps the M-circle, we have to take the radius  $R$  of this circle to be large keeping other scales fixed to obtain an  $AdS_3$  near-horizon geometry of a long black string. This implies that in the original heterotic frame in which we have labeled the charges (2.3), the radius  $R$  goes to infinity keeping other scales and moduli fixed. Since the 2-torus in the heterotic frame is also a simple product of two circles  $S^1 \times \tilde{S}^1$ , the moduli in (2.20) take the form

$$G_{ij} = \begin{pmatrix} \tilde{R}^2 & 0 \\ 0 & R^2 \end{pmatrix}, \quad B_{ij} = 0 \quad (11.31)$$

which implies that for the three moduli  $S, T, U$  we have

$$S = a + i \frac{R\tilde{R}}{g_6^2}, \quad T = iR\tilde{R}, \quad U = i \frac{R}{\tilde{R}}, \quad (11.32)$$



where  $g_6$  is the six-dimensional string coupling constant of heterotic compactified on  $T^4$  and we have allowed a nonzero axion field  $a$ . This means in particular that

$$S_2 \sim R, \quad T_2 \sim R, \quad U_2 \sim R \quad (11.33)$$

and for the charge configuration (11.30), the central charge matrix scales as

$$\mathcal{Z} \sim \begin{pmatrix} R & 0 \\ 0 & 0 \end{pmatrix} + \frac{1}{R} \begin{pmatrix} 2n & \ell + ag_6^2/\tilde{R} \\ \ell + ag_6^2/\tilde{R} & 2m + (Q_1^2/\tilde{R}^2 + Q_5^2\tilde{R}^2) \end{pmatrix}. \quad (11.34)$$

From the contour prescription (6.10), we conclude that the imaginary parts for the Fourier integral scale as

$$\text{Im}(\sigma) \sim R \quad (11.35)$$

$$\text{Im}(\tau) \sim 1/R \quad (11.36)$$

$$\text{Im}(z) \sim 1/R. \quad (11.37)$$

Therefore, in the region of the moduli space corresponding to the M-theory limit,  $p := \exp(2\pi i\sigma)$  is becoming much smaller compared to  $q := \exp(2\pi i\tau)$  and  $y := \exp(2\pi iz)$ . Hence one must first expand around  $p = 0$ :

$$\frac{1}{\Phi_{10}(\Omega)} = \sum_{m=-1}^{\infty} p^m \psi_m(\tau, z). \quad (11.38)$$


Since the function  $\Phi_{10}(\Omega)$  has double zeros at  $z = 0$ , the function  $\psi_m(\tau, z)$  is meromorphic with double poles at  $z = 0$  and its translates.

We conclude that the function  $\psi_m(\tau, z)$  can be interpreted as the asymptotic partition function for counting the BPS excitation of this MSW M5-brane for fixed value of the magnetic charge  $M^2/2 = m$  and is a *meromorphic* Jacobi form of weight  $-10$  and index  $m$ . This is the essential new ingredient compared to other situations encountered earlier such as the partition function of the D1-D5 string [107] which is a *holomorphic* weak Jacobi form<sup>26</sup>.

Note that  $\psi_m(\tau, z)$  is the asymptotic counting function in the canonical ensemble for a fixed chemical potential  $z$  where one sums over all M2-brane charges  $\ell$  for a fixed chemical potential  $z$ . To obtain the micro-canonical partition function for a fixed M2-brane charge, one has to perform the Fourier integral in  $z$ . There is an infinity of double poles in the  $z$  plane at  $z = \alpha\tau$  for  $\alpha \in \mathbb{Z}$ . The  $z$  contour can be made to lie between the poles at  $(\alpha - 1)\tau$  and  $\alpha\tau$  depending on how  $\text{Im}(z)$  compares with  $\text{Im}(\tau)$  given the  $O(1/R)$  terms in the central charge matrix (11.34). It is easy to see that by varying the moduli, the contour can be made to cross all poles. Thus, there are an infinity of walls still accessible in the regime when the radius of the M-circle becomes large.

---

<sup>26</sup>Here we mean holomorphic or meromorphic in the  $z$  variable that is conjugate to  $\ell$ . This should not be confused with the nomenclature used in §4 while discussing growth conditions (4.5).



**Figure 1:** The diagram on the left shows the moduli space projected onto the  $\Sigma$  upper-half plane divided into chambers separated by walls where a quarter-BPS state decays into two half-BPS states. The M-theory limit, shown on the right, corresponds to taking  $\text{Im}(\Sigma)$  to be very large. In this limit several walls are no longer visible. There are still an infinite number of walls which can be crossed by varying  $\text{Re}(\Sigma)$ . Spectral flow transformation maps one chamber to another chamber.

To display this graphically it is useful to define a complex scalar field  $\Sigma = \Sigma_1 + i\Sigma_2$  by

$$z \sim \frac{1}{\Sigma_2} \begin{pmatrix} |\Sigma|^2 & \Sigma_1 \\ \Sigma_1 & 1 \end{pmatrix}. \quad (11.39)$$

For fixed charges, the MSW region which corresponds to taking  $\Sigma_2$  large with  $\Sigma_1$  fixed corresponding to vanishing axion. Varying  $\Sigma_1$  allows one to access the infinite number of chambers separated by walls as shown in Fig. 1. Degeneracies in these different chambers can be obtained simply by choosing the Fourier contour in  $z$  for an appropriate value of  $\alpha$ . Note that the field  $\Sigma_1$  depends both on the moduli and on the charges and thus varies as moduli are varied for fixed charges or as the charges are varied for fixed moduli. In particular, even for fixed values of moduli, one can cross all walls by varying  $\ell$  appropriately.

With this embedding into M-theory, we now outline what we regard as a consistent physical picture to interpret  $\widehat{\psi}_m^F$  even though we do not yet fully understand the details. In the limit of large radius  $R$ , we effectively have a five-dimensional theory obtained from compactifying M-theory on  $K3 \times \widehat{T}^2$ . For a given a set of M5-brane charges  $\{p^i\}$  at asymptotic infinity, the five-dimensional supergravity admits a single-centered black string solution carrying magnetic charges  $\{p^i\}$ . The near horizon geometry of this solution is an  $AdS_3$  geometry where all vector multiplet moduli of the five-dimensional supergravity are fixed to their attractor values. Following the usual reasoning of holography, we expect that the superconformal field theory describing the low energy excitations of the M5-brane will be dual to this near horizon  $AdS_3$ .

The indexed partition function of this SCFT is expected to describe the BPS gravitational configurations inside the  $AdS_3$ .

What is the indexed partition function of this SCFT? It cannot possibly be the meromorphic Jacobi form  $\psi_m$  that exhibits many wall-crossings, because all vector-multiplet moduli are fixed in the attractor  $AdS_3$  geometry and no wall-crossings are possible. It cannot possibly be  $\psi_m^F$ , because it is not modular as would be expected from the symmetry under the  $SL(2, \mathbb{Z})$  mapping class group of the boundary torus. We propose that the most natural object to identify with the indexed partition function of the  $SCFT_2$  dual to the  $AdS_3$  is the modular completion  $\widehat{\psi}_m^F(\tau, z)$ . It satisfies (in a rather nontrivial way) both requirements of being properly modular and not having any wall-crossings. The non-holomorphy is not something that we would naively expect for the indexed partition function but this can be a reflection of noncompactness as we comment upon later. The quarter-BPS black holes discussed in this paper are excitations inside this single  $AdS_3$  represented as BTZ black holes and their degeneracies should then be captured by the Fourier coefficients of  $\psi_m^F(\tau, z)$  which is indeed the case for  $n > m$ .

This *microscopic* picture is consistent with the *macroscopic* analysis of multi-centered black string solutions [40] of five dimensional supergravity. These solutions can be obtained by lifting multi-centered black hole solutions of four-dimensional supergravity. According to this supergravity analysis, given a set of asymptotic magnetic charges  $\{p^i\}$  of a wrapped M5-brane, the single-centered  $AdS_3$  throat described above is not the only allowed solution [40]. There exist multi-centered solutions with multiple  $AdS_3$  throats with the same total asymptotic charge. Now, as we have seen from the microscopic analysis, there are still infinite chambers separated by walls visible in the M-theory limit as shown in Fig 1. These chambers can thus correspond to chambers in which different multi-string configurations exist as stable solutions. The wall-crossings can result not from decays of multi-centered black holes inside a single  $AdS_3$  throat but from the decays of these multi-string solutions. Thus, it is reasonable to identify  $\widehat{\psi}_m^F$ , which has no wall-crossings, with the (indexed) partition function of the CFT dual to the single  $AdS_3$  throat.

#### 11.4 Open problems and comments

The interpretation of  $\widehat{\psi}_m^F(\tau, z)$  proposed above raises a number of interesting questions. The essentially new feature of our proposal is the fact that the modular partition function  $\widehat{\psi}_m^F(\tau, z)$  is nonholomorphic and correspondingly the holomorphic counting function  $\psi_m^F$  is mock modular. It would clearly be important to understand the physical origin of this nonholomorphy and mock modularity from the boundary and bulk perspectives.

- *Mock modularity from the boundary perspective:*

We have seen in §11.2 that the mock modularity of  $\psi_m^F$  is tied to the noncompactness of the asymptotic SCFT. A more interesting question is to understand this mock modularity

directly from the point of view of the near horizon SCFT. It is natural to assume that the near horizon SCFT is also noncompact and this noncompactness is what is responsible for the mock modularity. The connection between mock modularity/nonholomorphy and noncompactness has been noted earlier in physics. For example, the partition function of topological  $SO(3)$   $\mathcal{N} = 4$  Yang-Mills theory on  $\mathbb{CP}^2$  is a mock modular form and it was suggested in [109] that it is related to the noncompactness of flat directions in field space. The nonholomorphy of the elliptic genus has been noted also in the context superconformal field theories [49, 108] with noncompact  $SL(2, \mathbb{R})$  as the target space. In this context, the nonholomorphy can be understood in complete detail [108, 4] is a consequence of the continuum of spectrum of the noncompact SCFT. It would be interesting to see if the holomorphic anomaly (11.23) of the completion of  $\widehat{\psi}_m^F$  be understood from such a path integral perspective.

In the framework of  $AdS_2$  holography, the quantity of interest is the integer  $d^*(n, m, \ell)$  which is identified with the partition function of the boundary  $CFT$ . Using the fact that the completion  $\widehat{\psi}_m^F(\tau, z)$  is modular, it should be possible to develop a Rademacher-like expansion for the  $d^*(n, m, \ell)$  along the lines of [13, 14, 15].

- *Mock modularity from the bulk perspective:*

If mock modularity is indeed a consequence of the noncompactness of the boundary  $CFT$ , what are the implications of this noncompactness in the bulk  $AdS_3$ ? In particular, if the spectrum of conformal dimensions is continuous, then the bulk theory must have a continuum of masses. Perhaps, these are to be identified with some extended long-string like excitations. It would be interesting to see if the mock modularity of the elliptic genus can be understood in terms of such bulk excitations.

The M-theory limit corresponds to the attractor contour for  $n > m$  and  $0 \leq \ell < 2m$  but the converse is not necessarily true, as explained in §11.1. For  $n \leq m$ , we have the possibility that there will be multi-centered configurations. In particular, if  $n \leq m$ , then even if  $\ell < 2m$ , it is possible to have states with non-positive discriminant ( $\Delta = 4mn - \ell^2 \leq 0$ ). Since, the discriminant must be positive for a black hole horizon, such states cannot possibly correspond to single-centered black holes and must be realized in supergravity as multi-centered configurations if there are smooth solutions corresponding to them at all. Hence, if  $\widehat{\psi}_m^F$  is indeed the elliptic genus of a single-centered  $AdS_3$ , then there must be supergravity configurations corresponding to these states for  $n \leq m$  that fit inside a single  $AdS_3$ .

This microscopic prediction is plausible from the supergravity perspective. According to [40], all two-centered black hole solutions where each center is a D4-D2 bound state, are scaled out of a single  $AdS_3$  throat. However, the two-centered solutions that have

one center carrying D6-brane charge and the other carrying anti-D6-brane charge can continue to exist as stable solutions inside a single  $AdS_3$ . Such configurations can give the supergravity realization of the negative discriminant states. Further analysis is required to verify these consequences of our proposed interpretation of  $\widehat{\psi}_m^F$ .

- *Generalization to  $\mathcal{N} = 2$  compactifications:*

One motivation for the present investigation is to abstract some general lessons which could be applicable to the more challenging case of dyons in  $\mathcal{N} = 2$  compactifications on a general Calabi-Yau three-fold. In the  $\mathcal{N} = 4$  case, the mock modularity of  $\psi_m^F$  was a consequence of meromorphy of  $\psi_m$ . As we have seen in §11.2, this meromorphy is in turn a consequence of noncompactness of Taub-NUT factor  $\sigma(\text{TN})$  in the asymptotic SCFT (11.25)<sup>27</sup>. This suggests that the SCFT dual to the near horizon  $AdS_3$  is also noncompact. In the  $\mathcal{N} = 2$  context also, there is no a priori reason to expect that the MSW SCFT will be compact in general everywhere in the moduli space. If this SCFT is noncompact, then our results suggest that mock Jacobi forms would provide the right framework for making the holographic  $SL(2, \mathbb{Z})$  symmetry manifest for the indexed partition function considered, for example, in [44, 39, 81]. In the  $\mathcal{N} = 4$  case, the mock modularity is closely correlated with the wall-crossing phenomenon. The wall-crossing phenomenon in  $\mathcal{N} = 2$  case corresponds to jumps in the Donaldson-Thomas invariants and is known to be much more complicated [43, 70, 68, 80]. It would be interesting though to define the analog of the counting function for the immortal degeneracies in this case. Since these degeneracies do not change under wall-crossing, such a counting function is likely to have an interesting mathematical interpretation. Moreover, it is expected to have nice (mock) modular properties from the considerations of holography discussed above.

- *Mock modularity and indefinite theta series:*

Mock modular forms appear in Zwegers’s work as members of three quite different families of functions:

1. Fourier coefficients of meromorphic Jacobi forms,
2. Appell-Lerch sums,
3. Indefinite theta series.

We have seen that the first two families appear naturally in the context of black hole physics with natural physical interpretations. This suggests that the indefinite theta

---

<sup>27</sup>Note that this is not to be confused with  $\mathbb{R}^3$  factor corresponding to the center of mass motion. It corresponds to the relative motion of the KK-monopole and the D1D5 system [37] and thus would belong to what is called the “entropic” factor [84] for the MSW string in the M-theory frame.

series could also play a role in this physical context. It would be interesting to develop this connection further. For some earlier work in this direction, see [82, 1].

## Acknowledgments

We would like to acknowledge useful conversations with Kathrin Bringmann, Miranda Cheng, Jan de Boer, Rajesh Gopakumar, Jeff Harvey, Jan Manschot, Shiraz Minwalla, Boris Pioline, Jan Troost, Erik Verlinde, Sander Zwegers, and especially Ashoke Sen. We would also like to thank Alexander Weisse for technical help. It is a pleasure to thank the organizers of the 2007 workshop on ‘Black Holes and Modular Forms’ in Paris and of the 2008 workshop on ‘Number Theory and Physics at Crossroads’ in Banff for hospitality where this research was initiated. The work of A. D. is supported in part by the Excellence Chair of the Agence Nationale de la Recherche (ANR). The work of S. M. is presently supported by the ERC Advanced Grant no. 246974, “*Supersymmetry: a window to non-perturbative physics*”.

## A. Appendix: Tables of special mock Jacobi forms

This appendix contains three collections of tables. The first gives the first Fourier coefficients of the weight 2 mock Jacobi forms  $\mathcal{Q}_M$  for all products of two primes less than 50, the second gives coefficients of the primitive mock Jacobi forms  $\Phi_{1,m}^0$  for all known cases having optimal growth, and the third gives coefficients of the weight 1 mock Jacobi forms  $\mathcal{Q}_M$  for all products of three primes less than 150.

### A.1 Table of $\mathcal{Q}_M$ (weight 2 case)

For each  $M$  with  $\mu(M) = 1$  with  $1 < M < 50$  we tabulate the value of  $c_M$  and the Fourier coefficients  $c(\mathcal{F}_M; n, r_{\min})$  for  $n$  ranging from 0 to some limit and all values of  $r_{\min}$ , (defined as the minimal positive representatives prime to  $M$  of their squares modulo  $4M$ ). Here  $c_M$  is the smallest positive rational number such that  $\mathcal{F}_M = -c_M \mathcal{Q}_M$  has integral coefficients, as defined in (9.52). The asterisk before a value of  $r_{\min}$  means that the corresponding row contains at least one negative non-polar coefficient.

$$\mathbf{M} = \mathbf{6} \quad c_M = 12$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12
1	-1	35	130	273	595	1001	1885	2925	4886	7410	11466	16660	24955

$$\mathbf{M} = \mathbf{10} \quad c_M = 6$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13
1	-1	21	63	112	207	306	511	693	1071	1442	2037	2709	3766	4788
3	0	9	35	57	126	154	315	378	625	819	1233	1491	2268	2772

$$\mathbf{M} = \mathbf{14} \quad c_M = 4$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13
1	-1	15	42	65	120	150	255	312	465	575	819	975	1355	1605
3	0	10	30	51	85	120	195	230	360	465	598	765	1065	1235
5	0	3	15	20	51	45	113	105	195	215	345	348	595	615

$$\mathbf{M} = \mathbf{15} \quad c_M = 3$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	-1	14	29	54	83	128	172	230	331	430	537	726	924	1116	1409
2	0	4	24	28	58	52	136	120	224	232	376	368	626	632	920
4	0	10	28	40	88	98	176	208	304	386	552	632	888	1050	1372
7	0	2	9	26	36	50	84	116	155	226	269	340	488	580	712

**M = 21**  $c_M = 2$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	8	15	32	40	56	79	96	135	176	191	248	336	368	434	576
2	0	8	26	28	56	64	112	104	180	184	272	300	416	432	606	636
4	0	2	12	8	32	14	60	40	80	64	148	96	208	166	272	268
5	0	8	17	32	49	64	85	120	150	200	241	288	374	448	534	656
8	0	2	12	20	32	38	76	72	112	138	176	200	304	302	408	472
11	0	0	3	8	11	24	26	32	59	64	74	112	138	144	189	248

**M = 22**  $c_M = 12$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13
1	-5	49	132	176	308	357	572	604	951	1061	1391	1572	2227	2325
3	0	44	103	169	272	301	536	580	771	976	1307	1380	2004	2121
5	0	25	91	116	223	229	427	405	716	700	1071	1121	1608	1633
7	0	13	48	79	127	180	272	272	452	549	632	764	1120	1133
9	0	1	23	24	91	25	180	92	247	205	403	229	715	453

**M = 26**  $c_M = 2$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	9	21	30	51	54	87	96	138	149	207	219	308	315	399	468
3	0	7	21	27	43	51	90	76	132	144	183	201	297	283	393	423
5	0	6	15	23	39	42	69	75	102	128	165	165	243	258	324	371
7	0	3	12	14	33	27	57	51	90	84	132	126	201	183	261	267
9	0	1	6	12	15	21	36	34	57	63	70	93	138	118	162	210
11	0	0	3	2	12	3	26	6	30	24	57	15	86	45	87	81

**M = 33**  $c_M = 6$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	-5	33	55	127	127	160	242	259	347	468	468	534	749	826	903
2	0	21	66	43	142	120	252	174	340	262	526	394	658	580	844
4	0	24	88	88	165	154	286	244	418	376	572	596	869	728	1122
5	0	14	36	47	80	113	113	160	215	259	229	372	460	386	507
7	0	33	44	99	143	167	209	308	365	440	506	541	748	847	915
* 8	0	-4	29	18	62	-4	146	-8	124	76	234	54	336	116	380
10	0	11	50	66	116	110	248	220	292	358	474	435	738	622	892
13	0	0	21	54	77	120	131	153	230	273	317	405	482	548	613
16	0	0	3	22	58	36	94	124	127	146	248	182	350	380	408
19	0	0	0	7	18	7	40	40	58	113	58	113	186	153	120



**M = 34**      $c_M = 3$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	-2	14	36	43	69	82	123	117	183	195	253	252	364	352	481
3	0	13	29	44	77	61	125	116	164	185	244	234	381	334	427
5	0	11	28	39	59	67	106	107	162	157	214	242	321	306	391
7	0	7	26	28	53	54	100	74	149	144	194	177	292	252	378
9	0	5	14	27	41	43	79	72	92	125	165	141	236	234	262
11	0	1	13	12	36	26	57	43	100	69	124	100	196	140	236
13	0	0	5	12	16	18	34	32	54	65	53	85	128	80	139
* 15	0	0	2	1	13	-2	35	0	29	14	59	-2	94	28	72

**M = 35**      $c_M = 1$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	10	14	18	31	44	49	60	76	86	98	134	147	168	200	226
* 2	0	-2	8	4	14	-2	24	8	30	12	56	0	70	22	72	44
3	0	2	6	16	11	18	24	26	43	50	44	58	84	70	93	124
4	0	6	10	16	28	26	46	40	56	70	90	96	128	128	152	176
6	0	6	16	16	30	32	56	52	78	80	104	108	152	150	208	216
8	0	2	10	8	20	12	42	24	52	40	74	48	104	90	122	120
9	0	2	3	8	14	18	17	16	30	42	39	58	69	60	71	100
11	0	2	5	16	22	20	30	42	52	68	67	84	114	118	121	166
13	0	0	5	10	9	18	24	34	33	42	59	52	79	92	99	126
16	0	0	2	0	12	12	16	8	28	18	50	40	56	36	88	56
18	0	0	0	4	6	6	24	8	30	24	32	40	64	46	80	80
23	0	0	0	0	1	2	5	8	7	16	11	8	33	26	19	50

**M = 38**      $c_M = 4$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	-3	21	44	59	103	97	159	149	238	240	322	300	465	445	540
3	0	16	46	62	80	102	166	138	226	231	287	318	449	391	559
5	0	15	43	48	91	75	151	148	194	208	303	269	427	354	512
7	0	14	29	43	82	83	119	117	179	197	256	252	360	358	433
9	0	6	30	36	61	55	124	77	179	165	201	193	360	256	404
11	0	4	16	31	47	45	88	91	108	136	168	155	257	254	300
13	0	0	14	14	40	26	72	45	103	64	169	105	207	137	249
15	0	0	5	11	14	29	43	25	57	75	58	76	160	88	133
* 17	0	0	1	1	20	-7	33	-5	47	21	60	-14	109	15	83

**M = 39**       $c_M = 1$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	6	8	16	22	26	31	42	47	60	72	72	94	110	115	154
2	0	4	14	16	26	20	52	32	64	60	86	76	128	100	148	152
4	0	2	12	6	20	16	36	22	48	32	68	46	96	66	124	86
5	0	6	11	18	22	34	37	44	64	66	71	102	116	120	137	170
7	0	2	3	10	12	12	16	18	28	42	28	40	64	44	57	88
8	0	4	12	12	28	22	44	40	56	62	88	74	120	108	156	148
* 10	0	0	4	0	10	-4	22	4	18	0	36	0	46	16	42	24
11	0	2	7	16	17	24	34	36	47	66	62	76	107	104	117	154
14	0	0	6	8	14	20	28	24	46	40	62	60	86	72	118	116
17	0	0	1	6	10	12	15	24	31	34	35	44	61	72	71	86
20	0	0	0	2	4	6	16	8	16	20	24	30	48	32	44	60
23	0	0	0	0	1	4	3	2	11	12	4	18	19	6	25	32

**M = 46**       $c_M = 12$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13
1	-11	57	161	184	276	299	483	414	667	644	874	897	1219	1023
3	0	69	115	184	299	253	493	437	585	690	838	736	1288	1127
5	0	46	138	147	299	285	414	368	667	607	782	676	1205	1044
7	0	46	123	161	215	230	414	353	560	575	744	790	1043	905
9	0	30	92	138	230	168	421	329	467	444	766	559	1065	766
11	0	23	75	98	167	230	282	276	443	443	558	535	817	765
13	0	5	69	69	166	120	281	143	442	350	465	424	810	516
15	0	0	27	73	119	92	211	188	211	349	376	303	606	487
17	0	0	26	26	92	26	187	95	256	121	325	233	466	236
19	0	0	2	23	25	71	94	48	163	163	96	140	349	188
* 21	0	0	0	1	47	-22	93	-22	70	1	208	-68	324	-21

## A.2 Table of $\Phi_{1,m}^0$ of optimal growth

We give the Fourier coefficients of the primitive weight 1 mock Jacobi form  $\Phi_{1,m}^0$  for the thirteen known values of  $m$  for which it has optimal growth, in the same format as was used in the  $k = 2$  case above. The number  $c$  is the factor relating the true value of  $\Phi_{1,m}^0$  to the form whose coefficients have been tabulated, and which has been normalized to have integral Fourier coefficients with no common factor except for the values  $C(\Delta = 0, r)$ , which are allowed to have a denominator of 2. (This happens for the indices 9, 16, 18, and 25.)

$$\mathbf{m} = \mathbf{2} \quad c = 24$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11
1	-1	45	231	770	2277	5796	13915	30843	65550	132825	260568	494385

$$\mathbf{m} = \mathbf{3} \quad c = 12$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13
1	-1	16	55	144	330	704	1397	2640	4819	8480	14509	24288	39765	63888
2	0	10	44	110	280	572	1200	2244	4180	7348	12772	21330	35288	56760

$$\mathbf{m} = \mathbf{4} \quad c = 8$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	-1	7	21	43	94	168	308	525	882	1407	2255	3468	5306	7931	11766
2	0	8	24	56	112	216	392	672	1128	1840	2912	4536	6936	10416	15456
3	0	3	14	28	69	119	239	393	693	1106	1806	2772	4333	6468	9710

$$\mathbf{m} = \mathbf{5} \quad c = 6$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	4	9	20	35	60	104	164	255	396	590	864	1259	1800	2541	3560
2	0	5	15	26	54	90	156	244	396	590	905	1320	1934	2751	3924	5456
3	0	4	11	24	45	80	135	220	350	540	810	1204	1761	2524	3586	5040
4	0	1	6	10	25	36	76	110	189	280	446	636	970	1360	1980	2750

$$\mathbf{m} = 6 \quad c = 24$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	-5	7	26	33	71	109	185	249	418	582	858	1184	1703	2291	3213
2	0	24	48	96	168	264	432	672	984	1464	2112	2976	4200	5808	7920
3	0	12	36	60	120	180	312	456	720	1020	1524	2124	3036	4140	5760
4	0	12	36	72	120	216	348	528	816	1212	1752	2520	3552	4920	6792
5	0	1	13	14	51	53	127	155	291	382	618	798	1256	1637	2369

$$\mathbf{m} = 7 \quad c = 4$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	2	3	5	10	15	21	34	48	65	94	129	175	237	312	413
2	0	2	6	10	16	24	40	54	84	116	164	222	310	406	552	722
3	0	3	6	11	18	29	45	66	95	137	192	264	361	486	650	862
4	0	2	6	8	18	24	42	58	90	122	180	240	338	448	612	794
5	0	1	3	7	11	18	28	41	63	91	125	177	245	328	441	590
6	0	0	2	2	6	6	16	18	32	40	66	80	126	156	224	286

$$\mathbf{m} = 8 \quad c = 4$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	1	2	4	6	7	14	18	25	35	48	63	87	110	146	190
2	0	2	4	6	10	16	22	32	46	62	86	116	152	202	264	340
3	0	2	5	7	13	18	29	38	58	77	108	141	195	250	333	424
4	0	2	4	8	12	18	28	40	56	80	108	144	196	258	336	440
5	0	1	4	5	11	14	24	32	50	63	94	122	170	215	294	371
6	0	0	2	4	6	10	16	22	32	46	62	86	116	152	202	264
7	0	0	1	1	4	3	9	8	17	20	33	36	60	70	101	124

$$\mathbf{m} = 9 \quad c = 3$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	0	2	2	3	6	8	10	15	20	27	36	46	58	78	98
2	0	2	3	4	8	9	15	20	27	37	51	63	87	109	142	178
3	0	2	3	6	9	12	18	26	36	48	63	84	111	144	183	234
4	0	1	4	5	9	13	21	25	39	50	70	90	122	151	202	251
5	0	2	3	6	8	12	18	26	34	48	65	84	112	146	186	240
6	0	$\frac{1}{2}$	3	3	7	9	15	18	30	36	54	66	93	115	156	192
7	0	0	1	2	4	6	9	12	19	26	33	46	62	78	103	134
8	0	0	1	1	3	1	6	5	10	10	18	19	33	35	52	60

$m = 10$      $c = 8$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-3	3	1	2	5	10	9	15	19	26	31	41	52	76	87	108
2	0	4	8	16	16	28	40	48	72	96	120	160	208	256	328	416
3	0	1	7	7	16	14	27	32	51	57	87	101	144	166	226	269
4	0	8	12	16	28	40	56	80	104	136	184	240	304	392	496	624
5	0	2	6	8	14	18	30	34	54	68	92	114	158	190	252	308
6	0	4	8	16	24	32	48	64	88	124	160	208	272	348	440	560
7	0	0	5	3	11	8	22	23	40	39	69	75	113	128	182	210
8	0	0	4	8	8	16	24	32	44	64	80	104	144	176	232	296
*9	0	0	1	-1	5	2	9	2	13	11	23	18	43	37	60	60

$m = 13$      $c = 2$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	1	1	0	1	1	2	3	3	4	6	7	7	10	12	14
2	0	0	1	2	2	3	4	4	6	8	10	12	16	18	24	28
3	0	1	1	2	3	4	5	7	9	11	13	17	22	26	32	39
4	0	1	2	2	4	4	6	8	11	12	18	20	26	32	40	46
5	0	1	2	3	3	5	7	8	11	15	18	23	29	34	43	53
6	0	1	2	2	4	4	8	8	12	14	19	22	30	34	44	52
7	0	1	1	2	4	5	6	8	11	13	17	22	27	34	41	51
8	0	0	2	2	3	3	6	6	10	12	16	18	26	28	38	44
9	0	0	1	2	2	3	4	6	7	10	13	15	20	25	30	37
10	0	0	1	0	2	2	4	4	6	6	10	10	16	17	24	26
11	0	0	0	1	1	1	2	2	4	5	5	8	10	11	14	18
12	0	0	0	0	1	0	2	0	2	1	4	2	6	4	8	8

$m = 16$       $c = 2$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	0	0	0	1	0	1	2	1	1	2	3	3	4	5	5
2	0	0	1	1	1	1	2	2	3	4	4	5	7	7	9	11
3	0	1	1	1	2	2	3	3	4	5	7	7	9	11	13	15
4	0	1	1	1	2	3	3	4	5	6	8	9	11	14	16	19
5	0	0	1	2	2	2	4	4	6	7	9	10	14	14	19	22
6	0	1	1	2	2	3	4	5	6	8	9	11	14	17	20	24
7	0	1	2	1	3	3	4	5	7	7	10	11	15	17	21	23
8	0	$\frac{1}{2}$	1	2	2	3	4	4	6	8	9	12	14	16	20	24
9	0	0	1	1	2	2	4	4	6	6	9	10	14	14	19	22
10	0	0	1	1	2	2	3	4	5	6	8	9	11	14	17	20
11	0	0	1	1	2	2	3	2	5	5	7	7	11	11	15	16
12	0	0	0	1	1	1	2	3	3	4	5	6	8	9	11	14
13	0	0	0	0	1	0	2	1	3	2	4	4	6	6	9	9
14	0	0	0	0	0	1	1	1	1	2	2	3	4	4	5	7
15	0	0	0	0	1	0	1	0	1	1	2	0	3	1	3	2

$m = 18$       $c = 3$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
*1	-2	0	-1	0	-1	2	1	0	0	1	0	3	1	3	2	4
2	0	3	0	3	3	3	3	6	6	6	9	9	12	15	15	21
*3	0	-1	2	1	1	0	3	1	4	2	5	5	7	4	9	8
4	0	0	3	3	3	6	6	6	12	12	15	18	21	24	30	36
5	0	2	1	1	2	3	3	3	4	6	5	6	9	11	12	13
6	0	3	3	3	6	6	9	12	12	15	18	24	27	33	39	45
*7	0	-1	1	0	3	1	3	2	6	4	8	7	11	8	15	15
8	0	3	3	6	6	6	9	12	12	18	21	24	30	36	39	48
9	0	0	1	1	2	2	3	3	5	5	7	7	10	11	14	15
10	0	0	3	3	3	6	9	9	12	15	18	21	27	30	39	45
11	0	0	2	1	3	0	4	3	5	3	7	5	10	8	14	12
12	0	0	$\frac{3}{2}$	3	3	6	6	6	9	12	15	18	21	24	30	36
*13	0	0	0	-1	1	1	2	0	3	3	4	3	8	6	9	8
14	0	0	0	3	3	3	3	6	6	9	9	9	15	18	18	24
*15	0	0	0	0	2	-1	1	1	3	0	4	1	5	2	5	4
16	0	0	0	0	0	0	3	0	3	3	3	6	6	6	9	12
*17	0	0	0	0	1	0	2	0	2	-1	1	0	3	1	3	0

$$\mathbf{m} = 25 \quad c = 1$$

$r_{\min}$ \backslash $n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	0	0	0	0	0	0	0	0	0	1	0	0	1	1	1
2	0	0	0	1	0	0	1	0	1	1	1	1	1	1	1	2
3	0	0	0	0	1	0	1	1	1	1	1	1	2	2	2	2
4	0	1	1	0	1	1	1	1	1	1	2	2	2	3	3	3
5	0	0	1	1	0	1	1	1	1	2	2	2	3	3	3	4
6	0	0	0	0	1	1	1	1	2	2	2	2	3	3	4	4
7	0	1	0	1	1	1	1	2	2	2	2	3	3	4	4	5
8	0	0	1	1	1	1	1	1	2	2	3	3	4	3	5	5
9	0	1	1	1	1	1	2	2	2	2	3	3	4	4	5	6
10	0	$\frac{1}{2}$	1	0	1	1	2	2	2	2	3	3	4	4	5	5
11	0	0	0	1	1	1	1	1	2	3	2	3	4	4	5	6
12	0	0	1	1	1	1	2	1	2	2	3	3	4	4	5	5
13	0	0	1	0	1	1	1	2	2	2	3	3	3	4	5	5
14	0	0	1	1	1	1	2	1	2	2	3	3	4	3	5	5
15	0	0	0	1	1	1	1	1	2	2	2	2	3	4	4	5
16	0	0	0	0	1	0	1	1	2	1	2	2	3	3	4	4
17	0	0	0	1	0	1	1	1	1	2	2	2	3	3	3	4
18	0	0	0	0	1	0	1	1	1	1	2	1	3	2	3	3
19	0	0	0	0	1	1	1	1	1	1	1	2	2	2	2	3
20	0	0	0	0	$\frac{1}{2}$	0	1	0	1	1	1	1	2	1	3	2
21	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	2
22	0	0	0	0	0	0	1	0	1	0	1	0	1	1	1	1
23	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	1
24	0	0	0	0	0	0	1	0	0	0	1	0	1	0	1	0

### A.3 Table of $\mathcal{Q}_M$ (weight 1 case)

We give the coefficients of  $\mathcal{Q}_M$  for all products of three primes less than 150 for which  $\mathcal{Q}_M$ . We also show the factor  $c_M$  relating  $\mathcal{Q}_M$  to  $\mathcal{F}_M$  and the minimal discriminant of  $\mathcal{F}_M$ , since it is no longer assumed to be  $-1$ . This time an asterisk means simply that the corresponding row contains (non-polar) coefficients of opposite signs, since here there are many examples of values of  $r \pmod{2m}$  for which the scalar factor  $\kappa_r$  that determines the sign of the coefficients  $C(\Delta, r)$  for large  $\Delta$  is negative rather than positive for  $r = r_{\min}$ .

$$\mathbf{M} = 30 \quad c_M = 3 \quad \Delta_{\min} = -1$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	1	1	2	1	3	2	3	3	5	3	6	5	7	7	9
7	0	1	2	2	3	3	4	4	6	5	7	8	9	9	12	12

$$\mathbf{M} = 42 \quad c_M = 2 \quad \Delta_{\min} = -1$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	-1	0	-1	-1	-1	0	-2	-1	-2	-1	-2	-1	-3	-2	-3
5	0	1	1	1	2	1	2	2	2	3	3	2	4	4	4	4
11	0	1	1	2	1	2	2	3	2	3	3	4	4	5	4	6

$$\mathbf{M} = 66 \quad c_M = 12 \quad \Delta_{\min} = -25$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10
1	-5	-5	-15	-60	-125	-313	-620	-1270	-2358	-4394	-7698
5	-1	15	65	175	450	989	2105	4140	7930	14508	25915
7	0	16	55	155	385	852	1816	3597	6880	12645	22627
13	0	5	22	60	155	357	781	1567	3070	5725	10367
19	0	0	-5	-33	-99	-268	-605	-1320	-2623	-5104	-9398

$$\mathbf{M} = 70 \quad c_M = 1 \quad \Delta_{\min} = -1$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	0	-1	-1	0	0	-1	0	-1	-1	-1	-1	-1	0	-1	-1
3	0	0	1	0	1	0	0	1	1	0	1	1	1	0	1	1
9	0	1	0	1	0	1	1	1	0	1	1	1	1	1	1	2
11	0	-1	0	0	0	-1	0	-1	0	0	0	-1	0	-1	0	-1
13	0	1	1	0	1	1	1	1	1	1	1	1	1	2	2	1
23	0	0	1	1	1	1	1	1	1	1	2	1	2	1	2	2

$$\mathbf{M} = 78 \quad c_M = 1 \quad \Delta_{\min} = -1$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	-1	0	0	-1	-1	0	0	-1	0	0	-1	0	-1	-1	-1	-1
5	0	1	0	1	0	1	0	1	1	1	0	1	1	1	1	1
7	0	-1	0	0	0	0	0	-1	0	-1	0	0	0	-1	0	0
11	0	1	1	0	1	0	1	1	1	1	1	1	1	1	1	1
17	0	1	0	1	1	1	1	1	0	1	1	1	1	2	1	2
23	0	0	1	1	1	0	1	1	1	1	1	1	2	1	1	1



$$\mathbf{M} = \mathbf{102} \quad c_M = 6 \quad \Delta_{\min} = -25$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11
1	-3	-1	-8	-14	-31	-56	-102	-173	-293	-461	-732	-1129
5	-1	7	16	41	76	153	262	454	745	1215	1886	2941
7	0	3	12	21	45	76	146	236	408	636	1024	1555
11	0	3	13	26	59	103	197	327	560	890	1427	2187
13	0	6	15	31	68	125	229	389	657	1055	1688	2603
19	0	2	7	16	35	61	120	205	349	574	924	1433
*25	0	0	1	-1	-4	-11	-16	-42	-61	-115	-178	-303
31	0	0	0	-3	-12	-27	-61	-113	-209	-358	-605	-972

$$\mathbf{M} = \mathbf{105} \quad c_M = 1 \quad \Delta_{\min} = -4$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	0	1	1	1	2	1	2	3	3	3	4	4	5	6	7	7
2	-1	-1	-1	-1	-1	-1	-2	-2	-2	-3	-3	-3	-4	-4	-5	-6
4	0	1	0	1	1	1	1	1	2	2	2	2	3	4	3	4
8	0	0	1	1	0	1	2	1	2	2	2	3	3	3	4	5
11	0	0	1	0	1	1	1	1	2	1	3	2	3	2	4	4
13	0	1	0	1	1	1	1	2	1	2	2	3	3	4	3	5
16	0	1	1	1	2	2	2	3	3	4	4	5	6	7	8	8
17	0	-1	0	-1	0	-1	0	-2	0	-1	-1	-2	-1	-3	-1	-3
19	0	1	1	2	1	2	2	3	3	4	4	5	5	7	7	9
23	0	0	1	0	1	1	2	1	2	1	3	3	4	3	5	4
32	0	0	0	0	0	1	0	0	1	1	1	1	1	1	2	2
34	0	0	0	1	1	1	1	2	2	2	2	3	4	4	4	6

$$\mathbf{M} = \mathbf{110} \quad c_M = 3 \quad \Delta_{\min} = -9$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	-2	-1	-2	-3	-3	-5	-5	-9	-9	-13	-15	-22	-25	-32	-37
3	-1	0	2	1	4	4	6	7	11	14	18	21	28	34	44
7	0	1	1	3	2	3	4	5	6	9	9	13	16	20	21
9	0	2	3	2	5	5	9	10	13	15	22	24	33	39	50
13	0	1	2	4	6	7	10	14	18	23	31	37	49	58	74
17	0	2	3	4	5	8	10	12	17	22	27	34	43	52	65
19	0	1	2	2	2	3	5	5	6	9	10	13	16	19	24
23	0	0	2	1	4	4	6	8	13	13	20	23	31	37	49
*29	0	0	1	0	1	0	1	-2	0	-3	-2	-4	-2	-9	-6
*39	0	0	0	0	1	-1	-1	-2	-3	-6	-5	-10	-11	-16	-20

$$\mathbf{M} = \mathbf{114} \quad c_M = 4 \quad \Delta_{\min} = -25$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13
1	-1	4	9	18	37	62	110	181	291	449	695	1034	1537	2235
5	-1	0	4	3	11	16	32	41	83	113	188	266	413	574
7	0	6	12	30	53	99	166	282	444	706	1067	1622	2387	3498
11	0	3	11	21	42	74	133	216	351	547	849	1271	1897	2757
13	0	3	8	16	37	59	111	178	295	457	718	1062	1606	2327
17	0	2	9	19	38	70	127	205	345	538	840	1267	1904	2778
23	0	0	3	5	15	24	48	75	135	204	333	496	765	1110
29	0	0	-1	-2	-6	-10	-20	-37	-64	-100	-165	-257	-388	-583
35	0	0	0	-3	-7	-18	-32	-67	-111	-194	-310	-501	-767	-1184

$$\mathbf{M} = \mathbf{130} \quad c_M = 2 \quad \Delta_{\min} = -9$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	-1	0	-1	0	-1	-1	-2	-2	-3	-3	-5	-5	-7	-7	-12
3	-1	-1	-1	-2	-1	-3	-3	-5	-5	-7	-7	-11	-11	-16	-17
7	0	1	2	2	3	3	6	6	8	9	13	14	19	21	27
9	0	1	2	2	4	4	6	8	10	12	16	18	24	28	35
11	0	1	1	2	2	3	4	4	6	8	9	10	14	17	19
17	0	1	1	2	2	3	4	5	5	8	9	12	14	17	19
19	0	1	2	2	4	5	6	8	11	13	17	20	26	31	38
21	0	1	2	3	3	5	6	8	10	13	15	20	23	29	35
27	0	0	1	0	1	1	2	0	3	1	3	3	5	3	7
29	0	0	1	1	2	2	4	4	6	6	10	11	15	16	22
37	0	0	0	0	0	-1	0	-1	-2	-2	-2	-4	-4	-6	-7
47	0	0	0	0	0	-1	0	-2	-2	-3	-3	-6	-5	-9	-10

$$\mathbf{M} = \mathbf{138} \quad c_M = 2 \quad \Delta_{\min} = -49$$

$r_{\min} \backslash n$	0	1	2	3	4	5	6	7	8	9	10	11
1	-5	5	20	57	141	323	658	1308	2449	4450	7786	13373
5	-2	9	27	56	125	235	451	812	1442	2463	4182	6884
7	-1	13	44	111	265	555	1138	2170	4032	7183	12541	21237
11	0	14	46	115	261	546	1091	2055	3787	6702	11610	19591
13	0	9	29	68	151	308	612	1139	2080	3675	6323	10653
17	0	6	28	68	174	354	753	1419	2678	4775	8407	14255
*19	0	2	6	-2	-12	-69	-157	-397	-784	-1581	-2884	-5240
25	0	0	-3	-23	-69	-192	-427	-926	-1817	-3473	-6310	-11189
31	0	0	-1	-11	-33	-100	-220	-491	-973	-1890	-3456	-6203
*37	0	0	0	-1	-1	-5	6	8	49	110	263	490
43	0	0	0	0	2	13	47	128	309	677	1369	2653

## B. Bibliography

- [1] Sergei Alexandrov, Jan Manschot, and Boris Pioline, *D3-instantons, Mock Theta Series and Twistors*, (2012), [arxiv:1207.1109](#).
- [2] Murad Alim, Babak Haghighat, Michael Hecht, Albrecht Klemm, Marco Rauch, and Thomas Wotschke, *Wall-crossing holomorphic anomaly and mock modularity of multiple M5-branes*, (2010), [arxiv:1012.1608](#).
- [3] O. Alvarez, T. P. Killingback, M. L. Mangano, and P. Windey, *String theory and loop space index theorems*, Commun. Math. Phys. **111** (1987) 1.
- [4] S. K. Ashok and J. Troost, *A Twisted Non-compact Elliptic Genus*, JHEP **1103** (2011) 067, [arxiv:1101.10593](#).
- [5] Shamik Banerjee and Ashoke Sen, *Duality orbits, dyon spectrum and gauge theory limit of heterotic string theory on  $T^6$* , JHEP **0803** (2008) 022, [arxiv:0712.0043](#).
- [6] ———, *S-duality action on discrete T-duality invariants*, JHEP **0804** (2008) 012, [arxiv:0801.0149](#).
- [7] S. Banerjee, A. Sen, and Y. K. Srivastava, *Generalities of quarter BPS dyon partition function and dyons of torsion two*, JHEP **0805** (2008) 101, [arxiv:0802.0544](#).
- [8] ———, *Partition functions of torsion  $> 1$  dyons in heterotic string theory on  $T^6$* , JHEP **0805** (2008) 098, [arxiv:0802.1556](#).
- [9] ———, *Genus two surface and quarter BPS dyons: The contour prescription*, JHEP **0903** (2009) 151, [arxiv:0808.1746](#).
- [10] N. Banerjee, I. Mandal and A. Sen, *Black hole hair removal*, JHEP **0907** (2009) 091, [arxiv:0901.0359](#).
- [11] B. Bates and F. Denef, *Exact solutions for supersymmetric stationary black hole composites*, JHEP **1111** (2011) 127, [hep-th/0304094](#).
- [12] J. C. Breckenridge, Robert C. Myers, A. W. Peet, and C. Vafa, *D-branes and spinning black holes*, Phys.Lett. **B391** (1997) 93–98, [hep-th/9602065](#).
- [13] K. Bringmann and K. Ono, *Coefficients of harmonic Maass forms*, Proceedings of the 2008 University of Florida Conference on Partitions, q-series, and modular forms.
- [14] K. Bringmann and O. Richter, *Zagier-type dualities and lifting maps for harmonic Maass-Jacobi forms*, Adv. Math. **225** no. **4**, (2010), 2298–2315.
- [15] K. Bringmann and J. Manschot, *From sheaves on  $P^2$  to a generalization of the Rademacher expansion*, [arXiv:1006.0915](#).

- [16] K. Bringmann and K. Mahlburg, *An extension of the Hardy-Ramanujan Circle Method and applications to partitions without sequences*, American Journal of Math 133 (2011), pages 1151-1178.
- [17] J. H. Bruinier, G. van der Geer, G. Harder and D. Zagier, *The 1-2-3 of Modular Forms*, Universitext, Springer-Verlag, Berlin–Heidelberg (2008).
- [18] G. L. Cardoso, J. R. David, B. de Wit, and S. Mahapatra, *The mixed black hole partition function for the STU model*, JHEP **12** (2008), 086, [arxiv:0810.1233](#).
- [19] Alejandra Castro and Sameer Murthy, *Corrections to the statistical entropy of five dimensional black holes*, JHEP **0906** (2009), 024, [arxiv:0807.0237](#).
- [20] Miranda C.N. Cheng, *K3 surfaces,  $N = 4$  Dyons, and the Mathieu group  $M_{24}$* , (2010), [arxiv:1005.5415](#).
- [21] Miranda C. N. Cheng, John F. R. Duncan and Jeffrey A. Harvey, *Umbral Moonshine*, [arXiv:1204.2779](#) .
- [22] Miranda C. N. Cheng and Erik Verlinde, *Dying dyons don't count*, JHEP **0709** (2007) 070, [arxiv:0706.2363](#).
- [23] Young Ju Choie and Subong Lim, *The heat operator and mock Jacobi forms*, The Ramanujan Journal Volume 22, Number 2 (2010), 209-219.
- [24] Atish Dabholkar, *Exact counting of black hole microstates*, Phys. Rev. Lett. **94** (2005), 241301, [hep-th/0409148](#).
- [25] Atish Dabholkar, Frederik Denef, Gregory W. Moore, and Boris Pioline, *Exact and asymptotic degeneracies of small black holes*, JHEP **08** (2005), 021, [hep-th/0502157](#).
- [26] ———, *Precision counting of small black holes*, JHEP **0510** (2005) 096, [hep-th/0507014](#).
- [27] Atish Dabholkar, Davide Gaiotto, and Suresh Nampuri, *Comments on the spectrum of CHL dyons*, JHEP **01** (2008), 023, [hep-th/0702150](#).
- [28] Atish Dabholkar, Gary W. Gibbons, Jeffrey A. Harvey, and Fernando Ruiz Ruiz, *Superstrings and solitons*, Nucl. Phys. **B340** (1990), 33–55.
- [29] Atish Dabholkar, Joao Gomes, and Sameer Murthy, *Counting all dyons in  $N = 4$  string theory*, (2008), [arxiv:0803.2692](#).
- [30] ———, *Quantum black holes, localization and the topological string*, JHEP **1106** (2011) 019 [arxiv:1012.0265](#).
- [31] ———, *Localization & Exact Holography*, [arxiv:1111.1161](#).

- [32] Atish Dabholkar, Joao Gomes, Sameer Murthy, and Ashoke Sen, *Supersymmetric index from black hole entropy*, (2010), [arxiv:1009.3226](#).
- [33] A. Dabholkar, M. Guica, S. Murthy and S. Nampuri, *No entropy enigmas for  $N = 4$  dyons*, JHEP **1006** (2010) 007 [arxiv:0903.2481](#).
- [34] Atish Dabholkar and Jeffrey A. Harvey, *Nonrenormalization of the superstring tension*, Phys. Rev. Lett. **63** (1989), 478.
- [35] Atish Dabholkar, Renata Kallosh, and Alexander Maloney, *A stringy cloak for a classical singularity*, JHEP **12** (2004), 059, [hep-th/0410076](#).
- [36] Atish Dabholkar and Suresh Nampuri, *Spectrum of dyons and black Holes in CHL orbifolds using Borchers lift*, JHEP **11** (2007), 077, [hep-th/0603066](#).
- [37] Justin R. David and Ashoke Sen, *CHL dyons and statistical entropy function from D1-D5 system*, JHEP **11** (2006), 072, [hep-th/0605210](#).
- [38] L. J. Dixon, J. A. Harvey, C. Vafa and E. Witten, *Strings on Orbifolds*, Nucl. Phys. B **261** (1985) 678.
- [39] Jan de Boer, Miranda C. N. Cheng, Robbert Dijkgraaf, Jan Manschot, and Erik Verlinde, *A Farey tail for attractor black holes*, JHEP **11** (2006), 024, [hep-th/0608059](#).
- [40] J. de Boer, F. Denef, S. El-Showk, I. Messamah and D. Van den Bleeken, *Black hole bound states in  $AdS_3 \times S^2$* , JHEP **0811** (2008) 050, [arXiv:0802.2257](#).
- [41] F. Denef, *Supergravity flows and D-brane stability*, JHEP **0008** (2000) 050, [hep-th/0005049](#).
- [42] F. Denef, *Quantum quivers and Hall/hole halos*, JHEP **0210** (2002) 023, [hep-th/0206072](#).
- [43] Frederik Denef and Gregory W. Moore, *Split states, entropy enigmas, holes and halos*, (2007), [hep-th/0702146](#).
- [44] Robbert Dijkgraaf, Juan Martin Maldacena, Gregory Winthrop Moore, and Erik P. Verlinde, *A black hole Farey tail*, (2000), [hep-th/0005003](#).
- [45] Robbert Dijkgraaf, Gregory W. Moore, Erik P. Verlinde, and Herman L. Verlinde, *Elliptic genera of symmetric products and second quantized strings*, Commun. Math. Phys. **185** (1997), 197–209, [hep-th/9608096](#).
- [46] Robbert Dijkgraaf, Erik P. Verlinde, and Herman L. Verlinde, *Counting dyons in  $N = 4$  string theory*, Nucl. Phys. **B484** (1997), 543–561, [hep-th/9607026](#).
- [47] Tohru Eguchi, Hiroshi Ooguri, and Yuji Tachikawa, *Notes on the  $K3$  surface and the Mathieu group  $M_{24}$* , (2010), [arxiv:1004.0956](#).

- [48] Tohru Eguchi, Hiroshi Ooguri, S-K. Yang and Anne Taormina, *Superconformal algebras and string compactification on manifolds with  $SU(N)$  Holonomy*, Nucl. Phys. **B315** (1989), 193.
- [49] T. Eguchi and Y. Sugawara,  *$SL(2, R)/U(1)$  supercoset and elliptic genera of noncompact Calabi-Yau manifolds*, JHEP **0405** (2004) 014. [hep-th/0403193](#)
- [50] Tohru Eguchi, Yuji Sugawara and Anne Taormina, *Liouville field, modular forms and elliptic genera*, JHEP **0703** (2007), 119. [hep-th/0611338](#)
- [51] ———, *On the unitary representation of  $N = 2$  and  $N = 4$  superconformal algebras*, Phys.Lett. **B210** (1988),125.
- [52] M. Eichler and D. Zagier, *The Theory of Jacobi Forms*, Birkhäuser, 1985.
- [53] Sergio Ferrara, Renata Kallosh, and Andrew Strominger,  *$N = 2$  extremal black holes*, Phys. Rev. **D52** (1995), 5412–5416, [hep-th/9508072](#).
- [54] Davide Gaiotto, *Re-recounting dyons in  $N = 4$  string theory*, (2005), [hep-th/0506249](#).
- [55] Davide Gaiotto, Andrew Strominger, and Xi Yin, *New connections between 4D and 5D black holes*, JHEP **02** (2006), 024, [hep-th/0503217](#).
- [56] ———, *The M5-brane elliptic genus: Modularity and BPS states*, JHEP **08** (2007), 070, [hep-th/0607010](#).
- [57] Jerome P. Gauntlett, Nakwoo Kim, Jaemo Park, and Piljin Yi, *Monopole dynamics and BPS dyons in  $N = 2$  super-Yang-Mills theories*, Phys. Rev. **D61** (2000), 125012, [hep-th/9912082](#).
- [58] Paul H. Ginsparg, *Applied conformal field theory*, (1988), [hep-th/9108028](#).
- [59] L. Goettsche, *The Betti numbers of the Hilbert scheme of points on a smooth projective surface*, Math. Ann. **286** (1990), 193.
- [60] V. Gritsenko, “Elliptic genus of Calabi-Yau manifolds and Jacobi and Siegel modular forms,” [math/9906190](#)
- [61] V. A. Gritsenko and V. V. Nikulin, *Automorphic forms and Lorentzian Kac–Moody Algebras. Part I*, (1996) [alg-geom/9610022](#).
- [62] V. A. Gritsenko and V. V. Nikulin, *Automorphic forms and Lorentzian Kac–Moody Algebras. Part II*, (1996) [alg-geom/9611028](#).
- [63] V. A. Gritsenko, N.-P. Skoruppa, D. Zagier, *Theta blocks I: Elementary theory and examples*, in preparation.
- [64] D. R. Hickerson, *On the seventh order mock theta functions*, Inv. Math. **94**, (1988), 661–677.

- [65] F. Hirzebruch and D. Zagier, *Intersection numbers on curves on Hilbert modular surfaces and modular forms of Nebentypus*, *Inv. Math.* **36** (1976), 57.
- [66] C. M. Hull and P. K. Townsend, *Unity of superstring dualities*, *Nucl. Phys.* **B438** (1995), 109–137, [hep-th/9410167](#).
- [67] D. P. Jatkar, A. Sen and Y. K. Srivastava, *Black hole hair removal: non-linear analysis*, *JHEP* **1002** (2010) 038, [arXiv:0907.0593](#).
- [68] D. Joyce, Y. Song, *A Theory of Generalized Donaldson-Thomas Invariants*, *Memoirs of the AMS* 217 (2012), [arXiv:0810.5645](#).
- [69] Toshiya Kawai, Yasuhiko Yamada, and Sung-Kil Yang, *Elliptic genera and  $N = 2$  superconformal field theory*, *Nucl. Phys.* **B414** (1994), 191–212, [hep-th/9306096](#).
- [70] M. Kontsevich and Y. Soibelman, *Stability structures, motivic Donaldson-Thomas invariants and cluster transformations*, [arXiv:0811.2435](#).
- [71] Per Kraus and Finn Larsen, *Microscopic black hole entropy in theories with higher derivatives*, *JHEP* **09** (2005), 034, [hep-th/0506176](#).
- [72] ———, *Partition functions and elliptic genera from supergravity*, (2006), [hep-th/0607138](#).
- [73] Gabriel Lopes Cardoso, Bernard de Wit, and Thomas Mohaupt, *Corrections to macroscopic supersymmetric black-hole entropy*, *Phys. Lett.* **B451** (1999), 309–316, [hep-th/0412287](#).
- [74] ———, *Area law corrections from state counting and supergravity*, *Class. Quant. Grav.* **17** (2000), 1007–1015, [hep-th/9910179](#).
- [75] ———, *Deviations from the area law for supersymmetric black holes*, *Fortsch. Phys.* **48** (2000), 49–64, [hep-th/9904005](#).
- [76] ———, *Macroscopic entropy formulae and non-holomorphic corrections for supersymmetric black holes*, *Nucl. Phys.* **B567** (2000), 87–110, [hep-th/9906094](#).
- [77] Gabriel Lopes Cardoso, B. de Wit, J. Kappeli, and T. Mohaupt, *Asymptotic degeneracy of dyonic  $N = 4$  string states and black hole entropy*, *JHEP* **12** (2004), 075.
- [78] J. M. Maldacena and A. Strominger,  *$AdS(3)$  black holes and a stringy exclusion principle*, *JHEP* **9812** (1998) 005 [hep-th/9804085](#).
- [79] Juan M. Maldacena, Andrew Strominger, and Edward Witten, *Black hole entropy in M-theory*, *JHEP* **12** (1997), 002, [hep-th/9711053](#).
- [80] J. Manschot, B. Pioline and A. Sen, *Wall Crossing from Boltzmann Black Hole Halos*, *JHEP* **1107** (2011) 059 [arXiv:1011.1258](#).



- [81] Jan Manschot and Gregory W. Moore, *A Modern Farey Tail*, (2007), [arxiv:0712.0573](#).
- [82] Jan Manschot, *Stability and duality in  $N=2$  supergravity*, Commun.Math.Phys., **299**, 651-676 (2010), [arxiv:0906.1767](#).
- [83] J. A. Minahan, D. Nemeschansky, C. Vafa, and N. P. Warner, *E-strings and  $N = 4$  topological Yang-Mills theories*, Nucl. Phys. **B527** (1998), 581–623, [hep-th/9802168](#).
- [84] Ruben Minasian, Gregory W. Moore, and Dimitrios Tsimpis, *Calabi-Yau black holes and  $(0,4)$  sigma models*, Commun. Math. Phys. **209** (2000), 325–352, [hep-th/9904217](#).
- [85] Gregory W. Moore, *Les Houches lectures on strings and arithmetic*, (2004), [hep-th/0401049](#).
- [86] Sameer Murthy and Boris Pioline, *A Farey tale for  $N = 4$  dyons*, JHEP **0909** (2009), 022, [arxiv:0904.4253](#).
- [87] Serge Ochanine, *Sur les genres multiplicatifs definis par des integrales elliptiques*, Topology **26** (1987), 143.
- [88] H. Ooguri, *Superconformal symmetry and geometry of Ricci flat Kähler manifolds*, Int. J. Mod. Phys. **A4** (1989), 4303–4324.
- [89] C. N. Pope, *Axial vector anomalies and the index theorem in charged Schwarzschild and Taub - NUT spaces*, Nucl. Phys. **B141** (1978), 432.
- [90] Ashoke Sen, *Dyon-monopole bound states, selfdual harmonic forms on the multi-monopole moduli space, and  $Sl(2, \mathbb{Z})$  invariance in string theory*, Phys. Lett. **B329** (1994), 217–221, [hep-th/9402032](#).
- [91] ———, *Strong-weak coupling duality in four-dimensional string theory*, Int. J. Mod. Phys. **A9** (1994), 3707–3750, [hep-th/9402002](#).
- [92] ———, *Extremal black holes and elementary string states*, Mod. Phys. Lett. **A10** (1995), 2081–2094, [hep-th/9504147](#).
- [93] ———, *Black holes and the spectrum of half-BPS states in  $N = 4$  supersymmetric string theory*, (2005), [hep-th/0504005](#).
- [94] ———, *Walls of marginal stability and dyon spectrum in  $N = 4$  supersymmetric string theories*, JHEP **05** (2007), 039, [hep-th/0702141](#).
- [95] ———, *Entropy Function and AdS(2)/CFT(1) Correspondence*, JHEP **11** (2008), 075, [arxiv:0805.0095](#).
- [96] ———, *Quantum Entropy Function from AdS(2)/CFT(1) Correspondence*, (2008), [arxiv:0809.3304](#).

- [97] ———, *Negative discriminant states in  $N = 4$  supersymmetric string theories*, (2011), [arxiv:1104.1498](#).
- [98] David Shih, Andrew Strominger, and Xi Yin, *Recounting dyons in  $N = 4$  string theory*, JHEP **10** (2006), 087, [hep-th/0505094](#).
- [99] David Shih and Xi Yin, *Exact black hole degeneracies and the topological string*, JHEP **04** (2006), 034, [hep-th/0508174](#).
- [100] N.-P. Skoruppa, *Über den Zusammenhang zwischen Jacobi-Formen und Modulformen halbganzen Gewichts*, Dissertation, Universität Bonn (1984).
- [101] ———, *Developments in the theory of Jacobi forms*. In Automorphic functions and their applications Khabarovsk, 1988, Acad. Sci. USSR Inst. Appl. Math., Khabarovsk 1990, 167–185
- [102] ———, *Binary quadratic forms and the Fourier coefficients of elliptic and Jacobi modular forms*. J. reine Angew. Math. 411 (1990), 66–95
- [103] N.-P. Skoruppa, D. Zagier, *A trace formula for Jacobi forms* J. reine Angew. Math. 393 (1989) 168-198.
- [104] ———, *Jacobi forms and a certain space of modular forms*. Inventiones mathematicae 94, 113–146 (1988).
- [105] Andrew Strominger, *Macroscopic entropy of  $N = 2$  extremal black holes*, Phys. Lett. **B383** (1996), 39–43, [hep-th/9602111](#).
- [106] ———, *AdS(2) quantum gravity and string theory*, JHEP **01** (1999), 007, [hep-th/9809027](#).
- [107] Andrew Strominger and Cumrun Vafa, *Microscopic origin of the Bekenstein-Hawking entropy*, Phys. Lett. **B379** (1996), 99–104, [hep-th/9601029](#).
- [108] Jan Troost, *The non-compact elliptic genus: mock or modular*, JHEP **1006** (2010), 104, [arxiv:1004.3649](#).
- [109] Cumrun Vafa and Edward Witten, *A strong coupling test of S-duality*, Nucl. Phys. **B431** (1994), 3–77, [hep-th/9408074](#).
- [110] Katrin Wendland, Ph.D. Thesis, Bonn University (2000).
- [111] Edward Witten, *Dyons of charge  $e\theta/2\pi$* , Phys. Lett. B **86** (1979) 283.
- [112] ———, *Elliptic genera and quantum field theory*, Commun. Math. Phys. **109** (1987), 525.
- [113] ———, *String theory dynamics in various dimensions*, Nucl. Phys. **B443** (1995), 85–126, [hep-th/9503124](#).

- [114] Edward Witten and David I. Olive, *Supersymmetry algebras that include topological charges*, Phys. Lett. **B78** (1978), 97.
- [115] D. Zagier, *Nombres de classes et formes modulaires de poids 3/2*, C. R. Acad. Sc. Paris **281** (1975), 883.
- [116] ———, *The Rankin-Selberg method for automorphic functions which are not of rapid decay*, J. Fac. Sci. Tokyo 28 (1982) 415-438.
- [117] ———, *Traces of singular moduli*, in “Motives, Polylogarithms and Hodge Theory” (eds. F. Bogomolov, L. Katzarkov), Lecture Series **3**, International Press, Somerville (2002), 209–244.
- [118] ———, *Ramanujan’s mock theta functions and their applications [d’après Zwegers and Bringmann-Ono]*, Séminaire Bourbaki, 60ème année, 2006-2007, **986** (2007).
- [119] S. P. Zwegers, *Mock theta functions*, Thesis, Utrecht (2002), <http://igitur-archive.library.uu.nl/dissertations/2003-0127-094324/inhoud.htm>.